

Archivos

de medicina del deporte

Órgano de expresión de la Sociedad Española de Medicina del Deporte

ISSN: 0212-8799

190

Volume 36(2)
March - April 2019



ORIGINALS

Relationships among motor coordination, body mass index and physical activity in adolescents with different weight status

Time limit at peak speed without prior warm-up: Effects on test duration, heart rate and rating of perceived exertion

Acute effects of resistance and concurrent exercise on the lipid profile of postmenopausal women


Sweating and core temperature in athletes training in continuous and intermittent sports in tropical climate

The effect of tapering and *Nigella sativa* on the histological structure of the lung after increasing interval exercise training

REVIEWS

Androgens from physiology, through pharmacy and pharmacology to the status of lifestyle drugs - are we going in the right direction?

Sport classification regulations for athletes with differences in sexual development (DSD)



NUEVO ANALIZADOR PORTÁTIL DE LACTATO

Lactate Scout 4



150 € de Descuento *

Incluye Analizador Lactate 4 Solo

o
Analizador Lactate 4 Start +
y 24 tiras reactivas

* Con la entrega de un analizador en uso de Lactate PRO2, Lactate Plus o Lactate Scout +



- Volumen de muestra: 0.2 µl.
- Resultados en 10 segundos
- **NO** precisa calibración previa
- Conexión PC vía Bluetooth
- Software de análisis Lactate Scout Assistant (en presentaciones Start y Maletín)
- Compatible con las tiras reactivas actuales.

Nuevas Características

- Memoria de hasta 500 resultados
- Nueva pantalla para facilitar la visualización
- Diseño más pequeño, más ligero, más robusto
- **Gran Conectividad:**
 - Conexión a monitores de ritmo cardíaco compatibles
 - **App Android** disponible próximamente



Oferta vigente desde el 29 de Abril
al 30 de Junio del 2019

Más información en: www.biolaster.com



Sociedad Española de Medicina del Deporte

Junta de Gobierno

Presidente:

Pedro Manonelles Marqueta

Vicepresidente:

Carlos de Teresa Galván

Secretario General:

Luis Franco Bonafonte

Tesorero:

Javier Pérez Ansón

Vocales:

Miguel E. Del Valle Soto

José Fernando Jiménez Díaz

Juan N. García-Nieto Portabella

Teresa Gaztañaga Aurrekoetxea

José Naranjo Orellana

Edita

Sociedad Española de Medicina del Deporte

Iturrana, 43 bis.

31007 Pamplona. (España)

Tel. 948 267 706 - Fax: 948 171 431

femede@femede.es

www.femede.es

Correspondencia:

Ap. de correos 1207

31080 Pamplona (España)

Publicidad

ESMON PUBLICIDAD

Tel. 93 2159034

Publicación bimestral

Un volumen por año

Depósito Legal

Pamplona. NA 123. 1984

ISSN

0212-8799

Soporte válido

Ref. SVR 389

Indexada en: EMBASE/Excerpta Medica, Índice Médico Español, Sport Information Resource Centre (SIRC), Índice Bibliográfico Español de Ciencias de la Salud (IBECS), Índice SJR (SCImago Journal Rank), y SCOPUS



La Revista Archivos de Medicina del Deporte ha obtenido el Sello de Calidad en la V Convocatoria de evaluación de la calidad editorial y científica de las revistas científicas españolas, de la Fundación Española para la Ciencia y la Tecnología (FECYT).

La dirección de la revista no acepta responsabilidades derivadas de las opiniones o juicios de valor de los trabajos publicados, la cual recaerá exclusivamente sobre sus autores.

Esta publicación no puede ser reproducida total o parcialmente por ningún medio sin la autorización por escrito de los autores.

Cualquier forma de reproducción, distribución, comunicación pública o transformación de esta obra sólo puede ser realizada con la autorización de sus titulares, salvo excepción prevista por la ley.

Dirijase a CEDRO (Centro Español de Derechos Reprográficos, www.cedro.org) si necesita fotocopiar o escanear algún fragmento de esta obra.

Archivos de medicina del deporte

Revista de la Sociedad Española de Medicina del Deporte

Afiliada a la Federación Internacional de Medicina del Deporte, Sociedad Europea de Medicina del Deporte y Grupo Latino y Mediterráneo de Medicina del Deporte

Director

Pedro Manonelles Marqueta

Editor

Miguel E. Del Valle Soto

Administración

M^{ra} Ángeles Artázcoz Bárcena

Adjunto a dirección

Oriol Abellán Aynés

Comité Editorial

Norbert Bachl. Centre for Sports Science and University Sports of the University of Vienna. Austria. **Ramón Balius Matas.** Consell Català de l'Esport. Generalitat de Catalunya. España. **Araceli Boraita.** Servicio de Cardiología. Centro de Medicina del Deporte. Consejo Superior de deportes. España. **Mats Borjesson.** University of Gothenburg. Suecia. **Josep Brugada Terradellas.** Hospital Clinic. Universidad de Barcelona. España. **Nicolas Christodoulou.** President of the UEMS MJC on Sports Medicine. Chipre. **Demitri Constantinou.** University of the Witwatersrand. Johannesburg. Sudáfrica. **Jesús Dapena.** Indiana University. Estados Unidos. **Franchek Drobnic Martínez.** Servicios Médicos FC Barcelona. CAR Sant Cugat del Vallés. España. **Tomás Fernández Jaén.** Servicio Medicina y Traumatología del Deporte. Clínica Centro. España. **Walter Frontera.** Universidad de Vanderbilt. Past President FIMS. Estados Unidos. **Pedro Guillén García.** Servicio Traumatología del Deporte. Clínica Centro. España. **Dusan Hamar.** Research Institute of Sports. Eslovaquia. **José A. Hernández Hermoso.** Servicio COT. Hospital Universitario Germans Trias i Pujol. España. **Pilar Hernández Sánchez.** Universidad Católica San Antonio. Murcia. España. **Markku Jarvinen.** Institute of Medical Technology and Medical School. University of Tampere. Finlandia. **Anna Jegier.** Medical University of Lodz. Polonia. **Peter Jenoure.** ARS Ortopédica, ARS Médica Clínica, Gravesano. Suiza. **José A. López Calbet.** Universidad de Las Palmas de Gran Canaria. España. **Javier López Román.** Universidad Católica San Antonio. Murcia. España. **Alejandro Lucía Mulas.** Universidad Europea de Madrid. España. **Emilio Luengo Fernández.** Servicio de Cardiología. Hospital General de la Defensa. España. **Nicola Maffully.** Universidad de Salerno. Salerno (Italia). **Pablo Jorge Marcos Pardo.** Universidad Católica San Antonio. Murcia. España. **Alejandro Martínez Rodríguez.** Universidad de Alicante. España. **Estrella Núñez Delicado.** Universidad Católica San Antonio. Murcia. España. **Sakari Orava.** Hospital Universitario. Universidad de Turku. Finlandia. **Eduardo Ortega Rincón.** Universidad de Extremadura. España. **Nieves Palacios Gil-Antuñano.** Centro de Medicina del Deporte. Consejo Superior de Deportes. España. **Antonio Pelliccia.** Institute of Sport Medicine and Science. Italia. **José Peña Amaro.** Facultad de Medicina y Enfermería. Universidad de Córdoba. España. **Fabio Pigozzi.** University of Rome Foro Italico, President FIMS. Italia. **Yannis Pitsiladis.** Centre of Sports Medicine. University of Brighton. Inglaterra. **Per Renström.** Stockholm Center for Sports Trauma Research, Karolinska Institutet. Suecia. **Juan Ribas Serna.** Universidad de Sevilla. España. **Peter H. Schober.** Medical University Graz. Austria. **Jordi Segura Noguera.** Laboratorio Antidopaje IMIM. Presidente Asociación Mundial de Científicos Antidopajes (WAADS). España. **Giulio Sergio Roi.** Education & Research Department Isokinetic Medical Group. Italia. **Luis Serratosa Fernández.** Servicios Médicos Sanitas Real Madrid CF. Madrid. España. **Nicolás Terrados Cepeda.** Unidad Regional de Medicina Deportiva del Principado de Asturias. Universidad de Oviedo. España. **José Luis Terreros Blanco.** Subdirector Adjunto del Gabinete del Consejo Superior de Deportes. España. **Juan Ramón Valentí Nin.** Universidad de Navarra. España. **José Antonio Villegas García.** Académico de número de la Real Academia de Medicina de Murcia. España. **Mario Zorzoli.** International Cycling Union. Suiza.



UCAM
UNIVERSIDAD
CATÓLICA DE MURCIA



ANALIZADOR de LACTATO

LACTATE PLUS

CÓMODO

El analizador Lactate Plus no necesita calibración

RÁPIDO

Tiempo de medición de 13 segundos. Sólo requiere 0.7 microlitros de sangre

PRECISO

Numerosos estudios demuestran la exactitud del Lactate Plus

ECONÓMICO

Coste por Análisis significativamente más bajo que en otras marcas



Laktate

www.laktate.com

619 284 022 



Archivos de medicina del deporte

Volumen 36(2) - Núm 190. March - April 2019 / Marzo - Abril 2019

Summary / Sumario

Editorial

Strength training, the forgotten component when prescribing physical exercise for health
La fuerza, la olvidada en la prescripción del ejercicio físico para la salud

José Antonio de Paz Fernández.....66

Original articles / Originales

Relationships among motor coordination, body mass index and physical activity in adolescents with different weight status
Relaciones entre la coordinación motora, índice de masa corporal y la actividad física en adolescentes con diferentes estados de peso corporal

Daniel V. Chagas, Luiz A. Batista69

Time limit at peak speed without prior warm-up: Effects on test duration, heart rate and rating of perceived exertion
Tiempo límite en la velocidad máxima sin calentamiento previo: efectos sobre la duración de la prueba, frecuencia cardíaca y grado de esfuerzo percibido

Guilherme H.S. Kimura, Francisco A. Manoel, Diogo H. Figueiredo, Cecília S. Peserico, Fabiana A. Machado74

Acute effects of resistance and concurrent exercise on the lipid profile of postmenopausal women
Efectos agudos del ejercicio resistido y concurrente en el perfil lipídico de mujeres postmenopáusicas

Roberto Rebolledo-Cobos, Andrés Peñaloza Florez, Luz Sarmiento-Rubiano, Laura Ardila-Pereira, Jimmy Becerra Enriquez, Adel Rodríguez-Delgado.....79

Sweating and core temperature in athletes training in continuous and intermittent sports in tropical climate
Sudoración y temperatura interna en atletas durante entrenamiento para deportes continuos e intermitentes en clima tropical

Anita M. Rivera-Brown, Farah A. Ramírez-Marrero, Jessica Frontanés, Raúl A. Rosario, Osvaldo J. Hernández, Walter R. Frontera.....86

The effect of tapering and *Nigella sativa* on the histological structure of the lung after increasing interval exercise training
El efecto del tapering y *Nigella sativa* sobre la estructura histológica del pulmón después de aumentar el entrenamiento de ejercicio de intervalo

Shadmehr Mirdar, Forough Neyestani, Gholamreza Hamidian, Mohammad Ali Khalilzadeh.....92

Reviews / Revisiones

Androgens from physiology, through pharmacy and pharmacology to the status of lifestyle drugs - are we going in the right direction?
Andrógenos, de la fisiología, a través de la farmacia y la farmacología al estado de las drogas de vida, ¿vamos en la dirección correcta?

Nataša Duborija-Kovačević, Mitar Popović, Rade Kovač.....101

Sport classification regulations for athletes with differences in sexual development (DSD)
Reglamento de clasificación deportiva para atletas con diferencias en el desarrollo sexual (DSD)

Pedro Manonelles Marqueta, José Luis Terreros Blanco, Juan José Rodríguez Sendín109

VIII Jornadas Nacionales de Medicina del Deporte117

Books / Libros120

Agenda / Agenda121

Guidelines for authors / Normas de publicación125

Strength training, the forgotten component when prescribing physical exercise for health

La fuerza, la olvidada en la prescripción del ejercicio físico para la salud

José Antonio de Paz Fernández

Especialista en Medicina del deporte. Instituto de Biomedicina (IBIOMED). Facultad CCAA Física y el Deporte. Universidad de León.

The condition of being new in order to accept or reject a paradigm, is two sides of the same counterfeit coin.

Some 9 years have passed since the WHO published its recommendations on physical activity for health, offering the general public some minimum recommended targets for daily physical activity. The main new feature of this guide is the inclusion of the recommendation to perform muscle-strengthening activities (musculoskeletal strengthening) at least two or three times a week, regardless of age (in fact from 5 years upwards)¹. Even so, these recommendations are all too frequently not included in the general exercise guidelines provided by medical practitioners.

And it is even more disturbing to see how old beliefs, that are now known to be false, are still being maintained and conveyed, with regard to “how strength training for children is a bad thing” as it affects the growth cartilage, or produces myocardial hypertrophy ... or “strengthening activities are really bad for those with high blood pressure or heart conditions” etc. Numerous studies that demonstrate the safety and benefits of exercises of this type on healthy children²⁻⁴, obese individuals⁵, or with heart diseases⁶, have been published. However, despite all this scientific and professional evidence, it has not been possible to do away with these misconceptions.

Luckily for the health of the general public, physical exercise is currently being recommended for patients affected by most chronic or degenerative disorders. However, medical practitioners are still not clearly recommending muscle-strengthening activities and, in some cases, they are even expressly advising against such exercises. This point can be illustrated by a survey conducted on 272,887 non-hypertensive and 179,789 hypertensive Americans. The respondents were asked whether, during their visit to the doctor, they had been recommended

to exercise and, if so, whether they were recommended to do only aerobic exercises, solely strengthening exercises or a combination of the two⁷. 36.9% of the healthy respondents had not been advised to exercise, neither had 45.7% of the hypertensive respondents. Of those who had been advised to exercise, 23.1% and 15% were recommended to do combined exercises and 9.5% and 7.3% solely strengthening exercises, for healthy and hypertensive respondents respectively. And this is despite the consensus agreements on exercise published by the various medical specialty or patient welfare societies. Even so, unfortunately it is not uncommon for patients suffering from a chronic disease or with after-effects to be advised against strengthening exercises. This is despite the evidence to the contrary found in scientific publications, showing how important and safe it is to do strengthening exercises, either on their own or together with the traditional aerobic exercises. Such is the case for patients suffering from ischemic heart disease^{8,9}, hypertension¹⁰, diabetes¹¹, arthritis¹², osteoporosis¹³, multiple sclerosis¹⁴, cancer survivors in general¹⁵ and breast cancer survivors in particular^{16,17}, chronic obstructive pulmonary disease^{18,19}, chronic kidney disease²⁰... and a long list of other diseases and disorders.

Due to the discrepancy between what is known through science and what is recommended by medical practitioners, there are reduced possibilities of functional improvement and an improvement in the quality of life of the healthy population and infirm population alike. It would be advisable for those in sports medicine to intensify their pedagogical efforts towards general and specialty medicine with regard to the prescription of strengthening exercises. There are sufficient scientific consensus, stances and recommendations to make it possible to correctly prescribe strength training in the healthcare context, for the general public^{21,22}, children²³, and cardiac patients²⁴.

Correspondence: José Antonio de Paz Fernández
E-mail: japazf@unileon.es

Although it is true that the benefits of exercise on health are related to the duration, intensity and frequency of the exercises, it is no less true that there is a relationship between the amount of strength training and its benefits on health. Not every exercise in which a resistance is successively overcome can be considered to be suitable strength training or a beneficial exercise to improve strength.

Strength training or exercises are not contraindicated. However poor training (incorrect load selection, incorrect load progression, incorrect technical execution, inadequate and unsafe means, etc.) is contraindicated.

Bibliography

- World Health Organization. OMS | Recomendaciones mundiales sobre la actividad física para la salud. WHO. 2013. https://www.who.int/dietphysicalactivity/factsheet_recommendations/es/. Accessed January 21, 2019.
- Myers AM, Beam NW, Fakhoury JD. Resistance training for children and adolescents. *Transl Pediatr*. 2017;6(3):137-43.
- Faigenbaum AD. Youth Resistance Training: The Good, the Bad, and the Ugly—The Year That Was 2017. *Pediatr Exerc Sci*. 2018;30(1):19-24.
- Faigenbaum AD, Myer GD. Pediatric Resistance Training. *Curr Sports Med Rep*. 2010; 9(3):161-8.
- Goldfield GS, Kenny GP, Alberga AS, et al. Effects of aerobic training, resistance training, or both on psychological health in adolescents with obesity: The HEARTY randomized controlled trial. *J Consult Clin Psychol*. 2015;83(6):1123-35.
- Hollings M, Mavros Y, Freeston J, Fiatarone Singh M. The effect of progressive resistance training on aerobic fitness and strength in adults with coronary heart disease: A systematic review and meta-analysis of randomised controlled trials. *Eur J Prev Cardiol*. 2017;24(12):1242-59.
- Mu L, Cohen AJ, Mukamal KJ. Prevalence and predictors of resistance and aerobic exercise among hypertensive adults in the United States. *J Hum Hypertens*. 2015;29(6):394-5.
- Marzolini S, Oh PI, Brooks D. Effect of combined aerobic and resistance training versus aerobic training alone in individuals with coronary artery disease: A meta-analysis. *Eur J Prev Cardiol*. 2012;19(1):81-94.
- Xanthos PD, Gordon BA, Kingsley MIC. Implementing resistance training in the rehabilitation of coronary heart disease: A systematic review and meta-analysis. *Int J Cardiol*. 2017;230:493-508.
- de Sousa EC, Abrahim O, Ferreira ALL, Rodrigues RP, Alves EAC, Vieira RP. Resistance training alone reduces systolic and diastolic blood pressure in prehypertensive and hypertensive individuals: meta-analysis. *Hypertens Res*. 2017;40(11):927-31.
- Nery C, Moraes SRA De, Novaes KA, Bezerra MA, Silveira PVDC, Lemos A. Effectiveness of resistance exercise compared to aerobic exercise without insulin therapy in patients with type 2 diabetes mellitus: a meta-analysis. *Brazilian J Phys Ther*. 2017;21(6):400-15.
- Vincent KR, Vincent HK. Resistance Exercise for Knee Osteoarthritis. *PM&R*. 2012;4(5 Suppl):S45-S52.
- Bolam KA, van Uffelen JGZ, Taaffe DR. The effect of physical exercise on bone density in middle-aged and older men: A systematic review. *Osteoporos Int*. 2013;24(11):2749-62.
- Kjølhede T, Vissing K, Dalgas U. Multiple sclerosis and progressive resistance training: a systematic review. *Mult Scler J*. 2012;18(9):1215-28.
- Fuller JT, Hartland MC, Maloney LT, Davison K. Therapeutic effects of aerobic and resistance exercises for cancer survivors: a systematic review of meta-analyses of clinical trials. *Br J Sports Med*. 2018;52(20):1311.
- Cheema BS, Kilbreath SL, Fahey PP, Delaney GP, Atlantis E. Safety and efficacy of progressive resistance training in breast cancer: a systematic review and meta-analysis. *Breast Cancer Res Treat*. 2014;148(2):249-68.
- Santos WDN dos, Gentil P, de Moraes RF, et al. Chronic Effects of Resistance Training in Breast Cancer Survivors. *Biomed Res Int*. 2017;2017:1-18.
- Iepsen UW, Jørgensen KJ, Ringbaek T, Hansen H, Skrubbeltrang C, Lange P. A Systematic Review of Resistance Training Versus Endurance Training in COPD. *J Cardiopulm Rehabil Prev*. 2015;35(3):163-72.
- Liao W -h., Chen J -w., Chen X, et al. Impact of Resistance Training in Subjects With COPD: A Systematic Review and Meta-Analysis. *Respir Care*. 2015;60(8):1130-45.
- Chan D, Cheema BS. Progressive Resistance Training in End-Stage Renal Disease: Systematic Review. *Am J Nephrol*. 2016;44(1):32-45.
- American College of Sports Medicine. Progression Models in Resistance Training for Healthy Adults. *Med Sci Sport Exerc*. 2009;41(3):687-708.
- Kraemer WJ, Ratamess NA. Fundamentals of resistance training: progression and exercise prescription. *Med Sci Sports Exerc*. 2004;36(4):674-88. <http://www.ncbi.nlm.nih.gov/pubmed/15064596>. Accessed February 3, 2019.
- Behm DG, Faigenbaum AD, Falk B, Klentrou P. Canadian Society for Exercise Physiology position paper: resistance training in children and adolescents. *Appl Physiol Nutr Metab*. 2008;33(3):547-61.
- Williams MA, Haskell WL, Ades PA, et al. Resistance Exercise in Individuals With and Without Cardiovascular Disease: 2007 Update. *Circulation*. 2007;116(5):572-84.

Analizador Instantáneo de Lactato Lactate Pro 2

arkray
LT-1730

- Sólo 0,3 µl de sangre
- Determinación en 15 segundos
- Más pequeño que su antecesor
- Calibración automática
- Memoria para 330 determinaciones
- Conexión a PC
- Rango de lectura: 0,5-25,0 mmol/litro
- Conservación de tiras reactivas a temperatura ambiente y
- Caducidad superior a un año



Importador para España:



c/ Lto. Gabriel Miro, 54, ptas. 7 y 9
46008 Valencia Tel: 963857395
Móvil: 608848455 Fax: 963840104
info@bermellelectromedicina.com
www.bermellelectromedicina.com



Monografías Femede nº 12
Depósito Legal: B. 27334-2013
ISBN: 978-84-941761-1-1
Barcelona, 2013
560 páginas.



Dep. Legal: B.24072-2013
ISBN: 978-84-941074-7-4
Barcelona, 2013
75 páginas. Color



Índice

Foreward
Presentación
1. Introducción
2. Valoración muscular
3. Valoración del metabolismo anaeróbico
4. Valoración del metabolismo aeróbico
5. Valoración cardiovascular
6. Valoración respiratoria
7. Supuestos prácticos
Índice de autores

Índice

Introducción
1. Actividad mioeléctrica
2. Componentes del electrocardiograma
3. Crecimientos y sobrecargas
4. Modificaciones de la secuencia de activación
5. La isquemia y otros indicadores de la repolarización
6. Las arritmias
7. Los registros ECG de los deportistas
8. Términos y abreviaturas
9. Notas personales

Información: www.femede.es

Relationships among motor coordination, body mass index and physical activity in adolescents with different weight status

Daniel V. Chagas, Luiz A. Batista

Laboratory of Biomechanics and Motor Behavior, Institute of Physical Education and Sports, Rio de Janeiro State University, Rio de Janeiro, Brazil.

Received: 11.01.2018

Accepted: 25.05.2018

Summary

Purpose: To analyze the influence of physical activity level on associations between motor coordination and body mass index (BMI) in normal weight, overweight and obese adolescents.

Method: Thirty nine adolescents (13 normal weight, 13 overweight and 13 obese) aged 12-14 years old, participated in this study. The Physical Activity Questionnaire for Older Children (PAQ-C) and Körperkoordinationstest für kinder (KTK) tools were used in order to assess the levels of physical activity and motor coordination, respectively. Bivariate and partial correlations were used to analyze the interrelationships among motor coordination, BMI and physical activity. The analysis of covariance test was used in order to compare the levels of motor coordination among normal weight, overweight and obese adolescents, considering the physical activity level as a covariate.

Results: Moderate negative correlations ($p < 0.05$) were found between motor coordination and BMI in the sample as a whole, normal weight and obese groups. However, when controlled for physical activity levels, it was not observed a significant correlation in the normal weight group. Motor coordination and BMI were not significantly correlated in overweight adolescents with and without controlling for physical activity levels. Furthermore, motor coordination level demonstrated a trend to be lower in overweight and obese adolescents.

Conclusion: Physical activity level influenced the association between motor coordination and BMI in normal weight adolescents, but not in overweight and obese. The negative effect of excess body mass on motor coordination level may overlap the possible influence that physical activity level exerts on the association between motor coordination and BMI in overweight and obese adolescents.

Key words:

Motor coordination. Obesity.
Overweight. Adolescents.
Biomechanics.

Relaciones entre la coordinación motora, índice de masa corporal y la actividad física en adolescentes con diferentes estados de peso corporal

Resumen

Objetivo: Analizar la influencia del nivel de actividad física en las asociaciones entre la coordinación motora y el índice de masa corporal (IMC) en adolescentes de peso normal, con sobrepeso y obesos.

Método: Treinta y nueve adolescentes (13 con peso normal, 13 con sobrepeso y 13 obesos) con edades comprendidas entre 12 y 14 años participaron en este estudio. Las herramientas Physical Activity Questionnaire for Older Children (PAQ-C) y Körperkoordinationstest für kinder (KTK) se usaron para evaluar los niveles de actividad física y coordinación motriz, respectivamente. Se usaron correlaciones bivariadas y parciales para analizar las interrelaciones entre la coordinación motora, el IMC y la actividad física. El análisis de covarianza se utilizó para comparar los niveles de coordinación motora entre los adolescentes de peso normal, con sobrepeso y obesos, considerando el nivel de actividad física como una covariable.

Resultados: se encontraron correlaciones negativas y moderadas ($p < 0.05$) entre la coordinación motora y el IMC en la muestra como un todo, en los adolescentes con peso normal y obesos. Sin embargo, cuando se controlan los niveles de actividad física, no se observó una correlación significativa en el grupo de peso normal. La coordinación motora y el IMC no se correlacionaron significativamente en adolescentes con sobrepeso con y sin control de los niveles de actividad física. Además, el nivel de coordinación motora demostró una tendencia a ser menor en adolescentes con sobrepeso y obesos.

Conclusión: El nivel de actividad física influyó en la asociación entre la coordinación motora y el IMC en adolescentes de peso normal, pero no en adolescentes con sobrepeso y obesos. El efecto negativo del exceso de masa corporal en el nivel de coordinación motora puede superponerse a la posible influencia que ejerce el nivel de actividad física en la asociación entre la coordinación motora y el IMC en los adolescentes con sobrepeso y obesos.

Palabras clave:

Coordinación motora. Obesidad.
Sobrepeso. Adolescentes.
Biomecánica.

Correspondence: Daniel das Virgens Chagas

E-mail: chagas_daniel@yahoo.com.br

Introduction

Pediatric obesity has become a global epidemic¹, being recognized as one of the most serious public health concerns in the 21st century². In Brazil, more than 8 million children and adolescents are estimated to be obese³. This high prevalence is a concern due its associated health risks such as hypertension, hyperinsulinemia, glucose intolerance, type II diabetes and dyslipidemia².

Obesity seems also to be associated with poor motor coordination in children and adolescents⁴⁻⁶. Although few studies^{7,8} have not found significant associations between motor coordination level and body mass index (BMI), most evidence suggests these variables are negatively associated⁹⁻¹⁴. Overall, these previous studies suggest obese children tend to have lower motor coordination level than their normal weight peers.

A plausible explanation for a negative association between BMI and motor coordination level in children and adolescents is based on biomechanical issues. That is, the higher the amount of body mass, the higher the mechanical work required to performing motor tasks, especially those which demand body weight-bearing. Evidence^{10,15,16} has confirmed, at least partially, this weight-bearing hypothesis. However, the reported negative relationship between motor coordination level and BMI can be mediated by several alternative mechanisms¹⁷.

One of mechanisms that can mediate, or influence, the relationship between motor coordination level and BMI in pediatric population is the physical activity level^{4,18}. This assumption is based on evidence that physical activity level is positively associated with motor coordination level^{19,20}. Also, it is important to recognize that not all obese children and adolescents have poor motor coordination or are physically inactive²¹. Therefore, the hypothesis that physical activity level may influence the association between motor coordination level and BMI in children and adolescents seems to be plausible.

Chagas and Batista²² found physical activity level influences the association between motor coordination level and BMI in normal weight, but not in their overweight/obese peers. However, the comparisons realized in that study involved overweight and obese adolescents in the same group. Therefore, it remains unknown if physical activity level can influence the associations between motor coordination level and BMI in obese adolescents.

The purpose of this study was to analyze the influence of physical activity level on associations between motor coordination and BMI in normal weight, overweight and obese adolescents.

Material and method

One hundred and fifty five adolescents (69 boys and 86 girls) between seventh and ninth grade of elementary school (age ranged between 12 and 15 years old) from a Brazilian public school were initially recruited to participate in the study. Inclusion criteria required students to be under 15 years old with no history of injury or disease that could affect motor performance. Exclusion criteria consisted of subject's weight status classified as underweight. Ethical approval for this study was obtained from the University's Ethics

Committee and parental consent and child assent were obtained prior to participation.

After initial recruitment (N=155), eight subjects were excluded from the study due to their being underweight. Further, thirteen obese adolescents were detected (n=13). Thus, among the 134 remaining subjects (28 overweight and 106 normal weight), 13 overweight and 13 normal weight adolescents were randomly selected to compose the final sample of 39 participants (17 boys and 22 girls). There was not missing data.

Body mass was measured to the nearest 0.1 kg using an electronic scale, with participants wearing their school uniform. Standing height was measured while unshod with a stadiometer wall to the nearest 0.1 cm. BMI (kg/m²) was then calculated. World Health Organization²³ age-specific cut-off points for BMI were used in order to determine the weight status of participants (underweight, normal weight, overweight and obesity) according to gender.

The Physical Activity Questionnaire for Older Children (PAQ-C), a valid²⁴ self-applied 7-day recall instrument, was used to assess general levels of physical activity of participants. The PAQ-C is appropriate for elementary school-aged children approximately between 8–14 years old who are currently in the school system and have recess as a regular part of their school week. The summary score from the PAQ-C is the average of the sum of the nine item questions, each scored on a 5-point scale, with 1 being the lower level of physical activity and 5 the higher level.

Motor coordination level was assessed using the Körperkoordinationstest für Kinder (KTK). The KTK is appropriated to assess motor coordination level of participants because it is a reliable and valid instrument, with a teste-retest reliability coefficient of 0.97²⁵. KTK is one of most used tools for assessing children's motor coordination²⁶ and it consists of four test items. The first is walking backwards along balance beams (3m length) of decreasing width (6, 4.5 and 3 cm). Each beam was crossed three times where a maximum of eight steps per trial were allowed (72 steps overall); the sum of steps in all trials determined score 1. The second involved one-legged hopping over an obstacle, formed by an increasing pile of pillows (pillow size 60 cm × 20 cm × 5 cm; the maximum was 12 pillows or a height of 60 cm). Only three trials were allowed for each obstacle and three, two, or one point(s) were/was awarded for successful performance on the first, second, or third try, respectively. Therefore, a maximum of 39 points (including a ground level trial) could be scored for each leg; the points were summed to determine score 2. The third task was two-legged sideways jumping across a wooden slat (60 cm × 4 cm × 2 cm) for 15 s as quickly as possible. The number of jumps performed correctly was summed over two trials to determine score 3. The final task involved moving sideways on wooden boards (25 cm × 25 cm × 5.7 cm) as many times as possible in 20 s. One point was awarded for each time the plate was transferred and one more for stepping on it. The number of relocations was counted and summed over two trials to determine score 4. KTK takes into account motor coordination level is gender and age-related. Thus, the four scores acquired in each item test were gender and age-adjusted in according to KTK normative database. Finally, the motor coordination level for each participant was derived from the sum of the four adjusted scores obtained in the tests.

Descriptive statistics were determined for all variables. The Kolmogorov–Smirnov test confirmed acceptable normality of the data distribution. Pearson's correlation coefficients were used to examine the bivariate relationships between levels of motor coordination, BMI and physical activity. Partial correlations were used to analyse the relationship between motor coordination level and BMI, controlling for physical activity levels. Analysis of covariance (ANCOVA) was used to compare the motor coordination levels between normal weight, overweight and obese adolescents, considering the physical activity level as a covariate. A significance level of 5% ($\alpha = 0.05$) was adopted in all statistical tests. Data analysis was executed using Statistical Package for Social Sciences software (SPSS ver. 22.0, IBM, USA).

Results

The total sample (N=39) presented the following values for age, body weight, height, BMI, motor coordination and physical activity levels, respectively: 13.7y (± 0.6), 63.2kg (± 17.5), 1.61m (± 0.8), 24.1kg/m² (± 5.9), 78.6 (± 22.0) and 2.7 (± 0.9). Descriptive statistics of group 1 (normal weight), group 2 (overweight) and group 3 (obese) are provided in Table 1.

As shown in Figure 1, motor coordination levels decline as weight status worsens, that is, as higher BMI, the lower motor coordination levels. However, results of ANCOVA test confirmed significant differences in motor coordination levels only between normal weight and obese groups ($F=4.123, p=0.025$).

Pearson and partial correlations coefficients are displayed in Table 2. Moderate negative correlations were found between BMI and motor coordination levels in the group as a whole, as well as in the normal weight and obese groups. However, when statistically controlled for physical activity levels, significant correlations were only observed in the group as a whole and obese group.

Table 1. Descriptive statistics (mean \pm standard deviation) of age, body weight, height, BMI, motor coordination levels (MC) and physical activity levels (PA) of normal weight (group 1), overweight (group 2) and obese (group 3) adolescents.


	Group 1 (n=13)	Group 2 (n=13)	Group 3 (n=13)
Age (years)	13.7 (± 0.7)	13.6 (± 0.6)	13.9 (± 0.6)
Body Weight (kg)	45.0 (± 8.2)	63.8 (± 6.6)	80.8 (± 13.1)
Height (m)	1.58 (± 0.1)	1.64 (± 0.1)	1.62 (± 0.1)
BMI (kg/m ²)	18.0 (± 1.9)	23.6 (± 1.4)	30.8 (± 4.1)
MC	89.9 (± 13.5)	80.3 (± 21.7)	65.6 (± 23.5)
PA	2.9 (± 0.9)	2.7 (± 0.9)	2.5 (± 0.8)

Table 2. Pearson and partial correlation coefficients between BMI and motor coordination levels (two-tailed test).

	Bivariate correlations	Partial correlations [†]
Total sample	- 0.640**	- 0.622**
Group 1	- 0.569*	- 0.565 [†]
Group 2	- 0.464 [†]	- 0.465 [†]
Group 3	- 0.691**	- 0.643*

[†]Controlled for physical activity levels.
^{*}Correlation significant at the 0.05 level.
^{**}Correlation significant at the 0.01 level.
^{††}Correlation non-significant ($p > 0.05$).

Figure 1. Mean and standard error of motor coordination levels across three groups.



Discussion

The main aim of this study was to analyze the influence of physical activity level on associations between motor coordination and BMI in adolescents with different weight status. Our findings suggested motor coordination level and BMI are negatively associated in normal weight and obese adolescents. However, the association between these variables was not significant when controlled for physical activity level in normal weight adolescents. On the other hand, motor coordination level and BMI were significantly negatively correlated in obese adolescents regardless of physical activity level. These results suggest physical activity level can influence the association between motor coordination and BMI in normal weight adolescents. Furthermore, motor coordination level demonstrated a trend to be decline as weight status worsens.

Moderate negative correlations between motor coordination level and BMI were observed in the group as a whole, normal weight and obese adolescents. These findings are in line with previous studies⁹⁻¹⁴. Considering BMI as an indicator of body fat, it seems plausible to expect that individuals with higher BMI, i.e. higher adiposity, have higher difficulty to perform some motor tasks due to increased biomechanical

demands. That is, the higher BMI, higher physical difficulties experienced by individuals, such as increased moments of inertia and mechanical work, especially in weight-bearing tasks.

Nevertheless, our findings showed that motor coordination and BMI were not significantly correlated in overweight adolescents. Few previous studies^{7,8} have also not found significant associations between motor coordination level and BMI. These findings can be explained by the fact that individuals with the same weight status, or even with the same BMI, may have considerable differences in body composition. That is, overweight individuals with relatively high levels of lean mass and low levels of body fat may have no increased difficulty to perform motor tasks due their BMI status.

Negative correlations between motor coordination levels and BMI remained weak to moderate when controlled for physical activity level in all groups. However, motor coordination level and BMI were not significantly correlated in normal weight adolescents when controlled for physical activity level. This evidence suggests physical activity level can influence the association between motor coordination and BMI in normal weight adolescents, but not in overweight and obese. These findings are in line with a previous study²², corroborating the assumption that, in obese adolescents, motor coordination level and BMI are negatively associated regardless of physical activity levels. These results can be explained by negative effect of excess body mass on motor coordination level; it may overlap the possible influence that physical activity level exerts on the association between motor coordination and BMI in overweight and obese adolescents.

In line with previous findings^{5,10,17}, our results showed motor coordination level tend to decline as weight status worsens. These results add to the body of evidence suggesting obese adolescents tend to have poorer motor coordination level than their normal weight peers. These findings were expected due to increased biomechanical demands in some motor tasks experienced by individuals with higher BMI. However, significant differences in motor coordination level were observed only between normal weight and obese groups. That is, there were no significant differences in motor coordination level when overweight adolescents were compared with normal weight and obese peers. These findings can also be explained by possible differences in body composition, especially with regard to body fat, among participants whose weight status was classified as overweight.

In this study was hypothesized that the physical activity level, as a mediator mechanism, could influence the associations between motor coordination and BMI in adolescents with different weight status. Our findings corroborated only partially this hypothesis, because physical activity level influenced only the associations among normal weight adolescents. Besides the increased negative effect of body fat in overweight and obese, other complementary mechanisms, as perceived motor competence and physical fitness, can have influenced the approached relationship.

Another way to explain our findings concerns the role of physical activity level in this interrelationship. Indeed, it is expected the higher physical activity level among adolescents, higher their opportunities for practice. Considering that opportunities for practice is essential for motor development²⁷, it is expected that physical activity level is associated with motor coordination. However, the levels of physical

activity showed for adolescents may not necessarily to be linked with diversified and increased opportunities for practice. Thus, the influence of physical activity on relationship between motor coordination and BMI can be low or inexistent. In this sense, the role of physical activity as a mediator mechanism, such as proposed in the conceptual framework proposed by Stodden *et al.*²⁸, may vary in according to opportunities for practice experienced by individuals.

This study had some limitations. First, motor coordination and physical activity were assessed using a test composed by gross motor skills and a self-reported questionnaire, respectively. Thus, the inference of our results is limited to these specific measures in adolescents. Second, both instruments used for assessing physical activity and motor coordination were not adapted to Brazilian population. Also, the small sample size was relatively small. Future investigations using different measurement tools should be conducted in order to corroborate or refuse our results.

Conclusions

Physical activity level influenced the association between motor coordination and BMI in normal weight adolescents, but not in overweight and obese. The increased negative effects of excess body mass, as well as the variability of opportunities for practice according to physical activity level, seem to explain these findings. Also, complementary analyses assessing the potential mediating effects of physical activity on the relationship between motor coordination and BMI should be considered in future studies. As practical application, physical educators and physicians must encourage not only a health weight status, but also the development of motor coordination among adolescents. Moreover, elevated levels of physical activity associated with diversified and increased opportunities for practice should also be promoted by health care professionals, as physical educators and physicians.

Conflict of interest

The authors do not declare a conflict of interest.

Bibliography

1. Han JC, Lawlor DA, Kimm S. Childhood obesity – 2010: progress and challenges. *Lancet*. 2010;375:1737-48.
2. Güngör NK. Overweight and obesity in children and adolescents. *J Clin Res Pediatr Endocrinol*. 2014;6:129-43.
3. Aiello AM, Marques de Mello L, Souza Nunes M, Soares da Silva A, Nunes A. Prevalence of obesity in children and adolescents in Brazil: a meta-analysis of cross-sectional studies. *Curr Pediatr Rev*. 2015;11:36-42.
4. D'Hondt E, Deforche B, Gentier I, De Bourdeaudhuij I, Vaeyens R, Philippaerts R, Lenoir M. A longitudinal analysis of gross motor coordination in overweight and obese children versus normal-weight peers. *Int J Obes (Lond)*. 2013;37:61-7.
5. Antunes AM, Maia JA, Stasinopoulos MD, Gouveia ER, Thomis MA, Lefevre JA, et al. Gross motor coordination and weight status of Portuguese children aged 6-14 years. *Am J Hum Biol*. 2015;27:681-9.
6. D'Hondt E, Deforche B, Vaeyens R, Vandorpe B, Vandendriessche J, Pion J, Philippaerts R, Lenoir M. Gross motor coordination in relation to weight status and age in 5- to 12-year-old boys and girls: a cross-sectional study. *Int J Pediatr Obes*. 2011;6 (2-2):556-64.
7. Catenassi FZ, Marques I, Bastos CB, Basso L, Ronque E, Gerage A. Relação entre índice de massa corporal e habilidade motora grossa em crianças de quatro a seis anos. *Rev Bras Med Esporte*. 2007;13:227-30.

8. Spessato BC, Gabbard C, Valentini NC. The role of motor competence and body mass index in children's activity levels in physical education classes. *J Teach Phys Educ.* 2013;32:118-30.
9. Lopes V, Stodden D, Bianchi M, Maia J, Rodrigues L. Correlation between BMI and motor coordination in children. *J Sci Med Sport.* 2012;15:38-43.
10. D'Hondt E, Deforche B, I Bourdeaudhuij, Lenoir M. Relationship between motor skill and body mass index in 5- to 10-year old children. *Adapt Phys Activ Q.* 2009;26:31-7.
11. Lima RA, Bugge A, Pfeiffer KA, Andersen LB. Tracking gross motor coordination from childhood into adolescence. *Res Q Exerc Sport.* 2017;88:52-9.
12. Hardman CM, Wanderley Junior RS, Oliveira E, Barros M. Relationship between physical activity and BMI with level of motor coordination performance in schoolchildren. *Rev Bras Cineantropom Desempenho Hum.* 2017;19:50-61.
13. Logan SW, Scrabis-Fletcher K, Modlesky C, Getchell N. The relationship between motor skill proficiency and body mass index in preschool children. *Res Q Exerc Sport.* 2011;82:442-8.
14. Freitas J, Castro P, Rezende E, Werneck F, Lima J. Relationship between the overweight and the motor coordination in young athletes of athletics. *RBCE.* 2017;39:91-7.
15. Vandendriessche J, Vandorpe B, Coelho-e-Silva M, Vaeyens R, Lenoir M, Lefevre J, et al. Multivariate association among morphology, fitness, and motor coordination characteristics in boys age 7 to 11. *Pediatr Exerc Sci.* 2011;23:504-20.
16. Zhu YC, Wu SK, Cairney J. Obesity and motor coordination ability in Taiwanese children with and without developmental coordination disorder. *Res Dev Disabil.* 2011;32:801-7.
17. D'Hondt E, Deforche B, Gentier I, Verstuyf J, Vaeyens R, De Bourdeaudhuij I, Philippeaerts R, Lenoir M. A longitudinal study of gross motor coordination and weight status in children. *Obesity. (Silver Spring)* 2014;22:1505-11.
18. Chagas DV, Batista LA. Interrelationships among motor coordination, body fat percentage, and physical activity in adolescent girls. *Hum Mov.* 2015;16:4-8.
19. Lubans DR, Morgan PJ, Cliff DP, Barnett LM, Okely AD. Fundamental movement skills in children and adolescents: review of associated health benefits. *Sports Med.* 2010;40:1019-35.
20. Kambas A, Michalopoulou M, Fatouros IG, Christoforidis C, Manthou E, Giannakidou D, et al. The relationship between motor proficiency and pedometer-determined physical activity in young children. *Pediatr Exerc Sci.* 2012;24:34-44.
21. Morrison K, Bugge A, El-Naaman B, Eisenmann J, Froberg K, Pfeiffer K, et al. Interrelationships among physical activity, body fat, and motor performance in 6- to 8-year-old Danish children. *Pediatr Exerc Sci.* 2012;24:199-209.
22. Chagas DV, Batista LA. Associations between motor coordination and BMI in normal weight and overweight/obese adolescents. *J Hum Growth Dev.* 2016;26:380-4.
23. De Onis A, Onyango A, Borghi E, Siyam A, Nishida C, Siekmann J. Development of a WHO growth reference for school-aged children and adolescents. *Bull World Health Organ.* 2007;85:660-7.
24. Kowalski K, Crocker P, Faulkner R. Validation of the Physical Activity Questionnaire for Older Children. *Pediatr Exerc Sci.* 1997;9:174-86.
25. Vandorpe B, Vandendriessche J, Lefevre J, Pion J, Vaeyens R, Matthys S, et al. The KörperkoordinationsTest für Kinder: reference values and suitability for 6-12-year-old children in Flanders. *Scand J Med Sci Sports.* 2011;21:378-88.
26. Cools W, De Martelaer K, Samaey C, Andries C. Movement skill assessment of typically developing preschool children: a review of seven movement skill tools. *J Sports Sci Med.* 2009;8:154-68.
27. Gallahue D, Ozmun J, Goodway J. *Understanding motor development: infants, children, adolescents, adults.* New York: McGraw-Hill; 2012.
28. Stodden DF, Goodway JD, Langendorfer SJ, Robertson MA, Rudisill ME, Garcia C, et al. A developmental perspective on the role of motor skill competence in physical activity: an emergent relationship. *Quest.* 2008;60:290-306.

Time limit at peak speed without prior warm-up: Effects on test duration, heart rate and rating of perceived exertion

Guilherme H.S. Kimura, Francisco A. Manoel, Diogo H. Figueiredo, Cecília S. Peserico, Fabiana A. Machado

State University of Maringá - UEM, Brazil.

Received: 01.02.2018

Accepted: 28.06.2018

Summary

Background: Time limit (t_{lim}) at peak speed (V_{peak}), that is maximal time that an individual can keep running at maximal intensity, is used to prescribe interval durations during interval training. The aim of this study was to compare two protocols (without or with 15 minutes of warm-up) for the t_{lim} determination at 100% of V_{peak} in untrained men.

Material and method: Twelve untrained young men performed three running tests on a treadmill: one maximal incremental test to determine V_{peak} and two rectangular tests, performed in randomized order, without a warm-up (t_{lim0}) or with a 15-minute warm-up duration (t_{lim15}) at 60% of V_{peak} to determine the t_{lim} at 100% of V_{peak} after the warm-up the tests were performed at the speed of the individual V_{peak} until volitional exhaustion. During the tests, heart rate (HR) and rating of perceived exertion (RPE) were monitored and blood lactate sampling was collected prior to session (LA_{pre}); immediately post-exercise (LA_{0-min}); 3 min (LA_{3-min}), 5 min (LA_{5-min}) and 7 min (LA_{7-min}) post-exercise to determine lactate concentrations. The Shapiro-Wilk test was used and confirmed the normality of the data distribution, with maximal and submaximal values compared using Student's t test for dependent samples.

Results: Test duration at t_{lim0} was significant higher than that at t_{lim15} ($P = 0.02$). Additionally, different t_{lim} protocols influenced HR and RPE submaximal responses and did not modify lactate concentrations or maximal variables (HR_{max} and RPE_{max}).

Conclusions: These findings suggest that the determination of t_{lim} at 100% of V_{peak} without a prior warm-up led to a higher test duration in untrained men.

Key words:

Running. Exercise test. Physical endurance.

Tiempo límite en la velocidad máxima sin calentamiento previo: efectos sobre la duración de la prueba, frecuencia cardíaca y grado de esfuerzo percibido

Resumen

Objetivos: Tiempo límite (t_{lim}) en la velocidad máxima (V_{pico}), que es el tiempo máximo que un individuo puede permanecer corriendo en la intensidad máxima, se utiliza para describir la duración de los intervalos durante el entrenamiento interválico. El objetivo de este estudio fue comparar dos protocolos (sin o con 15 minutos de calentamiento) para la determinación de t_{lim} al 100% de la V_{pico} en hombres no entrenados.

Material y método: Doce jóvenes no entrenados realizaron tres pruebas de carrera en la cinta rodante: una prueba incremental máxima para determinar V_{pico} y dos pruebas rectangulares, realizadas en orden aleatorio, sin calentamiento (t_{lim0}) o con una duración de calentamiento de 15 minutos (t_{lim15}) al 60% de la V_{pico} para determinar el t_{lim} al 100% de la V_{pico} ; después del calentamiento las pruebas fueron realizadas en la velocidad de la V_{pico} individual hasta el agotamiento voluntario. Durante las pruebas, la frecuencia cardíaca (FC) y el grado de esfuerzo percibido (RPE) fueron monitorizadas y se tomaron muestras de sangre antes de la sesión (LA_{pre}); inmediatamente después del ejercicio (LA_{0-min}); 3 min (LA_{3-min}), 5 min (LA_{5-min}) y 7 min (LA_{7-min}) después del ejercicio para determinar las concentraciones de lactato. Se utilizó la prueba de Shapiro Wilk y se confirmó la normalidad de la distribución de los datos, con los valores máximos y submáximos comparados utilizando la prueba t de Student para muestras dependientes.

Resultados: La duración de la prueba en el t_{lim0} fue significativamente mayor que aquella en el t_{lim15} ($P = 0.02$). Además, los diferentes protocolos de t_{lim} influenciaron las respuestas submáximas de FC y RPE y no modificaron las concentraciones de lactato o las variables máximas (FC_{max} y RPE_{max}).

Conclusiones: Estos resultados sugieren que la determinación del t_{lim} en la 100% de la V_{pico} sin calentamiento previo lleva a una mayor duración de la prueba en hombres no entrenados.

Palabras clave:

Carrera. Prueba de ejercicio. Resistencia física.

Correspondence: Fabiana Andrade Machado

E-mail: famachado_uem@hotmail.com; famachado@uem.br

Introduction

Endurance racing has been gaining popularity recently; thus, the training prescriptions linked to running performance improvements have received great attention^{1,2}. Such prescriptions should be planned for each individual according to the physiological (*e.g.*, heart rate [HR]), psychological (*e.g.*, rating of perceived exertion [RPE]), and performance variables (peak speed [V_{peak}], time limit [t_{lim}]) that control and monitor effort intensity³.

The t_{lim} is the maximal time that an individual can maintain a given exercise intensity⁴, such as V_{peak} occurrence velocity. The original protocol for the t_{lim} determination included a prior 15-minute warm-up at an intensity equivalent to 60% of the reference velocity (*e.g.*, V_{peak} or velocity associated with the occurrence of maximal oxygen uptake [$\dot{V}O_{2\text{max}}$]); after the warm-up, without interval, the velocity is automatically increased to 100% intensity, at which the individual should remain as long as possible until volitional exhaustion⁴. However, warm-up duration (5, 10 or 15 minutes) modified the test duration (t_{lim})⁵.

Studies demonstrated that various warm-up types, such as stretching (static and dynamic)⁶, whole-body vibration⁷, and the traditional warm-up consisting of low-intensity cycling⁸, could positively or negatively influence aerobic performance. For instance, Tomaras & Macintosh⁹ investigated a sample of highly trained male track cyclists and compared the traditional warm-up (WU) for a 200-m sprint in a track cycling competition with an experimental WU that was designed to be shorter and less intense and examined the fatigue and cycling performance after traditional and experimental WU. Results from this study showed that peak active twitch torque was lower after the traditional than experimental WU when expressed as percentage of pre-warm-up amplitude, and Wingate test performance was better after experimental WU than traditional WU; indeed, the traditional track cyclist's WU results in significant fatigue, which corresponds with impaired peak power output, and shorter and lower intensity WU permits a better performance.

However, a recent study showed that warm-up duration (5, 10 or 15 minutes) modified the test duration (t_{lim}) of untrained men in which the warm-up comprising 15 minutes led the participants to exercise for a shorter time compared to other t_{lim} protocols⁵. Since t_{lim} is a variable that is used to determine the optimal stimulus durations in interval training sessions⁹⁻¹¹, this longer or short duration altered by warm-up will directly impact this prescription.

Nevertheless, to the best of our knowledge, there is still a lack of studies that analyzed the best duration of the warm-up⁵ and knowing the importance of warm-ups prior to exercise and that the duration modifies the t_{lim} ⁵, the aim of this study was to compare two protocols (without and with a 15-minute warm-up) for the t_{lim} determination at 100% of V_{peak} in untrained men. Our hypothesis is that the t_{lim} determined without a warm-up will differ from that of the t_{lim} performed with a 15-minute warm-up.

Material and method

Twelve untrained young men, not included in systematic running training programs, with training volume less than 20 km per week,

volunteered to participate in this study. The 5-km running times reported by participants were between 25 and 35 minutes, with a pace between 8.6 and 12 km·h⁻¹ (\cong 36.1 and 50.4 % of the world record). The main characteristics of the participants were: age 21.4 \pm 2.3 years, height 1.8 \pm 0.1 m, body mass 76.7 \pm 10.9 kg, body mass index (BMI) 24.2 \pm 2.8 kg·m⁻² and body fat 15.3 \pm 4.2%. Prior to testing, written informed consent was obtained from all participants. The experimental protocol was approved by the local Human Research Ethics Committee (#1.262.502/2015) and appropriate standards for human experimentation have been followed.

Experimental overview

Participants performed three tests on a motorized treadmill (Super ATL Inbrasport, Porto Alegre, Brazil), with the gradient set at 1%. In the first visit the evaluation of the anthropometric measurements was performed, and the participants were submitted to a maximal incremental test to determine V_{peak} . After, in a randomized order, two rectangular tests with warm-up durations of 15 minutes and without warm-up were performed to determine the t_{lim} at 100% of V_{peak} . The tests were performed over 2 weeks, with each test separated from the other by 48 hours. For all tests, participants were instructed to stay well-rested, well-nourished, and well-hydrated, wearing lightweight comfortable clothing. Participants were also instructed to avoid eating for 2 hours before the maximal exercise test, to abstain from caffeine and alcohol and to refrain from strenuous exercise for 48 hours before testing. Tests were conducted at the same time of the day, under normal laboratory conditions (temperature 20-22 °C, relative humidity 50-60%).

Incremental exercise test to determine V_{peak}

After a warm-up, comprised walking at 6 km·h⁻¹ for three minutes, the continuous protocol started with a speed of 8 km·h⁻¹ and increased by 1 km·h⁻¹ between each successive 3-minute stage until participants reached volitional exhaustion¹². The V_{peak} of the incremental test was calculated as the speed of the last complete stage added to the completed fraction of the incomplete stage¹³, calculated according to the equation:

$$V_{\text{peak}} = V_{\text{complete}} + (t/T \times \text{speed increment})$$

where V_{complete} is the running speed of the last complete stage, t the time in seconds sustained during the incomplete stage, and T the time in seconds required to complete a stage (*i.e.*, 180 seconds), and speed incremental is the speed load increment. The maximal effort was deemed to be achieved if the incremental test met two of the following criteria: 1) $LA_{\text{peak}} \geq 8 \text{ mmol} \cdot \text{L}^{-1}$ 14 2) $HR_{\text{max}} \geq 100\%$ of endurance-trained age-predicted HR_{max} using the age-based (207 - 0.7 \times age) equation¹⁵ and 3) $RPE_{\text{max}} \geq 19$ in the 6-20 Borg Scale¹⁶.

Rectangular tests to determine the t_{lim} at V_{peak}

The two rectangular tests differed only by presence or absence of the warm-up of 15 minutes. After the warm-up at 60% of V_{peak} , the treadmill speed was quickly increased (in approximately 6 seconds) to the individual at 100% of V_{peak} ⁴. Participants were also encouraged to invest maximal effort and the time of permanency in this intensity was considered the t_{lim} at 100% of V_{peak} .

Psychophysiological and physiological variables

Before testing participants were familiarized with the 6–20 Borg scale¹⁶, which was used to determine the rating of perceived exertion (RPE) during the last 15 seconds the stages of V_{peak} and every minute in t_{lim} . The highest RPE value was adopted as the maximal RPE (RPE_{max}). Heart rate (HR) was monitored throughout the tests (Polar RS800sd, Kempele, Finland) and in the last 10 seconds of each stage da V_{peak} and every minute in t_{lim} HR was registered; the maximal heart rate (HR_{max}) was defined as the highest HR value observed during the tests¹⁵. Earlobe capillary blood samples (25 μ L) were collected in a capillary tube to determine the lactate concentrations. These samples were collected before (LA_{pre}) all exercise tests, after the incremental test at the third (LA_{3-min}) and fifth (LA_{5-min}) minutes, and at the end (LA_{0-min}), at the LA_{3-min} , LA_{5-min} and seventh (LA_{7-min}) minutes after the rectangular tests. For the LA_{pre} the participants remained at rest for 15 minutes in a comfortable chair prior to the sampling procedure. For the LA_{0-min} blood sampled collection the participants remained standing upright on the treadmill, and for the LA_{3-min} , LA_{5-min} , LA_{7-min} samples the participants remained sitting in a comfortable chair. Peak blood lactate concentration (LA_{peak}) was defined for each participant as the highest post-exercise blood lactate concentration value. The samples were subsequently determined by electroenzymatic methods using the YSI 2300 STAT (Ohio, USA) automated analyzer (accuracy \pm 2%).

Statistical analysis

Data are presented as mean \pm SD and were analyzed using the Statistical Package for the Social Sciences software v. 20.0 (SPSS Inc., Chicago, IL, USA). The Shapiro-Wilk test was used and confirmed the normality of the data distribution. Maximal and submaximal values were compared using Student’s *t* test for dependent samples. Statistical significance was set at $P < 0.05$.

Results

The variables obtained during the exhaustion incremental test to determine the V_{peak} were: $V_{peak} = 12.7 \pm 1.2 \text{ km}\cdot\text{h}^{-1}$, test duration = $17.1 \pm 3.8 \text{ min}$, $HR_{max} = 187 \pm 8.7 \text{ bpm}$, $RPE_{max} = 19.8 \pm 0.5$ and $LA_{peak} = 8.6 \pm 3.2 \text{ mmol}\cdot\text{L}^{-1}$.

Table 1 compares t_{lim} test duration, maximal variables (HR_{max} , RPE_{max} and LA_{peak}) and post-test lactate concentrations obtained during the two tests. Test duration at t_{lim0} was significantly higher than that at t_{lim15} ($P = 0.02$). However, no significant difference was seen in the other variables.

Comparisons between HR and RPE values obtained during the two tests for t_{lim} determination are shown in Table 2. Only the minutes in which the participants remained throughout the two tests were analyzed. Significant differences were noted in the HR and RPE values until the fifth minute ($P \leq 0.01$), in the RPE value at the sixth minute ($P = 0.04$), and in HR value at seventh minute ($P = 0.02$).

Table 1. Comparison between the variables: test duration (min), HR_{max} (bpm), RPE_{max} (6-20, AU) and blood lactate concentrations ($\text{mmol}\cdot\text{L}^{-1}$) obtained during the t_{lim0} and t_{lim15} (N = 12).

Variables	t_{lim0}	t_{lim15}	P
Test duration (min)	9.4 \pm 2.2	6.0 \pm 2.0*	0.02
HR_{max} (bpm)	184 \pm 11.0	184 \pm 10.9	0.80
RPE_{max} (6-20, AU)	19.9 \pm 0.3	19.9 \pm 0.3	1.00
LA_{pre} ($\text{mmol}\cdot\text{L}^{-1}$)	1.1 \pm 0.3	1.0 \pm 0.2	0.30
LA_{0-min} ($\text{mmol}\cdot\text{L}^{-1}$)	8.5 \pm 2.0	7.1 \pm 2.4	0.07
LA_{3-min} ($\text{mmol}\cdot\text{L}^{-1}$)	8.6 \pm 2.7	8.2 \pm 3.1	1.00
LA_{5-min} ($\text{mmol}\cdot\text{L}^{-1}$)	8.4 \pm 2.3	7.7 \pm 3.3	0.11
LA_{7-min} ($\text{mmol}\cdot\text{L}^{-1}$)	8.1 \pm 2.3	7.5 \pm 3.3	0.42
LA_{peak} ($\text{mmol}\cdot\text{L}^{-1}$)	9.5 \pm 2.6	8.4 \pm 3.3	0.13

Note: AU: arbitrary units; bpm: beat per minute; HR_{max} : maximal heart rate; LA_{pre} : blood lactate concentration at the before of the test; LA_{0-min} : blood lactate concentration at the end of the test; LA_{3-min} : blood lactate concentration at the third minute after the test; LA_{5-min} : blood lactate concentration at the fifth minute after the test; LA_{7-min} : blood lactate concentration at the seventh minute after the test; LA_{peak} : peak blood lactate concentration; RPE_{max} : maximal rating of perceived exertion.

* $P < 0.05$ compared with t_{lim0} .

Table 2. Comparison between the HR (bpm) and RPE (6-20, AU) submaximal values (min) at each minute obtained during the t_{lim0} and t_{lim15} .

Time	HR (bpm)			RPE (AU)		
	t_{lim0}	t_{lim15}	P	t_{lim0}	t_{lim15}	P
1 min (n=12)	143 \pm 12.4	167 \pm 10.9*	< 0.01	8.0 \pm 2.4	13.8 \pm 2.6*	< 0.01
2 min (n=12)	159 \pm 11.6	175 \pm 10.3*	< 0.01	10.0 \pm 2.9	15.5 \pm 2.4*	< 0.01
3 min (n=11)	165 \pm 10.9	180 \pm 9.2*	< 0.01	13.0 \pm 3.3	16.9 \pm 2.1*	< 0.01
4 min (n=10)	170 \pm 10.8	183 \pm 9.2*	< 0.01	14.5 \pm 3.1	17.9 \pm 1.8*	< 0.01
5 min (n=9)	172 \pm 10.7	183 \pm 8.5*	0.01	16.0 \pm 2.7	18.6 \pm 1.7*	0.01
6 min (n=6)	174 \pm 10.3	184 \pm 7.9	0.07	17.0 \pm 2.8	18.8 \pm 1.2*	0.04
7 min (n=4)	177 \pm 9.8	185 \pm 10.5*	0.02	19.0 \pm 2.3	19.8 \pm 0.5	0.06
8 min (n=3)	179 \pm 10.5	190 \pm 2.0	0.07	19.5 \pm 2.2	20.0 \pm 0.0	0.42

Note: AU: arbitrary units; HR: heart rate; RPE: rating of perceived exertion.

* $P < 0.05$ compared with t_{lim0} .

Discussion

This study aimed to compare the two protocols (without and with a 15-minute warm-up) in the t_{lim} determination at 100% of V_{peak} in untrained men. The main finding of the present study was that the different protocols for the t_{lim} determination had different test durations;

in particular, the absence of the 15-minute warm-up period led participants to run for a longer duration than the heating test, confirming our hypothesis. Furthermore, the t_{lim} protocol influenced the HR and RPE during the test responses but did not modify lactate concentrations or the maximal variables (HR_{max} and RPE_{max}).

Previous investigations determined t_{lim} using a standard protocol with a 15-minute warm-up duration at 60% of maximal aerobic speed (MAS), vVO_{2max} or V_{peak} ^{4,5,17}. Other studies used different warm-up durations (t_{lim10} and t_{lim5})^{5,18,19}, but none omitted the warm-up (t_{lim0}). This is the first study to investigate performance in a protocol for determining t_{lim} without a previous warm-up and compared it with the standard protocol (t_{lim15})⁴. Our results demonstrated that the runners participated for a longer duration in the protocol without a warm-up than in the protocol with a warm-up. This result was similar to those of Da Cruz *et al.*⁵ and Bertuzzi *et al.*¹⁸, who used shorter warm-up protocols (*i.e.*, t_{lim10} and t_{lim5}) when testing untrained young men and recreational long-distance runners, respectively. When investigating the influence of different warm-up durations on determining t_{lim} (*i.e.*, t_{lim15} , t_{lim10} , t_{lim5}), Da Cruz *et al.*⁵ reported a longer participation time in the protocol with a shorter warm-up duration for t_{lim} determination. Similarly, Bertuzzi *et al.*¹⁸ observed that the participants remained in the protocol with a 5-minute warm-up longer than they did in the protocol with a 10-minute warm-up. Thus, a longer warm-up time for the t_{lim} determination showed a negative effect on test duration and may be a tiring factor prior to exercise. The longer warm-up duration seems to cause greater physiological wear in the participants at the beginning of the test as demonstrated by the higher values of HR and RPE in the t_{lim15} protocol than in the t_{lim0} protocol observed in the present study.

In addition to warm-up duration, t_{lim} seems to be influenced by participant fitness level since there was an inverse correlation between MAS and t_{lim} ^{4,20,21}. Renoux *et al.*²⁰ reported a mean t_{lim15} value of 4.5 ± 1.3 minutes in trained runners, similar to that observed by Billat *et al.*⁴ (*i.e.*, 5.01 ± 0.9 minutes) in a study of trained runners that showed a shorter t_{lim} duration than the results of our study for the t_{lim15} protocol (*i.e.*, 6.0 ± 2.0 minutes) as well as the study of Da Cruz *et al.*⁵ (*i.e.*, 5.9 ± 1.7 minutes) that evaluated untrained young males. These results show that it is possible to observe a difference in t_{lim} duration despite use of the same protocol (t_{lim15}) due to differences in participant fitness levels.

Since t_{lim} is a variable that is used to determine the optimal stimulus durations in interval training sessions^{9,11}, this longer duration found by a lack of a warm-up will directly impact this prescription. Previous studies using t_{lim} to prescribe interval training showed that if these sessions do not have the ideal duration, runners may not complete the training and/or show no improvement because of a low stimulus intensity^{10,11,22}. Billat *et al.*¹⁰ tested the training effect in that the stimuli had a duration of 50% of t_{lim} in vVO_{2max} in the interval training sessions, and no differences were found in the aerobic variables associated with performance after 4 weeks of training. Similarly, some authors^{11,22} tested different combinations of vVO_{2max} and its respective t_{lim} for individualized training with series of 60 and 75%; 60 and 70% of t_{lim} in vVO_{2max} , respectively. As a result, performance improvements were observed only for the groups that trained with the series duration at 60% of t_{lim} in vVO_{2max} . The results of these studies suggest that small changes in t_{lim} duration can have a great impact on the training prescription. Thus, the correct choice of the

protocol for determining the t_{lim} is important because the warm-up time (its lack or presence) will directly affect the t_{lim} duration and, therefore, the training prescription.

Regarding HR, RPE, and lactate concentration responses during the test, the protocols with different warm-up durations (*i.e.*, t_{lim0} and t_{lim15}) were not expected to affect the maximal variables (HR_{max}, RPE_{max}, and LA_{peak}). However, when during the test HR and RPE responses were analyzed during the different warm-up times, we obtained smaller values in the t_{lim0} protocol compared to the t_{lim15} protocol. A similar result was found by Da Cruz *et al.*⁵, who observed higher values for these variables in the protocol with a 15-minute warm-up compared to protocols with 5- and 10-minute warm-ups. This change can be explained by the increase in cardiovascular activity after exercises with durations > 10 minutes caused by changes in thermoregulatory mechanisms, energy substrates, and increased blood flow^{23,24}, which also affected the participants final performances.

Despite the important findings of our study, one limitation was that the determination of the V_{peak} (and warm-up based on V_{peak}) could lead to different individual intensities relative to the velocity of anaerobic threshold (vAT). Thus, despite of V_{peak} and vAT are correlated with endurance running performance^{12,25} no previous study examined the relationship between V_{peak} and vAT. Future studies should investigate the relationship between V_{peak} and vAT and to better understand how differences in relative intensities of vAT could affect the time limit performance.

Conclusion

Therefore, we conclude that the lack of a 15-minute warm-up, based on the protocol proposed by Billat *et al.*⁴, for determining t_{lim} leads to a longer test duration at 100% of V_{peak} and modifies the responses during the test variables (HR, RPE) in untrained men. This result may impact or interfere with the use of t_{lim} for interval training prescriptions. Hence, we suggest that further studies should be performed in training protocols with different warm-up durations to evaluate the impact of the use of time limits.

Conflict of interest

The authors do not declare a conflict of interest.

Bibliography

1. Salgado JVV, Chacon-Mikahil MPT. Corrida de rua: análise do crescimento do número de provas e de praticantes. *Conexões: Educ. Fís., Esporte e Saúde*. 2006;4:90-9.
2. Nakamura FY, Moreira A, Aoki MS. Monitoramento da carga de treinamento: a percepção subjetiva do esforço da sessão é um método confiável? *Rev. Educ. Fís/UEM*. 2010;21:1-11.
3. Midgley AW, McNaughton LR, Jones AM. Training to enhance the physiological determinants of long-distance running performance. *Sports Med*. 2007;37:857-80.
4. Billat VL, Renoux JC, Pinoteau J, Petit B, Koralsztejn JP. Times to exhaustion at 100% of velocity at VO_{2max} and modeling of the time-limit/velocity relationship in elite long-distance runners. *Eur J Appl Physiol*. 1994;69:271-3.
5. Da Cruz VHM, Peserico CS, Machado FA. Effect of prior warm-up duration on the time limit at peak speed in untrained men. *J Sports Med Phys Fitness*. 2017;57:1276-81.
6. Wallmann HW, Christensen SD, Perry C, Hoover DL. The acute effects of various types of stretching (static, dynamic, ballistic, and no stretch) of the iliopsoas on 40-yard sprint times in non-athletes. *Int J Sports Phys Ther*. 2012;7:540-7.

7. Donahue RB, Vingren JL, Duplanty AA, Levitt DE, Luk HY, Kraemer WJ. Acute Effect of Whole-Body Vibration Warm-up on Footspeed Quickness. *J Strength Cond Res.* 2016;30:2286-91.
8. Tomaras EK, Macintosh BR. Less is more: standard warm-up causes fatigue and less warm-up permits greater cycling power output. *J Appl Physiol.* 2011;111:228-35.
9. Manoel FA, da Silva DF, de Lima JRP, Machado FA. Peak velocity and its time limit are as good as the velocity associated with VO_{2max} for training prescription in runners. *Sports Med Int Open.* 2017;1:E8-E15.
10. Billat VL, Flechet B, Petit B, Muriaux G, Koralsztein JP. Interval training at VO_{2max} : effects on aerobic performance and overtraining markers. *Med Sci Sports Exerc.* 1999;31:156-63.
11. Smith TP, Mcnaughton LR, Marshall KJ. Effects of 4-wk training using V_{max}/T_{max} on VO_{2max} and performance in athletes. *Med Sci Sports Exerc.* 1999;31:892-6.
12. Machado FA, Kravchychyn AC, Peserico CS, da Silva DF, Mezzaroba PV. Incremental test design, peak "aerobic" running speed and endurance performance in runners. *J Sci Med Sport.* 2013;16:577-82.
13. Kuipers H, Rietjens G, Verstappen F, Schoenmakers H, Hofman G. Effects of stage duration in incremental running tests on physiological variables. *Int J Sports Med.* 2003;24:486-91.
14. Astrand PO. *Experimental studies of physical working capacity in relation to sex and age.* Copenhagen: Ejnar Munksgaard. 1952; p. 23-27, 92-102.
15. Tanaka H, Monahan KD, Seals DR. Age-predicted maximal heart rate revisited. *J Am Coll Cardiol.* 2001;37:153-6.
16. Borg GA. Psychophysical bases of perceived exertion. *Med Sci Sports Exerc.* 1982;14:377-81.
17. Da Silva DF, Simões HG, Machado FA. vVO_{2max} versus V_{peak} , what is the best predictor of running performances in middle-aged recreationally-trained runners? *Sci Sports.* 2015;30: e85-e92.
18. Bertuzzi RC, Bueno S, Pasqua LA, Acquesta FM, Batista MB, Roschel H, *et al.* Bioenergetics and Neuromuscular Determinants of the Time to Exhaustion at Velocity Corresponding to VO_{2max} in Recreational Long-Distance Runners. *J Strength Cond Res.* 2012;26:2096-102.
19. Caputo F, Denadai B. Resposta do VO_2 e tempo de exaustão durante a corrida realizada na velocidade associada ao VO_{2max} : aplicações para o treinamento aeróbio de alta intensidade. *Rev Bras Cienc Esporte.* 2004;26:19-31.
20. Renoux JC, Petiti B, Billat V, Koralsztein. Oxygen Deficit is Related to the Exercise Time to Exhaustion at Maximal Aerobic Speed in Middle Distance Runners. *Arch Physiol Biochem.* 1999;107:280-5.
21. Kachouri M, Vandewalle H, Huet M, Thomaidis M, Jousselin E, Monod H. Is the exhaustion time at maximal aerobic speed an index of aerobic endurance? *Arch Physiol Biochem.* 1996;104:330-6.
22. Smith TP, Coombes JS, Geraghty DP. Optimising high-intensity treadmill training using the running speed at maximal O_2 uptake and the time for which this can be maintained. *Eur J Appl Physiol.* 2003;89:337-43.
23. Coyle EF. Cardiovascular drift during prolonged exercise and the effects of dehydration. *Int J Sports Med.* 1998;19:121-4.
24. Buchfuhrer MJ, Hansen JE, Robinson TE, Suedy DY, Wasserman MJ, Whipp BJ. Optimizing the exercise protocol for cardiopulmonary assessment. *J Appl Physiol.* 1983;55:1558-64.
25. Machado FA, de Moraes SM, Peserico CS, Mezzaroba PV, Higino WP. The D_{max} is highly related to performance in middle-aged females. *Int J Sports Med.* 2011;32:672-6.

Acute effects of resistance and concurrent exercise on the lipid profile of postmenopausal women

Roberto Rebolledo-Cobos¹, Andrés Peñaloza Florez¹, Luz Sarmiento-Rubiano¹, Laura Ardila-Pereira², Jimmy Becerra Enriquez¹, Adel Rodríguez-Delgado³

¹Universidad Metropolitana de Barranquilla, Colombia. ²Universidad de Santander, Valledupar, Colombia. ³Universidad Autónoma del Caribe, Barranquilla, Colombia.

Received: 24.05.2018
Accepted: 26.06.2018

Summary

The postmenopausal condition is characterized by a systematic increase in cardiovascular risk factors, including negative alterations in lipid metabolism. Physical exercise has shown beneficial effects in the regulation of lipemic markers in different populations; however, its effects are not well understood in postmenopausal women. The aim of the study was to determine the effects of resisted exercises (RE), concurrent exercises (CE), compared with a control group (CG) on lipid profile in postmenopausal women. Thirty-two voluntary postmenopausal women were randomly assigned into three groups; one group performed a RE session (n = 11), other group a CE session (n = 11) and control group performed flexibility exercises (n = 10). The basal lipid profile, body composition, muscular fitness were evaluated and the basal metabolic rate (BMR) was calculated before the study. The lipid profile was evaluated before, immediately after and 12 hours after the specific exercise session. The results showed that there was no a statistical significant difference among all groups in plasma concentration of total cholesterol, triglycerides and LDL, in any of the evaluation moments ($p > 0.05$). After 12 hours of exercise session, the CE group decreased more VLDL concentrations than the RE group and control group ($p < 0.05$). The CE group increased significantly HDL concentrations in relation to the ER group ($p < 0.05$). Therefore, it can be concluded that CE had more favorable acute effects on lipid metabolism than ER, prolonging its effect up to 12 hours after being performed in overweight postmenopausal women.

Key words:

Postmenopause. Exercise.
Resistance training.
Muscle strength (MeSH).

Efectos agudos del ejercicio resistido y concurrente en el perfil lipídico de mujeres postmenopáusicas

Resumen

El estado postmenopáusico se caracteriza por el aumento sistemático de los factores de riesgo cardiovascular, incluyendo las alteraciones negativas en metabolismo de lípidos. El ejercicio físico ha demostrado efectos benéficos en la regulación de marcadores lipémicos en diferentes poblaciones, sin embargo, en mujeres postmenopáusicas no están bien dilucidados sus efectos. El objetivo del estudio fue determinar el efecto que tiene sobre el perfil lipídico de mujeres postmenopáusicas la realización de una sesión de ejercicios resistidos (ER) o de ejercicios concurrentes (EC), comparados con un grupo control (GC). La metodología del estudio consto de una división aleatoria en tres grupos de 32 mujeres posmenopáusicas voluntarias, un grupo realizó una sesión de ER (n=11), otro grupo una sesión de EC (n=11) y un grupo control realizó ejercicios de flexibilidad (n=10). Inicialmente fue evaluado el perfil lipídico basal, la composición corporal, la *fitness* muscular y se calculó la tasa metabólica basal (TMB). Fue evaluado el perfil lipídico antes, inmediatamente después y 12 horas después de la sesión de ejercicio específica. Como resultados se encontró que entre los tres grupos, no existieron diferencias significativas en las concentraciones plasmáticas de colesterol total, triglicéridos y LDL en ninguno de los momentos de evaluación ($p > 0.05$). El grupo EC disminuyó las concentraciones de VLDL 12 horas después de la sesión de ejercicios, en comparación al grupo ER y GC ($p < 0.05$); y aumentó significativamente las concentraciones de HDL con relación al grupo ER ($p < 0.05$). De este estudio se puede concluir que en las mujeres postmenopáusicas con sobrepeso participantes, el EC puede tener efectos agudos más favorables en el metabolismo de lípidos que el ER, prolongando su efecto hasta 12 horas después de realizarse.

Palabras clave:

Postmenopausia. Ejercicio físico.
Entrenamiento de resistencia.
Fuerza muscular (MeSH).

Note: The financial support of the Metropolitan University of Barranquilla was provided to carry out this bibliographical study.

Correspondence: Roberto Carlos Rebolledo Cobos
E-mail: rrebolledo@unimetro.edu.co

Introduction

High lipid concentrations are a significant risk factor for the development of cardiovascular diseases (CVD), given that they commonly precede endothelial dysfunction, playing a key role in the formation of atherosclerotic plaque and the subsequent inflammatory response¹. These phenomena are associated with an increase in the secretion of cytokines, the precursors of inflammation, the expression of adhesion molecules (ICAM, VCAM-1), fibrinogen and the activity of oxidation-promoting substances².

Physical exercise has been shown to have beneficial effects on the regulation of lipemia markers in different populations, associating the total energy expenditure during exercise as the main determinant of the subsequent metabolic phenomena^{3,4}. The performance of an aerobic exercise session can reduce the lipid plasma concentrations associated with atherosclerosis pathogenesis, such as low-density lipoproteins (LDL) and very low-density lipoproteins (VLDL)⁵, as well as increasing the activity of high-density lipoproteins (HDL) which play a significant role in the systemic regulation of excess cholesterol⁶. More recent studies have reported favourable changes in the lipid profile after a single aerobic exercise session^{7,8}.

On the other hand, muscle strengthening exercises with external resistances or resistance exercises (RE) are primarily associated with responses in the functional sense of anaerobic resistance and strength⁹. RE are classed as a means for the prevention of chronic diseases and acute catastrophic events, such as infarcts or cerebrovascular ischemia¹⁰. The preventive role of RE could be due to their impact on the assimilation of lipids and the regulation of the pro- and anti-inflammatory markers on the organism, during, and primarily after exercising^{11,12}. Only a few studies have assessed how an RE session modifies the lipid markers in the blood stream.

The combination of RE and aerobic exercises, currently known as concurrent exercises (CE), has been shown to have beneficial effects on the state of health and general physical condition if performed regularly¹³. However, benefits have been observed, as well as improvements in the lipid profile and general physical fitness in vulnerable populations, such as persons with obesity and postmenopausal women¹⁴.

After the menopause, the risk of women developing CVD increases⁴. This increased vulnerability is related to the reduction in oestrogen levels, a hormone that appears to exercise a protective role against CVD in women, by promoting an atherogenic lipid profile, an anti-inflammatory immunological profile and by direct action on the blood vessel walls, thereby avoiding the dysfunction of the vascular endothelium¹⁵. The associated physiological, metabolic and morphological changes are characterised by a fall in the level of physical activity, increased body mass index (BMI), a decrease in muscle mass (sarcopenia), a decrease in the basal metabolic rate, an increase in triglyceride concentrations (TG) and LDL¹². The inclusion of physical exercise in the life style of this population, may have a positive effect on a number of cardiovascular risk factors, such as a reduction in body fat and the acute regulation of some of the lipemia markers⁶.

This study is directed at determining the acute effect of an RE session and a CE session on the lipid profile of postmenopausal women, compared to a control group.

Material and method

This investigation corresponds to a quasi-experimental study with single blind randomisation, exposure-response based on the acute changes provided in two experimental groups, two types of physical exercise with specific biochemical blood markers, compared to a control group.

Subjects

The sample comprised a total of 32 postmenopausal women volunteers, workers from the Fundación Hospital Universitario Metropolitano in the city of Barranquilla, with an average age of 54.66 ± 4.04 years, recruited through the occupational welfare office. The inclusion criteria were postmenopausal women with no hormone replacement treatment. Excluded from the study were those women who had taken part in some type of resistance training during the 6 months prior to the study and those with a medical history of serious endocrine, metabolic and/or neuromuscular disorders, determined during screening by a doctor specialising in internal medicine. Before taking part, each of the subjects of the study was carefully informed of the study design, particularly the possible risks and discomforts related to the procedures, subsequently giving their written informed consent.

Ethical considerations

This study was conducted in accordance with the ethical standards of exercise sciences¹⁶, of the World Medical Association and of the Declaration of Helsinki. The protocol was approved by the Institutional Investigation and Ethics Committee of the Universidad Metropolitana de Barranquilla, Colombia.

Experimental design

The total number of study subjects ($n=32$) were randomly divided into 3 groups, two experimental groups who performed a session of resistance exercises (RE, $n=11$) and a session of concurrent exercises (CE, $n=11$), respectively; and a control group with no exercises with significant energy expenditure (CG, $n=10$).

Prior to the experimental protocol, an appointment was made for the women for an initial assessment of the musculoskeletal variables (maximum dynamic force and lower-limb functional capacity), body composition (% of fat and lean mass), anthropometric assessment (height, weight, waist circumference), basal metabolic rate and monitoring of dietary behaviour based on the protocols described below.

Maximum dynamic force

In order to determine the maximum dynamic force of the women's different body segments, maximum repetition tests (1RM) were used

for the muscle groups performing the resistance exercises on which the experimental groups were based. The maximum load for each subject was determined with no more than 5 tries with a 4-minute recovery between tries.

Maximum dynamic force

In order to determine the maximum dynamic force of the different body segments of the women, tests of a maximum repetition (1RM) were used for the muscular groups that perform the resisted exercises on which the experimental groups are based. The maximum load of each subject was determined with no more than 5 attempts with a recovery of 4 minutes between attempts.

Assessment of the lower-limb functional capacity

The 30 second sit-to-stand test measures the functional capacity of the lower limbs. The test was performed as follows: the initial position was seated in the centre of a chair (height 43 cm), with the back straight and the feet placed flat on the floor, shoulder width apart, the arms crossed at chest height, with an approximate angle of 90° flexion of the hip and knee. At a verbal signal, the participants stand up and then return to the initial seated position. Participants were encouraged to repeat this as many times as possible within a 30 second period.

Anthropometric assessment, body composition and basal metabolic rate

The anthropometric assessment was made of the women, determining height, weight, waist circumference, using a digital scale, stadiometer and tape measure, SECA brand, according to the instructions set out in resolution 2465 of the Ministry of Health of Colombia¹⁷. The data were used to calculate the BMI. Using the bioelectrical middle ear analyser, Tanita brand (TBF-300WA Wrestling Body Composition Analyzer) the body composition was determined, calculating the percentage of body fat and lean mass. The Basal Metabolic Rate (BMR) was calculated with the Harris-Benedict formula, modified with the physical activity factor (+20%). $BMR = [655.0955 + (9.5634 \times \text{weight in kg}) + (1.8449 \times \text{height in cm}) - (4.6756 \times \text{age in years})]$.

Monitoring and guiding nutritional behaviour

On the day of the initial assessment, the study subjects were given a printed document and, prior to a detailed explanation by a nutritionist, they recorded their nutritional behaviour during the 3 days prior to performing the experimental protocol. This was for the purpose of compensating for any improbable lipid profile results, derived from consuming excessively high fat food. The aim was to reduce alterations in the results obtained, derived from atypical dietary behaviour.

On the day of the experimental protocol, the food intake was monitored, standardising the calorie consumption prior to doing the exercises to 60% of the BMR with a physical activity factor for each case. The strategy included breakfast (cereals and fruit), snack (cereals) and lunch (chicken,

potatoes and vegetables). All the food was prepared, proportioned and monitored by a nutritionist. The aim of this procedure was to guarantee that, when performing the exercise session, the lipid concentrations were influenced by similar nutritional conducts and in specific quantities for each individual case. The last food intake for each case was programmed at least 3 hours prior to the exercises, encouraging participants to keep hydrated just by drinking water.

Baseline lipid profile

All subjects underwent this procedure one day prior to doing the exercises and after at least a 12-hour fast. The women in the groups went to the facilities of the Fundación Hospital Universitario Metropolitano in the morning, to have 5ml of blood drawn in order to determine the baseline lipid profile, which includes the analysis of the serum concentrations of Total Cholesterol (TC), HDL, LDL, VLDL and TG.

Experimental protocol

In a maximum period of one week following the muscular fitness assessment, another appointment was scheduled with the study subjects, at 5:00 pm at the gym facilities of the Physiotherapy Department of the Fundación Hospital Universitario Metropolitano. On the subjects' arrival, the nutritionist initially authorised the exercise session for those women with no atypical eating disorders that could interfere with the results and who correctly followed the dietary procedures, based on the review of the nutrition monitoring document.

Blood samples were drawn from the women by a bacteriological expert, for the assessment of the lipid profile before the exercise session, and then immediately after it ended and 12 hours later, the following morning (under fasting conditions). Immediately after being drawn, the blood samples were taken to the laboratory for analysis. The exercise sessions for each experimental group and the control group are described below:

Exercise protocol

The subjects of the RE group did a 10 minute warm-up consisting in walking (5 min) and general joint mobilization (5 min). Following this, they performed a total of 8 resistance exercises (1. Leg Press; 2. Machine knee extension; 3 Machine knee flexion; 4 Elbow flexion with dumbbells; 5 Elbow extension with dumbbells; 6 Shoulder adduction with dumbbells from abduction in a supine position; 7. Unilateral dumbbell row; and 8. Push-ups with free weight), each one with 3 sets and each set with 15 reps, with an intensity of 75% of MR1. The approximate total duration of this session was 40 minutes.

The CE group initially performed a 10 minute warm-up consisting in walking (5 min) and general joint mobilization (5 min), followed by the same exercises as for the RE group, except for the fact that each exercise was only performed for a set with 15 reps, at 75% of MR1. Subsequently, the subjects performed 20 minutes of continuous pedalling on a cycle ergometer at a frequency of between 70 to 80% of the maximum heart rate. The approximate total duration of this exercise session was 40 minutes.

On the implicit instructions of the ethics and scientific committee of the Universidad Metropolitana de Barranquilla, those subjects forming part of the Control Group (CG) ought to do some exercises to promote their physical health, from any point of view. It was therefore decided to get the members of this group to perform a set of passive stretching exercises, which were shown to have no metabolic implications on the lipid metabolism¹⁸. The session comprised a 10 minute warm-up (5 minutes free walk and 5 minutes of joint mobilization) and 30 minutes of directed stretching. The average duration of this session was 40 minutes.

At the end of each exercise session, the subjects were instructed to drink only water with no calorie content, until the following morning when, under fasting conditions, a blood sample was taken corresponding to 12 hours after the exercise session.

Statistical analysis

A descriptive and comparative statistical analysis was performed with measurements of averages with respective standard deviations. The Shapiro-Wilk test of normality was used and the Levene test of homogeneity. Subsequently, the Analysis of Variance Test (ANOVA) was used two-way, intra-group and inter-group. Whenever relevant, the Bonferroni post-hoc test was considered. The significance level was $p < 0.05$, considering this for all analyses.

Results

The results of the initial assessment of the women showed that the randomisation did not lead to any significant difference between any of the groups and for none of the performance variables before executing the experimental protocol (Table 1). For the total number of

women, the BMI ranged from 25.02 to 37.58, ranging from 25 to 30 in 23 women, greater than 30 for 8 women considered to have type I obesity and only one woman had a BMI of more than 35, considered to have type II obesity. The distribution of women with obesity was balanced in the three groups. The waist circumference value ranged between 84 to 110 cm, with all the women with abdominal obesity coming within the criteria established by resolution 2465 of 201617. The average values of the variables analysed in the initial evaluation of the women for the total population and for each of the experimental and control groups are set out in Table 1.

The statistical comparison of the average values between the experimental groups, of the variables of the baseline lipid profile under fasting conditions and at each of the times established in the experimental protocol (before, immediately after and 12 hours after exercise), showed no significant variations ($p > 0.05$) in the exercise sessions performed by the RE and CE groups in relation to total cholesterol, triglycerides and LDL. The assessment of the VLDL levels was statistically significant and was lower for the women in the CE group 12 hours after the exercise session, compared to the RE and CG groups. After 12 hours post-exercise, the HDL levels were also seen to be statistically higher in the CE group in relation to the RE group (Table 2).

Table 3 shows the within-group variations for each lipid profile component, under fasting conditions, comparing the baseline analysis with that collected 12 hours after the corresponding exercise session. No statistical differences were found for any of the variables in the comparison of averages in relation to times ($p > 0.05$). The average with the greatest variation in the CT analysis was that of the CG, dropping 10.1%, followed by the RE group and CE with drops of 9.8% and 3.5% respectively. The CE group showed a greater variation in the TG analy-

Table 1. Characteristics of the subjects, members of the different groups.

Variable	Total population (n=32)	Experimental groups			P ³ value
		RE (n=11)	CE (n=11)	CG (n=10)	
Age (years)	54.65±4.04	54.82±3.46	54.63±4.59	54.50±4.43	0.98
Anthropometric characteristics					
Weight (kg)	71.31±9.29	70.31±9.44	71.88±10.32	71.80±8.85	0.91
BMI (kg/m ²)	28.52±3.33	27.76±3.28	29.03±3.70	28.82±3.16	0.64
Body fat (%)	36.73±4.68	36.79±4.05	37.04±4.39	36.34±5.97	0.94
Lean mass (%)	44.20±5.69	44.48±3.58	42.66±8.45	45.60±3.58	0.50
Waist circumference	96.23±7.61	94.88±7.56	97.09±8.63	96.78±7.28	0.77
Strength and functional capacity					
1RM knee extension (kg)	22.78± 4.57	22.36±4.57	23.45±4.20	22.50±5.34	0.84
1RM elbow flexion (kg)	11.15± 3.69	11.45±4.11	10.36±2.50	11.70±4.47	0.68
30" sit-to-stand test (reps)	10.71±2.54	10.55±3.14	10.36±2.38	11.30±2.11	0.68
Energy requirements (Kcal)					
BMR	1374.4±104.26	1364.32±94.76	1374.95±116.11	1384.98±110.61	0.90
BMRAF	1649.3±125.11	1637.19±113.71	1649.92±139.33	1661.97±132.73	0.90

Average values of the variables analysed in the initial evaluation of the women for the total population and for each of the experimental groups. Value P³ shows the statistical significance when comparing the average values of the variables in the three experimental groups. BMR: basal metabolic rate; BMRAF: basal metabolic rate with physical activity factor.

Table 2. Inter-group comparison of the lipid profile for fasting baseline, before, immediately after and 12 hours fasting after the exercise sessions.

Lipid profile (mg/dL)	Experimental groups			P ³ value
	RE (n=11)	CE (n=11)	CG (n=10)	
TC				
12 hours fasting basal	214.36±21.86	217.52±33.05	217.10±38.09	0.93
Pre exercise	228.45±73.34	220.18±68.08	222.00±70.16	0.95
Immediately post-exercise	244.09±66.86	249.27±67.39	213.10±59.38	0.40
12 hours fasting post-exercise	193.18±28.18	209.81±33.80	195.20±35.61	0.43
TG				
12 hours fasting basal	171.09±71.73	167.36±44.12	172.00±51.76	0.98
Pre exercise	226.27±96.56	204.63±87.56	222.80±87.21	0.41
Immediately post-exercise	252.21±102.21	198.09±82.62	250.81±104.09	0.54
12 hours fasting post-exercise	166.18±63.93	144.00±29.50	175.50±54.90	0.36
LDL				
12 hours fasting basal	135.95±15.59	140.63±18.60	137.35±32.48	0.88
Pre exercise	131.50±54.36	126.87±49.33	119.05±52.36	0.85
Immediately post-exercise	137.61±49.15	148.65±50.12	105.00±48.63	0.12
12 hours fasting post-exercise	140.46±39.22	155.94±47.84	130.40±23.06	0.32
LDL				
12 hours fasting basal	31.05±13.84	30.04±9.55	32.02±14.26	0.92
Pre exercise	45.25±19.31	40.92±15.38	44.56±20.72	0.84
Immediately post-exercise	50.45±29.74	39.61±16.45	50.16±29.86	0.54
12 hours fasting post-exercise	33.23±12.78 ^a	23.34±5.50 ^b	37.10±20.80 ^a	<0.05
HDL				
12 hours fasting basal	42.59±7.19	48.76±6.53	46.13±8.24	0.15
Pre exercise	51.70±11.30	52.38±10.36	58.39±21.18	0.53
Immediately post-exercise	56.02±9.12	61.05±7.99	57.94±16.03	0.59
12 hours fasting post-exercise	48.42±9.05 ^a	56.90±7.40 ^b	52.71±8.43 ^{ab}	<0.05

Average values by group for TC: total cholesterol; TG: triglycerides; HDL: high density lipoproteins; LDL: low density lipoproteins; VLDL: very low density lipoproteins; baseline, before, immediately after and 12 hours after exercise. The comparison of the averages is significant between the three groups when P³<0.05. The averages in the lines that do not share the same superscript letter (a, b) are statistically different.

Table 3. Intra-group variations of the lipid profile components when fasting compared to physical exercise.

Variable	Basal	Post exercise	Δ%	P-value
Total cholesterol (mg/dL)				
RE	214.36±21.86	193.18±28.18	-9.8	0.06
CE	217.52±33.05	209.81±33.80	-3.5	0.59
CG	217.10±38.09	195.20±35.61	-10.1	0.16
Triglycerides (mg/dL)				
RE	171.09±71.73	166.18±63.93	-2.9	0.86
CE	167.36±44.12	144.00±29.50	-13.9	0.15
CG	172.00±51.76	175.50±54.90	+2.0	0.87
LDL (mg/dL)				
RE	135.95±15.59	140.46±39.22	+3.3	0.72
CE	140.63±18.60	155.94±47.84	+10.9	0.19
CG	137.35±32.48	130.40±23.06	-5.0	0.58
VLDL (mg/dL)				
RE	31.05±13.84	33.23±12.78	+7.0	0.70
CE	30.04±9.55	23.34±5.50	-22.3	0.39
CG	32.02±14.26	37.10±20.80	+15.8	0.53
HDL (mg/dL)				
RE	42.59±7.19	48.42±9.05	+13.6	0.13
CE	48.76±6.53	56.90±7.40	+16.7	0.13
CG	46.13±8.24	52.71±8.43	+14.26	0.06

Average values and proportion of variation of the lipid profile by groups. HDL: high density lipoproteins; LDL: low density lipoproteins; VLDL: very low density lipoproteins; baseline, before, immediately after and 12 hours after exercise.

ses, dropping 13.9% compared to 2.9% for the RE group while the CG increased 2.0%. For the RE and CE groups, the LDL showed a tendency to increase, while it dropped in the CG by 5.0%. Only the CE group showed lower VLDL plasma levels in the post-exercise period (22.3%), with the levels for this variable increasing in the RE group (7.0%) and CG (15.8%). The HDL trend in the post-exercise analysis was to increase for all groups.

Discussion

For each individual, the lipid metabolism at the systemic level is dependent on many factors, including diet, the physical health condition and the daily energy expenditure. This study aimed to demonstrate the influence of the different types of physical exercise considered, in a population of postmenopausal women with similar characteristics by comparing the results of the assessment times (baseline, pre-exercise, immediately post-exercise and 12 hours post-exercise) between the experimental groups. In addition, it attempted to show the possible influence of physical exercise on the variation in the lipid profile under fasting conditions within each group. Although it is clear that the influence of physical exercise is not the only determining factor in the lipid profile variations, given that, depending on time, the general energy shortage and the physiological replenishment mechanisms vary from subject to subject. However, it is extremely important to clarify that, although there are limitations with regard to the control of the biological and behavioural factors that influence the lipid profile, additional energy expenditure involved in exercising, influences the magnitude of the physiological responses regulating the lipid metabolism and other energy substrates.

The analyses of the results of this study are based on the acute responses of the types, levels and intensities of physical exercise considered on the lipids found in the blood plasma. These responses have not been fully clarified for postmenopausal women, a population that is relatively prone to suffering CVD. The interpretation of responses at times that are subsequent to the scheduled performance of physical exercise is extremely important given that it makes it possible to build a useful description of the physiology of the human body, endeavouring to support the notion that argues the benefits of physical exercise not only when it is being performed but also subsequently, during a period of physiological recovery and reorganisation.

Findings in the literature associate regular physical exercise with changes in the metabolic and functional variables, primarily in combinations of resistance exercises with aerobic exercises^{9,19}. In order to bring about changes in the BMI, weight and percentage of body fat, it is necessary to implement long-term exercise programs. This is a limiting factor in this study, given that a single exercise session does not produce changes in body composition. However, it has been observed that a single exercise session can modify the lipid profile of young people, adults and subjects with various disorders.

The systematic implementation of RE for more than 8 weeks, has shown findings associated with the reduction in serum concentrations of TC, TG and LDL, as well as increased concentrations of HDL²⁰, although

results from other studies differ, such as the study by Kelley *et al.*²¹, in which it is alleged that in different RE periods (for example, between 8 to 20 weeks) there are no changes in the lipoprotein concentrations in the blood plasma.

With regard to the acute effects of one exercise session, these effects are primarily associated with increased HDL concentrations. In a study on healthy adult men, Wallace *et al.*²² reported a 12% increase in HDL plasma concentrations within the first 24 hours following a high volume and moderate intensity session of RE (7 exercises of 3 sets and 12 reps at 80% of the maximum strength), as well finding a 20% reduction in TG concentrations for the same period. Very similar results were found in this present study, a 13.6% increase in HDL was recorded for the RE group in relation to its baseline records, not finding any great changes in the TG levels 12 hours after exercise.

Compared to the control group, the RE group subjects showed no differences in modifications of LDL and HDL, refuting the findings of the study made by Correa *et al.*²³ in which high or low volume resistance exercises did not reduce the TC concentrations in postmenopausal women, neither did they affect the LDL and VLDL levels, both with regard to baseline values and the postprandial analysis. In the population studied, these results would associate resistance exercises with weak acute responses with regard to the lipid metabolism. However, more evidence is needed to support this claim, particularly taking account of the fact that the variability in the prescription of resistance exercises with regard to volume, intensity, number of exercises, groups of muscles, recovery intervals and speed of contraction, may show divergent results for alternative combinations to those considered in this study.

A marked trend in the lipid profile results before and immediately after the exercise session was that none of the variables showed an intra-group statistical difference. However, 12 hours after exercise, the group that combined aerobic and resistance exercises showed differences with the RE group and control group, with a significant reduction in VLDL of approximately 22% in relation to the baseline records during fasting for the same group. It appears that aerobic exercises have a better impact on the energy metabolism of postmenopausal than just doing resistance exercises. For this population, Weise *et al.*⁵ reported that 12 hours after performing a session of aerobic exercises, the TG concentrations could be reduced by 8.5% while the HDL concentrations could be increased by 5%. In this study, the scope of the variation in TG and HDL within the CE group showed better results than in the aforementioned study, with a 13% reduction in TG and an increase in HDL up to 16.7%, 12 hours after performing the exercises. This finding may be an indicator that the combination of aerobic and anaerobic exercises have a better impact on the acute lipid metabolism of postmenopausal women.

The above results may suggest that high levels of resistance exercises do not stimulate the formation of HDL, due to the fact that the reverse mobilisation of lipids (towards the liver) is reduced by the energy demands of the exercised skeletal muscle, which requires the typical mobilisation of fatty acids. These are very similar results to those reported in the studies by Wooten *et al.*²⁴ and Zotou *et al.*²⁵, where they demonstrated that, although resistance exercises exert a positive influence on

the acute mitigation of the baseline and postprandial lipid markers for postmenopausal women, the high volumes of these exercises did not show positive modifications in the HDL blood concentrations.

There are many authors who have not only mentioned a reduction in the levels of physical activity as a crucial factor in increasing the cardiovascular risk in postmenopausal women, but also find physiological aggravating factors that can be controlled with periodical physical exercise, particularly in overweight women. According to the American College of Sports Medicine, the regular performance of resistance exercises may improve the overall physical health of women, and help prevent and treat cardiovascular risk related disorders, such as diabetes or high blood pressure. Therefore, these exercises, monitored by a specialist, are currently part of rehabilitation and physical fitness programs, directed at this population²⁶. There is little evidence-based literature that describes the acute metabolic implications of concurrent exercises on the lipid profile of postmenopausal women, although studies such as Figueroa *et al.*¹⁹ and Libardi *et al.*²⁷ show positive effects on the control of cardiovascular risk factors when this type of exercise is performed in the long term. Based on the most important findings of this study, it is possible to start to consider the idea of the combined performance of resistance and aerobic exercises, in order to guarantee positive acute influences on the blood lipid profile of postmenopausal women, which, in the long term, could become a protective factor against suffering CVD.

Based on the analysis of the results of this study, we can conclude that, in comparison to the performance of a session of resistance exercises, a session of concurrent exercises could positively affect the lipid profile of overweight postmenopausal women, principally 12 hours after doing the exercises, reducing the VLDL plasma concentrations and increasing the HDL concentration.

Conflict of interest

The authors do not declare a conflict of interest.

Bibliography

- Pirillo A, Norata G, Catapano A. Postprandial lipemia as a cardiometabolic risk factor. *Curr Med Res Opin.* 2014;30(8):1489-503.
- Chan D, Pang J, Romic G, Watts G. Postprandial hypertriglyceridemia and cardiovascular disease: current and future therapies. *Curr Atheroscler Rep.* 2013;15(3):309-13.
- Wooten J, Phillips M, Mitchell J, Patrizi R, Pleasant R, Hein R, *et al.* Resistance exercise and lipoproteins in postmenopausal women. *Int J Sports Med.* 2011;32(1):7-13.
- Tibana R, Pereira G, De Souza J, Tajra V, Vieira D, Campbell C, *et al.* Resistance training decreases 24 hour blood pressure in women with metabolic syndrome. *Diabetol Metab Syndr.* 2013;5(1):27-32.
- Weise S, Grandjean P, Rohack J, Womack J, Crouse S. Acute changes in blood lipids and enzymes in postmenopausal women after exercise. *J Appl Physiol.* 2005;99(2):609-15.
- Gilmore L, Crouse S, Carbuhn A, Klooster J, Calles J, Meade T, *et al.* Exercise attenuates the increase in plasma monounsaturated fatty acids and high-density lipoprotein cholesterol but not high-density lipoprotein 2b cholesterol caused by high-oleic ground beef in women. *Nutr Res.* 2013;33(12):1003-11.
- Comassi M, Vitolo E, Pratali L, Del Turco S, Dellanocce C, Rossi C, *et al.* Acute effects of different degrees of ultra-endurance exercise on systemic inflammatory responses. *Intern Med J.* 2015;45(1):74-9.
- Gonçalves V, Julio U, Diniz T, de Moura B, Lira F, Takito M, *et al.* Postprandial lipoprotein profile in two modes of high-intensity intermittent exercise. *J Exerc Rehabil.* 2016;12(5):476-82.
- Rebolledo-Cobos R, Correa C, Reischak-Oliveira A. Respuesta metabólica y adaptaciones musculares de mujeres posmenopáusicas al entrenamiento resistido de alto y bajo volumen. *Rev Mov Cient.* 2014;8(1):8-17.
- Agrinier N, Cournot M, Dallongeville J, Arveiler D, Ducimetière P, Ruidavets J, *et al.* Menopause and modifiable coronary heart disease risk factors: a population based study. *Maturitas.* 2010;65(3):237-43.
- Elliott K, Sale C, Cable N. Effects of resistance training and detraining on muscle strength and blood lipid profiles in postmenopausal women. *Br J Sports Med.* 2002;36(5):340-4.
- Moreau K, Deane K, Meditz A, Kohrt W. Tumor necrosis factor- α inhibition improves endothelial function and decreases arterial stiffness in estrogen-deficient postmenopausal women. *Atherosclerosis.* 2013;230(2):390-6.
- Spence A, Carter H, Naylor L, Green D. A prospective randomized longitudinal study involving 6 months of endurance or resistance exercise. Conduit artery adaptation in humans. *J Physiol.* 2013;591(5):1265-75.
- Ho S, Dhaliwal S, Hills A, Pal S. The effect of 12 weeks of aerobic, resistance or combination exercise training on cardiovascular risk factors in the overweight and obese in a randomized trial. *BMC Public Health.* 2014;12(704):2-10.
- Costa R, Lima A, Tagliari M, Krueger L. Effects of resistance training on the lipid profile in obese women. *J Sports Med Phys Fitness.* 2011;51(1):169-77.
- Harris D, Atkinson G. Ethical standards in sport and exercise science research. *Int J Sports Med.* 2011;32(12):819-21.
- Ministerio de Salud y Protección Social de Colombia. Resolución 2465 de junio 14. 2016. (Consultado 06/07/2017) Disponible en: <https://ids.gov.co/web/2016/resoluciones/2465.pdf>
- Lewis-Frank L, Sorensen B, Yasui Y, Tworoger S, Schwartz R, Ulrich C, *et al.* Effects of exercise on metabolic risk variables in overweight postmenopausal women: a randomized clinical trial. *Obes Res.* 2005;13(3):615-25.
- Figueroa A, Park S, Seo D, Sanchez-Gonzalez M, Baek Y. Combined resistance and endurance exercise training improves arterial stiffness, blood pressure, and muscle strength in postmenopausal women. *Menopause.* 2011;18(9):980-4.
- Campbell W, Haub M, Wolfe R, Ferrando A, Sullivan D, Apolzan J, *et al.* Resistance training preserves fat-free mass without impacting changes in protein metabolism after weight loss in older women. *Obesity (Silver Spring).* 2009;17(7):1332-9.
- Kelley G, Kelley K. Impact of progressive resistance training on lipids and lipoproteins in adults: a meta-analysis of randomized controlled trials. *Am J Prev Med.* 2009;48(1):9-19.
- Wallace M, Moffatt R, Haymes E, Green N. Acute effects of resistance exercise on parameters of lipoprotein metabolism. *Med Sci Sports Exerc.* 1991;23(2):199-204.
- Correa C, Teixeira B, Macedo R, Bittencourt A, Krueger R, Gross J, *et al.* Resistance exercise at variable volume does not reduce postprandial lipemia in postmenopausal women. *Age.* 2014;36(2): 869-79.
- Wooten J, Phillips M, Mitchell J, Patrizi R, Pleasant R, Hein R, *et al.* Resistance exercise and lipoproteins in postmenopausal women. *Int J Sports Med.* 2011;32(1):7-13.
- Zotou E, Magkos F, Koutsari C, Fragopoulou E, Nomikos T, Sidossis L, *et al.* Acute resistance exercise attenuates fasting and postprandial triglyceridemia in women by reducing triglyceride concentrations in triglyceride-rich lipoproteins. *Eur J Appl Physiol.* 2010;110(4):869-74.
- Chodzko-Zajko W. Exercise and physical activity for older adults. *Kinesiol Rev.* 2014;3(1):101-6.
- Libardi C, De Souza G, Cavaglieri C, Madruga V, Chacon-Mikahil M. Effect of resistance, endurance, and concurrent training on TNF- α , IL-6, and CRP. *Med Sci Sports Exerc.* 2012;44(1):50-6.

Sweating and core temperature in athletes training in continuous and intermittent sports in tropical climate

Anita M. Rivera-Brown^{1,2}, Farah A. Ramírez-Marrero³, Jessica Frontanés⁴, Raúl A. Rosario⁵, Osvaldo J. Hernández⁶, Walter R. Frontera^{1,2}

¹Department of Physical Medicine, Rehabilitation and Sports Medicine. University of Puerto Rico, San Juan, Puerto Rico. ²Department of Physiology and Biophysics, University of Puerto Rico, San Juan, Puerto Rico. ³Department of Physical Education and Recreation, Rio Piedras Campus, University of Puerto Rico, San Juan, Puerto Rico. ⁴St. Luke's University Health Network, Bethlehem, Pennsylvania. ⁵Mayo Clinic Sports Medicine, Jacksonville, FL, United States. ⁶Department of Physical Education and Recreation, San German Campus. Interamerican University of Puerto Rico, San German, Puerto Rico.

Received: 08.02.2018

Accepted: 26.06.2018

Summary

Higher sweat rate values have been reported during intermittent compared to continuous type exercise in hot conditions in the laboratory. Studies in the training field are scarce.

Objective: Document sweat rate, rehydration and core temperature (T_c) during training in long duration-continuous (CON) and intermittent (INT) type sports, and determine the presence of a mutation in the CFTR gene in athletes with high sodium concentration ($[Na^+]$) in sweat.

Methods: Athletes (CON =50 and INT=123) were tested during training in tropical climate. Sweat rate, sweat $[Na^+]$, T_c , dehydration level, and presence of $\Delta F508$ gene mutation in the CFTR gene in athletes with high sweat $[Na^+]$ were evaluated.

Results: Sweat rate was higher in CON (1.5 ± 0.4 L/h) compared to INT (1.2 ± 0.5 L/h) and CON athletes finished training with higher dehydration (2.1 ± 0.8 vs $1.2 \pm 0.7\%$) $P < 0.05$. Sweat $[Na^+]$ was higher in INT (62.0 ± 21.1 mmol/L) compared to CON (53.9 ± 18.1 mmol/L), $P < 0.05$. A tennis player with high $[Na^+]$ was heterozygous for the $\Delta F508$ mutation. Average and highest T_c was similar for CON (38.4 ± 0.3 and 38.8 ± 0.4 °C) and INT (38.3 ± 0.3 and 38.7 ± 0.4 °C), $P > 0.05$.

Conclusion: During training in a tropical climate, sweat loss and dehydration level are lower, and fluid intake is higher in intermittent compared to continuous type sports. Core temperature may rise to a similar level in intermittent type sports due to the repeated high intensity bouts and/or the effects of clothing worn while training in hot venues. Healthy athletes with high $[Na^+]$ in sweat who are heterozygous carriers of CFTR mutations may be at increased risk for hyponatremic dehydration and whole-body muscle cramps.

Key words:

Sweating. Tropical climate. Dehydration. Athletes. Body temperature. Sodium.

Sudoración y temperatura interna en atletas durante entrenamiento para deportes continuos e intermitentes en clima tropical

Resumen

Se reportan tasas de sudoración más altas durante ejercicio intermitente comparado con continuo en condiciones de calor en el laboratorio. Estudios en el campo de entrenamiento son escasos.

Objetivo: Documentar la tasa de sudoración, rehidratación y temperatura central (T_c) durante entrenamiento para deportes de tipo continuo (CON) e intermitente (INT), y determinar presencia de mutación genética en el gen CFTR en atletas con alta concentración de sodio ($[Na^+]$) en sudor.

Metodología: Se evaluó la tasa de sudoración, la $[Na^+]$ en sudor, la T_c y el nivel de deshidratación en atletas (CON =50; INT =123) durante entrenamiento en clima tropical, y la presencia de la mutación genética $\Delta F508$ en el gen CFTR en aquellos con alta $[Na^+]$ en sudor.

Resultados: La tasa de sudoración fue mayor en CON ($1,5 \pm 0,4$ L/h) comparado con INT ($1,2 \pm 0,5$ L/h) y los atletas en CON terminaron el entrenamiento con mayor deshidratación ($2,1 \pm 0,8$ vs $1,2 \pm 0,7\%$) $P < 0,05$. La $[Na^+]$ en sudor fue más alta en INT ($62,0 \pm 21,1$ mmol/L) comparado con CON ($53,9 \pm 18,1$ mmol/L), $P < 0,05$. Un tenista con alta $[Na^+]$ era heterocigoto para la mutación $\Delta F508$. La T_c promedio y más alta fueron similares para CON ($38,4 \pm 0,3$ y $38,8 \pm 0,4$ °C) e INT ($38,3 \pm 0,3$ y $38,7 \pm 0,4$ °C), $P > 0,05$.

Conclusión: Durante el entrenamiento en clima tropical, la pérdida de sudor y el nivel de deshidratación son más bajos, y la ingesta de líquido es más alta en deportes intermitentes que en deportes continuos. La temperatura interna puede aumentar a nivel similar en deportes intermitentes debido a periodos repetidos de alta intensidad y/o la vestimenta usada durante el entrenamiento. Atletas saludables con alta $[Na^+]$ en sudor que son heterocigóticos para mutaciones de CFTR pueden estar en mayor riesgo de deshidratación hiponatémica y calambres musculares.

Palabras clave:

Sudoración. Clima tropical. Deshidratación. Atletas. Temperatura corporal. Sodio.

Support: This study was supported in part by the "Comité Organizador de los Juegos Centroamericanos Mayagüez 2010" and the Puerto Rico Science, Technology & Research Trust.

Correspondence: Anita M. Rivera-Brown
E-mail: aniriver.brown@gmail.com

Introduction

Exercise in tropical environments may cause considerable elevations in core temperature that contribute to high sweat production and sodium loss, impairment in athletic performance¹ and heat related illnesses². The risk is higher if athletes begin training in a state of body fluid deficit and do not drink enough during exercise to replenish losses². Sweat losses of 1 to 2 L/h are common during training and competition in hot and humid tropical environments and may reach or exceed 2.5 L/h in heavy sweaters. Heat acclimatized elite athletes with high sweat production that train for long duration-continuous type (CON) sports, are most prone to dehydration. Athletes in intermittent type (INT) sports, such as team, ball and combat sports, are also at risk since multiple high intensity time periods elevate metabolic heat production and core temperature and induce heavy sweating³. This may be particularly true in athletes who wear heat-retaining clothing and/or protective equipment that decreases the effectiveness of heat loss through the evaporation of sweat.

Athletes with high sweat rates are more susceptible to large electrolyte losses in sweat, which have been implicated in the development of skeletal muscle cramps⁴. Athletes born and raised in the tropics may be protected from large sodium losses because regular exposure to high environmental temperatures induces physiological adaptations in the sweat glands such as an increased sodium reabsorption which result in a decreased sweat sodium concentration ($[Na^+]$) for a given sweat rate^{5,6}. However, a proportion of athletes exhibit "salty sweat". Those whose sweat $[Na^+]$ is typically higher than 55 mmol/L^{4,7} may have a reduced Cl^- and Na^+ reabsorption across the sweat duct membrane.

Studies have identified persons without cystic fibrosis (CF) have sweat Na^+ levels like those of CF patients⁸. It is possible that some healthy "salty sweaters," are heterozygous for a CF mutation and have a malfunction of plasma membrane CF transmembrane conductance regulator (CFTR) in the sweat glands that results in sweat with high sodium chloride concentration like CF patients⁹. Approximately 1 in 46 Hispanics carry one mutation of the CFTR gene that causes hypo-absorption of Na^+ in sweat gland ducts and high sweat $[Na^+]$ ¹⁰. In the United States, the most common mutation in the CFTR gene associated to the development of CF is the deletion of phenylalanine 508 ($\Delta F508$)¹¹ and its prevalence in athletes with salty sweat is currently unknown.

Studies in the laboratory have shown that prolonged variable intensity exercise with periods of high intensity exercise (≥ 15 minutes) may elicit similar or higher heat storage, core temperature (T_{c}), and sweat production, as prolonged constant intensity exercise with no rest periods^{3,12-14}. Studies in the playing field show a predisposition to heat strain due to hyperthermia and dehydration in American Football players¹⁵ but very little is known about other intermittent type or endurance sports in tropical environments. Particularly lacking are data in female athletes.

The aims of the present observational field study were to: 1) document the sweating response and rehydration during training sessions in tropical climate in CON and INT sports in both genders; 2) investigate the presence of the $\Delta F508$ mutation in the CFTR gene in athletes with high sweat $[Na^+]$; and 3) compare the T_{c} response in a subset of male and female athletes in CON and INT sports. We hypothesized that athletes

in INT sports would show a lower sweat rate, a lower average T_{c} and a higher sweat $[Na^+]$.

Material and method

Subjects

One hundred and seventy three competitive athletes participating in long duration-continuous (endurance=50; 30 males [M] and 20 females [F]) or intermittent (ball/team= 94, 45 M and 49 F; combat =29, 15 M and 14 F) sports (Table 1) were tested during a typical training session for their sport. The subjects were recruited from teams in a Central American and Caribbean competition in the summer in Puerto Rico and from National teams, top ranked university teams, and recreational athletes ranked in the top of their age categories in Puerto Rico. All were natives of tropical countries with a predominantly warm and humid climate all year long.

All subjects were informed of the purpose and procedures of the study and written informed consent was obtained. There were 52 athletes between the ages of 17 and 21 who gave their written assent, and written consent was also obtained from their parents. The study was approved by the Institutional Review Board of the University of Puerto Rico, Medical Sciences Campus.

Study design and procedures

Athletes were studied under environmental conditions typical of training sessions for their sport, and wearing uniforms and equipment used in competition. This field study design allowed us to collect information in conditions that are difficult to simulate in the laboratory. Athletes completed a questionnaire about heat exposure during exercise and past history of muscle cramps. General health status was also documented to assure they did not have a medical condition that would affect thermoregulatory or body fluid balance variables.

All subjects were tested during the summer months in Puerto Rico. The average WBGT (10 am to 4 pm) was 29.9 ± 1.6 °C¹⁶. The runners, cyclists, triathletes, soccer, tennis, and beach volleyball players trained outdoors exposed to direct sunlight, while the judo, basketball, team

Table 1. Athletes in long duration-continuous (CON) and intermittent (INT) sports.

CON (N=50) Endurance Sports	INT (N=123) Team/Ball Sports		F	M	
	F	M			
Running	8	13	Soccer	10	18
Cycling	7	9	Tennis	3	10
Triathlon	5	8	Basketball	7	0
			Team Handball	13	0
			Beach Volleyball	16	17
			Combat Sports		
			Judo	7	3
			Fencing	7	12
Total	20	30		63	60

handball, and fencing athletes trained indoors in non-air conditioned gyms. The triathletes were tested while running. All athletes wore official competition uniforms required for their sport. The judokas and fencers wore t-shirts and shorts underneath the uniforms.

- *Pre-training session.* Subjects urinated into a pre-weighed container. Nude body weight was determined using an electronic scale accurate to 50 g (precision health scale uc321). After cleaning the area with distilled water, absorbent patches (Tegaderm +pad, 3m Health Care, Borken, Germany) were applied to the subject's right posterior forearm and right anterior thigh for regional sweat collection in all but the beach volleyball players, in which it was applied to the right anterior thigh only. Athletes were then fitted with an accelerometer (Actigraph GTX3; Actigraph, Pensacola, FL, USA) that was programmed to record activity counts in one-minute epochs. The cyclists wore the accelerometer on the right thigh.
- *Training session.* The duration of training was similar to a typical session for the sport. Training sessions were organized and directed by the athletes' coaches. Environmental conditions were measured before, every 30 min, and at the end of training, using a heat stress monitor (Questemp 32, Quest Technologies, Wisconsin), which was placed in the training area. Athletes drank voluntarily from labeled bottles with water and/or sports drinks. Any subject who needed to urinate during the training session did so into a pre-weighed container and the volume was measured. The absorbent patches were removed when saturated and placed in sealed plastic tubes.
- *Post-training session.* Athletes removed the accelerometer and clothes, towel dried and pre-training measures were repeated (urine sample, nude body weight).
- *Measures and calculations*
- *Hydration status.* Urine specific gravity (USG) was used as an index of pre-exercise hydration status¹⁷ and determined using a hand-held refractometer (URC-NE, Atago Clinical, Japan).
- *Environmental conditions.* Dry bulb (DB), wet bulb (WB), globe temperature (GT), and relative humidity (RH) were measured on site. The wet bulb globe temperature (WBGT) heat stress index was calculated using the following equations: CON sports: $(WB \times 0.7) + (GT \times 0.2) + (DB \times 0.1)$; INT sports: $(WB \times 0.7) + (GT \times 0.3)$ ¹⁸.
- *Regional sweat [Na⁺].* Sweat was extracted from each patch using a 5-ml syringe and two 100 μ l samples were analyzed using an ion selective electrolyte analyzer (Easylyte Plus, Medica, Bedford, MA) to determine the [Na⁺].
- *Body fluid balance.* The bottles with the fluid that was used for hydration during the session were weighed before and after the session with a scale accurate to 1 g (CS2000 Compact Scale). Sweat loss was calculated as: change in body mass, corrected for fluid intake and urine loss. Sweat rate was calculated as sweat loss divided by exercise duration. The level of dehydration was determined using the following formula: $(\text{body mass pre-training} - \text{body mass post-training}) / \text{body mass pre-training}$.
- *Core temperature.* Core temperature was measured in 62 athletes that lived and trained in Puerto Rico (females, CON= 10 and INT=34; males, CON=13 and INT=5) using ingestible sensors (Cortemp™, HQ Inc.) before, every 10 minutes, and at the end of training. We measured T_c only in athletes who could satisfy requirements for

valid measurements (sensor calibration, timing of ingestion and electromagnetic interference). Eight hours prior to the beginning of the exercise session, each subject swallowed a disposable pill that contained a temperature sensor for T_c determination. Pre-exercise T_c was the value taken after sitting quietly for five minutes in the exercise area before the start of training. The T_c of the exercise session was the average of measures taken every 10 minutes. The change in T_c was calculated as the highest minus the pre-exercise T_c . A final measure was taken immediately after the coach indicated the session had finished.

Genetic analysis

Athletes with a sweat $[Na^+] \geq 70$ mmol/l were identified for subsequent genetic analysis¹⁹. Athletes were asked to provide two ml of saliva which were collected into a DNA genotek's oragene® DNA self-collection vial. The container had approximately 2 ml of cell lysis solution or DNA-preserving fluid, that was mixed with the saliva. Genomic DNA was assayed for the Δ F508 mutation in the CFTR gene, using restriction fragment length polymorphism (Ambry Genetics, Aliso Viejo, CA). As per standard procedures, DNA was extracted from saliva and purified. The purified DNA was digested using restriction endonucleases. The restriction fragments produced during DNA fragmentation were analyzed using gel electrophoresis.

Statistical Analysis

Means and standard deviations were calculated for each variable. Student's *t*-tests were used to compare CON and INT in pre-exercise urine specific gravity, sweat rate, percent dehydration, percent rehydration, minutes of exposure to exercise in the heat in the past month, activity counts/min, sweat $[Na^+]$, pre, average, change and highest T_c , and the WBGT index. An alpha level of $P < 0.05$ was considered significant.

Results

Subjects, environmental conditions and training sessions

Descriptive characteristics of the subjects and training session characteristics are presented in Table 2. The WBGT index for CON and INT was $> 28^\circ\text{C}$, which is typical of tropical countries during the summer and indicates that all subjects were exposed to a high level of heat stress without exhibiting symptoms of heat illness. Activity counts per minute²⁰ revealed that in INT sports the training sessions were of predominantly light- to- moderate intensity whereas in CON sports they were shorter and of vigorous intensity. The average time spent training exposed to solar radiation in the month preceding the study was 17% higher in CON.

Hydration status and fluid balance

Athletes in INT arrived to the training session in a state of hypo-hydration as evidenced by a mean pre-exercise USG of 1.021 ± 0.006 . Athletes in CON had a higher sweat rate ($P < 0.05$) and finished training with higher ($P < 0.05$) level of dehydration (2.1 ± 0.8 vs $1.2 \pm 0.7\%$) due

Table 2. Descriptive characteristics of the subjects, environmental conditions and training sessions in long duration-continuous (CON) and intermittent (INT) type sports.

	CON (N=50)			INT (N=123)		
	Females N=20	Males N=30	Total N=50	Females N=63	Males N=60	Total N=123
Age (y)	26.0 ± 6.5	26.2 ± 5.4	26.1 ± 5.8	23.4 ± 5.2	23.9 ± 4.4*	23.6 ± 4.8
Height (cm)	159.7 ± 5.5	173.9 ± 7.1	168.2 ± 9.6	167.2 ± 9.4*	179.0 ± 8.9*	173.1 ± 10.9*
Weight (kg)	53.8 ± 5.9	68.5 ± 10.7	62.6 ± 11.6	65.8 ± 10.2*	76.5 ± 11.2*	71.0 ± 11.9*
Training exposed to sun (min/day in past month)	122.0 ± 63.2	164.3 ± 83.1	147.4 ± 78.0	96.7 ± 84.4	147.5 ± 90.0	121.9 ± 90.5
Duration of Training Session (min)	69.6 ± 20.6	78.7 ± 24.0	75.0 ± 22.9	91.6 ± 19.1*	86.8 ± 20.1	89.2 ± 19.7*
Activity counts (counts/min)	7,695.0 ± 1,666.7	7,189.7 ± 2,367.7	7,414.3 ± 2,078.9	2,421.9 ± 539.6*	2,909.0 ± 882.4*	2,654.7 ± 760.9*
Minutes at light intensity (< 1,952 counts/min)	1.7 ± 1.9	8.6 ± 18.9	5.6 ± 14.4	46.4 ± 18.2*	32.6 ± 17.2*	39.8 ± 19.0*
Minutes at moderate intensity (1,952 – 5,724 counts/min)	10.4 ± 21.7	16.6 ± 30.1	13.8 ± 26.6	37.8 ± 11.6*	45.5 ± 17.1*	41.5 ± 14.9*
Minutes at vigorous intensity (> 5,724 counts/min)	57.1 ± 28.2	54.9 ± 28.3	55.9 ± 27.9	6.8 ± 5.0*	8.7 ± 8.4*	6.9 ± 0.6*
WBGT heat stress index (°C)	30.9 ± 1.1	30.8 ± 1.7	30.8 ± 1.1	28.9 ± 1.4*	30.2 ± 1.7	29.5 ± 1.7*

*Significant difference between CON and INT, $P < 0.05$.

Table 3. Hydration status and fluid balance in long duration-continuous (CON) and intermittent (INT) type sports.

	CON			INT		
	Females N=20	Males N=30	Total N=50	Females N=63	Males N=60	Total N=123
USG Pre (mg/dL)	1.017 ± 0.001	1.018 ± 0.001	1.017 ± 0.000	1.020 ± 0.001	1.022 ± 0.000*	1.021 ± 0.006*
Sweat rate (L/h)	1.3 ± 0.3	1.6 ± 0.4	1.5 ± 0.4	0.9 ± 0.3*	1.5 ± 0.4	1.2 ± 0.5*
Dehydration (%IBW)	1.9 ± 0.7	2.2 ± 0.8	2.1 ± 0.8	0.9 ± 0.6*	1.4 ± 0.8*	1.2 ± 0.7*
Fluid replaced (% of fluid loss)	30.9 ± 20.9	29.3 ± 20.0	29.9 ± 20.1	57.8 ± 27.5*	52.8 ± 19.3*	55.4 ± 23.9*
Sweat [Na ⁺] (mmol/L)	57.2 ± 20.6	57.2 ± 18.2	53.9 ± 18.1	63.2 ± 22.6	67.7 ± 18.7*	62.0 ± 21.1*

*Significant difference between CON and INT, $P < 0.05$.


to lower fluid replacement. Sweat [Na⁺] was higher ($P < 0.05$) in INT compared to CON and there was no difference between genders and history of muscle cramps.

Sixty athletes (28% of the group) had a sweat [Na⁺] >70 mmol/L and sweat [Cl⁻] > 60 mmol/l. Forty-nine of those (82%) provided saliva samples for genetic analysis. One male tennis player was identified as having heterozygous presence of the $\Delta F508$ mutation in the CFTR gene. He had a history of whole body muscle cramps and his sweat [Na⁺] was 74.9 mmol/L (average of thigh and forearm regional sweat samples).

Core temperature

We measured core temperature in 23 athletes in CON (females=10; males=5) and in 39 athletes in INT (females=34; males=5). For this sub-group, the CON sports were running, cycling, and triathlon and the INT sports were soccer, basketball, handball, and judo. For the whole group, the average and highest T_c for the session was similar for CON (38.4 ± 0.3 and 38.8 ± 0.4 °C) and INT (38.3 ± 0.3 and 38.7 ± 0.4 °C), $P < 0.05$.

In Figure 1 we present the T_c before, during and at the end of the INT training sessions. The judo athletes, who were exposed to a high

Figure 1. Core temperature before, during, and at the end of the intermittent intensity training sessions.

+Soccer higher than tennis and basketball; *Judo higher than soccer and handball; #judo higher than all; $P < 0.05$.

level of heat stress while wearing two layers of clothing, showed the highest values for T_c during the final part and at the end of the INT exercise sessions.

Discussion

The present study examined the body hydration status and thermoregulatory responses to exercise under tropical conditions in athletes in continuous and intermittent type sports. The main findings were: 1) sweat rate and the change in core temperature were higher in CON compared to INT; 2) low fluid intake resulted in a higher level of dehydration in CON; 3) a proportion of athletes exhibit "salty sweat" and one was a healthy carrier of a CFTR gene mutation; and 4) athletes in INT exhibited maximal T_c similar to athletes in CON who trained at high intensity for longer periods.

Hydration status and fluid balance

Athletes in our study are indigenous to the tropics and have physiological adaptations induced by repeated exposure to a hot and humid environment that lead to increased capacity and sensitivity of the sweat glands²¹. The sweat rate values in males (1.6 ± 0.4 L/h) and females (1.3 ± 0.3 L/h) in CON and in males in INT (1.5 ± 0.4 L/h) in the present study are higher than normative data reported for adult male (1.42 ± 0.72 L/h) and female (1.10 ± 0.57 L/h) athletes²².

Whole body sweat rate comparisons between continuous and intermittent type exercise in the heat in controlled, artificial, thermoneutral^{13,23}, and hot-dry³ conditions have revealed mixed results. Drust *et al.*¹³ found a similar sweat rate during a laboratory-based soccer protocol in male athletes that ran on a treadmill for 45 minutes either intermittently (7.5 min exercise/1.2 min rest) or continuously in a thermoneutral environment. In contrast, Mora-Rodriguez *et al.*³ found higher sweat rate during 90 min of variable intensity intermittent exercise (1.5 min high intensity/4.5 min low intensity) compared to same amount of work performed at a constant load in endurance-trained, heat-acclimated males exercising in hot-dry environment. In the present field study, in which we did not control for exercise intensity or environmental conditions, we found higher sweat rates in athletes in CON which may be expected for two reasons: 1) a greater proportion of the CON sessions were at high exercise intensity; and 2) athletes in CON were exposed to a higher heat stress.

Our data are consistent with the observation that athletes replace less than 50% of their sweat losses during exercise and thus, they are hypohydrated at the end of training²⁴. In CON sports opportunities to drink may be frequent but athletes may be reluctant to drink, to avoid slowing down and the abdominal discomfort associated with large fluid consumption. On the other hand, in intermittent type sports, regular opportunities for hydration are available during substitutions, pauses in play, time-outs, intermissions, and change of sides of court, which leads to frequent and better fluid replacement.

Sweat sodium and genetic mutation in CFTR gene

When the sweat rate is high, sodium reabsorption capacity in the sweat gland can be exceeded resulting in a higher $[Na^+]$ appearing in

sweat. However, the linear increase seen in sweat $[Na^+]$ as sweat rate increases can be modulated by heat acclimation⁵. For example, Buono *et al.*⁵ showed that heat acclimation may increase the absorptive capacity of the sweat gland duct, reducing the sweat $[Na^+]$ for a given sweat rate up to $1 \text{ mg} \cdot \text{cm}^{-2} \cdot \text{min}^{-1}$. In the present study athletes in CON showed a lower sweat $[Na^+]$ despite their higher sweat rate in comparison to athletes INT, which may be attributed to their higher level of heat exposure and acclimatization. An advantage of a higher sweat rate coupled with a lower sweat $[Na^+]$ for athletes in CON is a lower risk for developing hyponatremia during exercise in the heat²⁵.

A factor proposed for high Na^+ excretion in healthy athletes is the presence of genetic mutations in the CFTR that affects the Cl^- channel in the sweat glands^{7,8}. The majority of CF patients have the $\Delta F508$ mutation and exhibit $[Na^+]$ in their sweat that may be up to five times higher than healthy individuals²⁶. It is estimated that 2% (1 in 46) of Hispanic Americans are heterozygous carriers of a mutated CFTR gene¹⁰ and may be genetically predisposed to high sweat Na^+ loss. The prevalence of CF mutations in athletes with salty sweat is currently unknown.

A reduced ductal luminal membrane expression of the Cl^- channel was found in six healthy recreationally active subjects with a mean sweat $[Na^+]$ of $94.9 \pm 15.2 \text{ mmol/l}$. However, in that study, none of the subjects was a carrier of any of the 39 most common CFTR mutations in the United States. The authors concluded that a full CFTR gene sequencing of the subjects would be required before excluding a possible genetic link between excessive Na^+ loss in sweat and CFTR mutations.

A novel finding of our study was that one of 45 Hispanic athletes (2%), classified as "salty sweaters" was heterozygous for the $\Delta F508$ mutation in the CFTR gene. That athlete was a tennis player with a history of whole body muscle cramps that lead to collapse on court requiring medical attention.

Core temperature response

During prolonged high intensity exercise, the heat stored is higher and T_c rises faster when the same amount of exercise is performed in a variable intensity mode that includes short (1.5 to 7.5 min) high intensity bouts followed by rest periods compared to constant intensity mode^{3,13,23,27}. However, if longer bouts (15-30 min) of high intensity exercise are used, a similar T_c rise is observed^{28,29}. We found that although athletes in INT spent the majority of the exercise session at lower exercise intensity with multiple resting periods, they exhibited a maximal T_c (38.7 ± 0.4 °C) similar to athletes in CON (38.8 ± 0.4 °C) who trained at high exercise intensity for most of the session. This response suggests that the heat stored during the few periods of high exertion was not dissipated significantly and T_c remained elevated instead of decreasing as expected during low intensity exercise periods.

The highest core temperature response in INT was observed in judokas who trained in a non-air-conditioned gym (WBGT= 30.0 °C). Similar to our previous report³⁰, judokas in the present study arrived to training with a fluid deficit, as evidenced by USG. The combination of a pre-exercise fluid deficit that worsened during training because of inadequate fluid intake, hot and humid environmental conditions, and insulation properties of the uniform that decreased heat dissipation, predisposed the athletes to hyperthermia during training. In fact, 8 of the 9 judokas felt overheated and 7 of them had maximal $T_c \geq 39.0$ °C.

Two athletes reached a T_{c} of 39.6 °C and felt dizzy, had a headache, and stopped training with symptoms of exhaustion. High T_{c} and increased physiologic strain have been reported in American football athletes wearing a full uniform while training in similar climatic conditions¹⁵. We believe this to be the first study documenting the rate of T_{c} rise in judo athletes during training in the heat. Our data indicate that training in full uniform in a non-air conditioned gym in the tropics predisposes judo athletes to heat illness and should be discouraged.

Conclusions

This observational field study demonstrates that during training in a tropical climate, sweat loss and dehydration level are lower, and fluid intake is higher in intermittent compared to continuous type sports. Nonetheless, core temperature may rise to a similar level in intermittent type sports due to the repeated high intensity bouts and/or the effects of clothing worn while training in hot venues. For athletes in continuous sports, high sweat rates may not provide a thermoregulatory advantage in a tropical climate because humidity hinders the evaporation of sweat. Coaches should pay special attention to factors that may increase fluid intake, such as availability of cool, palatable fluids, encouragement to drink, and frequent, short pauses. Healthy athletes with high $[Na^+]$ in sweat who are heterozygous carriers CFTR mutations may be at increased risk for hyponatremic dehydration and whole-body muscle cramps during exercise.

Acknowledgements

The authors thank the athletes and their coaches for their enthusiastic participation in the study, research assistants José Quiñones, Javier Ríos, Eddie Rodríguez, and David Colón, and medical personnel at the Center for Sports Health and Exercise Sciences for their help during testing.

Conflict of interest

The authors do not declare a conflict of interest.

Bibliography

- Gonzalez-Alonso J, Crandall CG, Johnson JM. The cardiovascular challenge of exercising in the heat. *J Physiol*. 2008;586(1):45-53.
- Sawka MN, Burke LM, Eichner ER, Maughan RJ, Montain SJ, Stachenfeld NS. American College of Sports Medicine position stand. Exercise and fluid replacement. *Med Sci Sports Exerc*. 2007;39(2):377-90.
- Mora-Rodriguez R, Del Coso J, Estevez E. Thermoregulatory responses to constant versus variable-intensity exercise in the heat. *Med Sci Sports Exerc*. 2008;40(11):1945-52.
- Stofan JR, Zachwieja JJ, Horswill CA, Murray R, Anderson SA, Eichner ER. Sweat and sodium losses in NCAA football players: a precursor to heat cramps? *Int J Sport Nutr Exerc Metab*. 2005;15(6):641-52.
- Buono MJ, Ball KD, Kolkhorst FW. Sodium ion concentration vs. sweat rate relationship in humans. *J Appl Physiol*. 2007;103(3):990-4.
- Kirby CR, Convertino VA. Plasma aldosterone and sweat sodium concentrations after exercise and heat acclimation. *J Appl Physiol*. 1986;61:967-70.
- Brown MB, Haack KK, Pollack BP, Millard-Stafford M, McCarty NA. Low abundance of sweat duct Cl⁻ channel CFTR in both healthy and cystic fibrosis athletes with exceptionally salty sweat during exercise. *Am J Physiol Regul Integr Comp Physiol*. 2011;300(3):R605-615.
- Sato K, Sato F. Na⁺, K⁺, H⁺, Cl⁻, and Ca²⁺ concentrations in cystic fibrosis eccrine sweat in vivo and in vitro. *J Lab Clin Med*. 1990;115(4):504-11.
- Eichner ER. Genetic and other determinants of sweat sodium. *Curr Sports Med Rep*. 2008;7:S36-S40.
- Bobadilla JL, Macek M, Jr, Fine JP, Farrell PM. Cystic fibrosis: a worldwide analysis of CFTR mutations--correlation with incidence data and application to screening. *Hum Mutat*. 2002;19(6):575-606.
- Riordan JR, Rommens JM, Kerem B, et al. Identification of the cystic fibrosis gene: cloning and characterization of complementary DNA. *Science*. 1989;245:1066-72.
- Sawka MN, Latzka WA, Montain SJ, et al. Physiologic tolerance to uncompensable heat: intermittent exercise, field vs laboratory. *Med Sci Sports Exerc*. 2001;33(3):422-30.
- Drust B, Reilly T, Cable NT. Physiological responses to laboratory-based soccer-specific intermittent and continuous exercise. *J Sports Sci*. 2000;18(11):885-92.
- Gagnon D, Kenny GP. Exercise-rest cycles do not alter local and whole body heat loss responses. *Am J Physiol Regul Integr Comp Physiol*. 2011;300(4):R958-68.
- Armstrong LE, Johnson EC, Casa DJ, et al. The American football uniform: uncompensable heat stress and hyperthermic exhaustion. *J Athl Train*. 2010;45(2):117-27.
- Armstrong LE, Casa DJ, Millard-Stafford M, Moran DS, Pyne SW, Roberts WO. American College of Sports Medicine position stand. Exertional heat illness during training and competition. *Med Sci Sports Exerc*. 2007;39(3):556-72.
- Casa DJ, Armstrong LE, Hillman SK, et al. National Athletic Trainers' Association position statement: Fluid replacement for athletes. *J Athl Train*. 2000;35:212-24.
- Yaglou C, Minard D. Control of heat casualties at military training centers. *Arch Ind Health*. 1957;16:302-5.
- Brown MB, McCarty NA, Millard-Stafford M. High-sweat Na⁺ in cystic fibrosis and healthy individuals does not diminish thirst during exercise in the heat. *Am J Physiol Regul Integr Comp Physiol*. 2011;301(4):R1177-85.
- Freedson PS, Melanson E, Sirard J. Calibration of the Computer Science and Applications, Inc. accelerometer. *Med Sci Sports Exerc*. 1998;30(5):777-81.
- Inoue Y, Havenith G, Kenney WL, Loomis JL, Buskirk ER. Exercise- and methylcholine-induced sweating responses in older and younger men: effect of heat acclimation and aerobic fitness. *Int J Biometeorol*. 1999;42:210-6.
- Baker LB, Barnes KA, Anderson ML, Passe DH, Stofan JR. Normative data for regional sweat sodium concentration and whole-body sweating rate in athletes. *Journal of Sports Sci*. 2016/02/16 2016;34(4):358-68.
- Eklom B, Greenleaf CJ, Greenleaf JE, Hermansen L. Temperature regulation during continuous and intermittent exercise in man. *Acta Physiol Scand*. 1971;81(1):1-10.
- Broad EM, Burke LM, Cox GR, Hecley P, Riley M. Body weight changes and voluntary fluid intakes during training and competition sessions in team sports. *Int J Sport Nutr*. 1996;6:307-20.
- Montain SJ, Chevront SN, Sawka MN. Exercise associated hyponatremia: quantitative analysis to understand the aetiology. *Br J Sports Med*. 2006;40(2):98-105.
- Quinton PM. Cystic fibrosis: a disease in electrolyte transport. *FASEB J*. 1990;4(10):2709-17.
- Kraning KK, Gonzalez RR. Physiological consequences of intermittent exercise during compensable and uncompensable heat stress. *J Appl Physiol*. 1991;71(6):2138-45.
- Belding HS, Hertig BA, Kraning KK. Comparison of man's responses to pulsed and unpulsed environmental heat and exercise. *J Appl Physiol*. 1966;21(1):138-42.
- Lind AR. A physiological criterion for setting thermal environmental limits for everyday work. *J Appl Physiol*. 1963;18:51-6.
- Rivera-Brown AM, De Felix-Davila RA. Hydration status in adolescent judo athletes before and after training in the heat. *Int J Sports Physiol Perform*. 2012;7(1):39-46.

The effect of tapering and *Nigella sativa* on the histological structure of the lung after increasing interval exercise training

Shadmehr Mirdar¹, Forough Neyestani¹, Gholamreza Hamidian², Mohammad Ali Khalilzadeh³

¹Department of Exercise Physiology, Faculty of Physical Education and Sport Sciences, University of Mazandaran, Babolsar, Iran. ²Department of Basic Sciences, Faculty of Veterinary Medicine, University of Tabriz, Tabriz, Iran. ³Department of Phytochemistry, Qaemshahr Branch, Islamic Azad University, Qaemshahr, Iran.

Received: 12.12.2017

Accepted: 09.07.2018

Summary

Objectives: During maturation period in which the immune system of lung tissue is not fully developed, physical exercises may have a negative effect and cause inflammation. This study aimed to investigate the effects of tapering and *Nigella sativa* (NS) hydro-alcoholic extract on the reduction of lung tissue inflammation caused due to increasing interval exercise training (IET) during maturation period by histological and stereological methods.

Methods: Ninety-five three weeks old rats after adaption were randomly divided into two control and exercise groups and 19 subgroups. The exercise group carried out a period of six weeks of undulating IET followed by three weeks of load reduction performed by three models in two different times. Rats entered the taper phase were administrated by NS supplement in tapering and control groups. The lung tissue samples were processed by standard paraffin embedding, stained by H&E and examined by using point counting method through systematic random sampling in stereological study. The results were analyzed using by two-way ANOVA and LSD post hoc in $\alpha=0.05$.

Results: The result showed that IET caused severe inflammation in lung tissue and an increase in infiltration of inflammatory cells and lymphocytes into the connective tissues surrounding the respiratory air ways, vessels and interstitial lamellae. This severity of inflammation was considerably and similarly more in comparison to the basic and control groups ($p=0.001$). Stereological analysis in the taper exercise training groups with NS and without NS as well, revealed a significant decrease in the degree and intensity of lung tissue inflammation in the examined times in comparison to the IET group ($p=0.001$).

Conclusion: Generally it can be concluded that performing NS and a three weeks period of tapering has a noticeable effect in the reduction of inflammation in lung tissue followed by interval exercise training.

Key words:

Increasing interval exercise training. Tapering. *Nigella sativa*. Lung. Inflammation.

El efecto del tapering y *Nigella sativa* sobre la estructura histológica del pulmón después de aumentar el entrenamiento de ejercicio de intervalo

Resumen

Objetivos: Durante el período de maduración en el que el sistema inmunitario del tejido pulmonar no está completamente desarrollado, el ejercicio físico puede tener un efecto negativo y causar inflamación. Este estudio tuvo como objetivo investigar los efectos del tapering y del extracto hidroalcohólico de *Nigella sativa* (NS) sobre la reducción de la inflamación del tejido pulmonar causada por el aumento de entrenamiento interválico (IET) durante el período de maduración mediante métodos histológicos y estereológicos.

Métodos: Noventa y cinco ratas de tres semanas de edad, después de la adaptación, se dividieron aleatoriamente en dos grupos de control y ejercicio y 19 subgrupos. El grupo de ejercicio llevó a cabo un período de seis semanas de IET ondulado seguido de tres semanas de tapering realizadas por tres modelos en dos momentos diferentes. Las ratas entraron en la fase de tapering y se les administró un suplemento de NS en ambos grupos. Las muestras de tejido pulmonar se procesaron mediante inclusión convencional de parafina, se tiñeron con H & E y se examinaron mediante el método de conteo puntual mediante muestreo aleatorio sistemático en un estudio estereológico. Los resultados se analizaron usando ANOVA de dos factores y LSD *post hoc* en $\alpha = 0,05$

Resultados: Los resultados mostraron que el IET causó inflamación severa en el tejido pulmonar y un aumento en la infiltración de células inflamatorias y linfocitos en los tejidos conectivos que rodean las vías respiratorias, los vasos y las lamelas intersticiales. Esta gravedad de la inflamación fue considerablemente mayor y similar en comparación con los grupos básico y de control ($p = 0,001$). El análisis estereológico en los grupos de tapering con NS y sin NS también, reveló una disminución significativa en el grado e intensidad de la inflamación del tejido pulmonar en las mediciones examinadas en comparación con el grupo IET ($p = 0,001$).

Conclusión: en general, se puede concluir que la realización de NS y un período de tapering de tres semanas tiene un efecto notable en la reducción de la inflamación en el tejido pulmonar seguida de entrenamiento de ejercicios a intervalos.

Palabras clave:

Aumento del entrenamiento de ejercicio interválico. Tapering. *Nigella sativa*. Pulmón. Inflamación.

Correspondence: Gholamreza Hamidian
E-mail: hamidian@tabrizu.ac.ir

Introduction

Physical exercises with different intensity create different responses on the immune system function^{1,2}. Increase in the volume and intensity of the exercise training along with a decrease in the recovery time may cause an overtraining in athletes and affect their immune system³. On the other hand, physical exercise can be a strong stimulus in the development of lung inflammation⁴. Findings of several studies shows that exercise training load beyond the normal range increases the risk of upper respiratory tract infection (URTI)^{2,5} while moderate exercise not only decrease these risks but also reinforce the function of respiratory immune system^{1,6}. Researches showed that the URTI symptoms in the most of the elite athletes are similar to general population⁶, however, usual seasonal patterns of URTI are not seen in these athletes⁷. It is found out that short and long term physical exercise trainings with different grades and intensities affect the immunity and inflammatory factors in childhood and adulthood period⁸. There are still many unanswered questions on physical exercise mechanisms influencing inflammatory index before and during puberty⁹.

Since mild inflammation in early stages is a helpful protective response against primary cellular damage factors which eliminates external invaders and necrotic tissue and also because too much inflammation causing severe damage to lung tissue can be life-threatening¹⁰, it seems necessary to take some strategies in exercise training programs including taper to prevent overtraining and immune function decline^{11,12}. Taper can be performed in the forms of frequency reduction, repetition and intensity of exercise training^{3,13} in different time periods. One of the hardest challenges for sport science researchers and trainers is considered as to determine the most appropriate taper program¹⁴. Limited studies about intensity of exercise training during taper period showed athletes can make use of training intensity reduction programs in order to improve their performance in competition season¹⁵. Some reports reveled that time execution of swimmers¹⁶, runners¹⁷ and bike riders¹⁸ improved due to taper programs. Previous studies suggest the favorable time period for taper is between 4 to 28 days or even more^{8,13}. Though many studies have confirmed a two-week period taper, there have been some reports on the improvement of athletes' performance due to very short or very long period tapers¹⁹. Thomas *et al.*³ concluded that the optimal time duration for taper depends on the training before the taper and no particular time limit can be specified for that.

Today, there are several methods other than taper such as medicinal plants using for reducing the symptoms of inflammation and boosting the immune system. Among these plants, *Nigella sativa L.* (*NS*) is a medicinal plant known in Iranian traditional medicine. This native plant has been used for the treatment of headaches, nasal congestion, asthma, and allergy as well as for boosting the immune system^{20,21}. The biological properties of the seed of this plant include Anti-oxidative, anti-inflammatory, anti-tumor, anti-viral and anti-microbial effects as well as strengthening the immune system²². Previous studies indicated that *NS* has a protective effect in lung injury and pulmonary fibrosis^{23,24}. Gholamzadeh *et al.*²⁵ revealed that *NS* has an anti-inflammatory effect and reduces pro/anti-inflammatory cytokine ratio in overtrained animals. They showed that this effect of *NS* is more pronounced in overtraining animals than control or moderate exercise animals. In fact, *NS* causes

an immunoregulatory effect which somehow homogenizes immune state during different physiological status²⁵.

Since in the previous studies, the compatibilities of the high intensity trainings, the effects of taper and *NS* use during maturation period as to the inflammatory response of the lung did not come to a clear conclusion, this study was designed to investigate the effect of the performing different patterns of taper following a high-intensity interval training (HIIT) as well as influence of interactive effect of taper and *NS* on the microscopic properties of inflammation in lung tissue of male Wistar rats during maturation period.

Material and method

Animals

In this study, 95 male Wistar rats with an approximate age of 3 weeks and an average weight of 68±9 g were obtained from Pasteur Institute of Iran. In order to adaptation, animals (5 rats per cage) were maintained in transparent polycarbonate cages under controlled environment with a temperature of 23±2 °C, humidity of 45-55 % and 12:12 hours light/dark cycle for two week. Throughout all stages of the study, ethics of working with laboratory animals such as free access to standard pellet diet and water *ad libitum*, euthanizing without pain, prevention of pain associated with surgery and sampling were taken into consideration according to the international recommendations about clinical and laboratory animals' researches, ratified in Helsinki and updated in 2008 by the American Physiology. After one weeks of familiarity with laboratory and manipulation, the rats were randomly divided into two control and exercise groups matched for their weight.

Exercise training program

The rats were divided into two control and exercise groups at first. Then, for getting acquainted with the treadmill, they performed the exercise training with the main pattern training including increasing interval training (IIT) but with a lower intensity for two weeks. After one week period of getting acquainted with the environment and the treadmill, they performed increasing interval undulating exercise training for six weeks and that was followed by 3 weeks of load reduction (taper) carried out by 3 models in two different times. Treatment and un-treatment with *NS* supplement was also observed during the tapering (Table 1). Finally, according to research design and sampling procedure after 6 weeks increasing interval exercise training, the first and third week of tapering, the animals were divided into 19 groups and each group included 5 rats.

The familiarity and compatibility phase included 4 sessions of interval exercise training per week with the speed of 10 to 25 meters per minute and the slope of zero percent lasting for 15 to 30 minutes. The increasing interval exercise training program was carried out in the form of 10 repetitions of 1 minute length and active rests of 2 minute length minutes in such a way that the total daily workout time for each rat was 30 minutes-long. The animals started the increasing interval exercise training at the speed of 25 m/min and ended it at the speed of 70 m/min²⁶. Apart from the main activity, 5 minutes was estimated for warm-

Table 1. General specification of research protocol.

Weeks of training	Familiarization	1	2	3	4	5	6	7	8	9
Age (week)	4	5	6	7	8	9	10	11	12	13
Control	Control orientation	Control						taper control with NS		
								taper control without NS		
Interval training	Orientation	Training period						taper with NS		
								taper without NS		

Table 2. Increasing interval 6-weeks training program.

Week	Familiarization	First	Second	Third	Fourth	Fifth	Sixth
Training speed (m/min)	10-25	25-35	30-45	45-55	50-65	60-70	65-70
Rest speed (m/min)	10	10-20	15-25	25-30	25-35	30-35	30-35
Training duration (min)	1	1	1	1	1	1	1
Rest time between replications (min)	2	2	2	2	2	2	2
Set number	10	10	10	10	10	10	7
Session number per week	4	5	6	6	6	6	5

Table 3. Three weeks training program for reduced training load.

Groups	Last week pattern of increasing interval training All groups	Taper program		
		First taper (volume)	Second taper (frequency)	Third taper (intensity)
Training duration (min)	70	70	70	50
Rest duration (min)	25	25	25	25
Training duration (min)	2	2	2	2
Rest time duration (min)	10	7	10	10
Replication	6	6	4	6

ing up and 5 minutes for cooling down. This program was conducted in 6 weeks and each week included 6 sessions (Table 2). Following the increasing interval exercise trainings, the rats entered the taper phase in which NS supplement was used for the taper and control groups²⁷.

Preparation of hydro-alcoholic extract

Fifty five grams of NS powder was weighed with a scale of 0.001 precision and then was soaked in 30% distilled water mixed with 70% ethanol solution for 72 hours. During this period, the container of the solution was well sealed with paraffin and was kept at 20 to 25 °C room temperature. The mixture was stirred with a glassy rod every six hours. After this period, the mixture was filtered through Whatman filter paper and its solvent was removed by mild temperature rotary (under 60 °C). Control and NS tapering groups was treated by extract via oral gavage at a dose of 500 mg/kg body weight.

Tissue sampling and histological studies

At the end of six week period of increasing interval exercise training, and at the end of the first and the third week of tapering (Table 3), the

animals were euthanized with a mixture of ketamine hydrochloride (50 mg/kg) and xylazine (10 mg/kg), intraperitoneally and left lung was removed and fixed in a 10% buffered formalin solution. Lung tissue samples were dehydrated by passing through a graded series of ethanol and cleared by xylene and impregnated by paraffin. Tissue processing was done by histokinette 2000 (Lica, Germany) and samples were embedded in paraffin blocks. Then, 20 to 25 non-serial 5 µm sections from each block were obtained using rotary microtome and stained with hematoxylin-eosin (H&E). For quantitative and qualitative microscopic analysis of lung tissue, at least 10 microscopic fields from each section were examined at × 400 magnification using point counting and based on systematic uniform random strategy and unbiased stereological studies by a version 9 stereo-investigator system software (MBF Bioscience, Micro Bright Field, Inc., Germany). In each microscopic field, 0.016 mm² of lung tissue were analyzed. Inflammation index of lung tissue was evaluated using grading scale described by Braber *et al.*²⁸ based on the frequency and manner of the inflammatory cell presence. A value of 0 was assigned when no inflammation was detectable, a value of 1 was adjudged for occasional cuffing with inflammatory cells, a value of 2 when most bronchi or vessels were surrounded by a thin layer (one to

five cells thick) of inflammatory cells, and a value of 3 was given when most bronchi or vessels were surrounded by a thick layer (more than five cells thick) of inflammatory cells. Total lung inflammation was defined as the average of the peribronchial and perivascular inflammation scores²⁸. All analyses were carried out by one evaluator who was blinded to the treatment groups.

Statistical analysis

All statistical analyses were performed using the SPSS software version 21. For the analysis of normal distribution of data, Kolmogorov-Smirnov test and for the comparison of the variables among groups, a two-way analysis of variance (ANOVA) followed by LSD post hoc test were used. Descriptive statistical data expressed as mean ± SD; differences of $p \leq 0.05$ was considered as significant and the rejection of null hypothesis.

Results

The study of histological structure of lung tissue has revealed that the lung parenchyma was normal in both control and basic groups (Figure 1-A and B). Structural integrity of lung tissue in taper groups treated by *NS* was greater and better than in the taper groups without *NS*. Furthermore, among the groups with *NS*, the third week taper groups


had a better structure and a more integrated alveolus wall. The results of microscopic investigations showed that IIT had a significant effect on the lung tissue of rats during maturation period. These results indicated that the IIT caused severe inflammation in lung tissue and infiltration of inflammatory cells and lymphocytes into the connective tissue around the respiratory airway, vessels and interstitial lamellae (Figure 1-C). The taper exercises could decrease these damages in the lung tissue (Figure 1-D to 1-F). Although a small amount of emphysema, and mild interstitial inflammation was observed in some taper groups, this damage was less in the groups with *NS*, but the relative inhibitory effect of all three taper types were rather good (Figure 1-G to 1-I). Among these groups, the taper group treated with *NS* in the third week showed a better improvement in comparison to the other groups.

As it is shown in Figure 2, results indicated that the severity of lung tissue inflammation in control groups increased until the second week of tapering (age of 13 weeks) and then decreased. The severity of lung tissue inflammation in control groups without *NS* increased as similarly and significantly as in the basic and interval control groups (age of 11 weeks), ($p=0.002$). A similar and significant increase of inflammation has also been observed in the second week control animals with *NS* comparing with the basic and interval control groups ($p=0.02$). There was not a significant difference in the severity of lung tissue inflammation in the second week taper groups (age of 14 weeks) with and without *NS* comparing with the basic group (respectively $p=0.07$ and $p=0.30$).

Figure 3 indicates that the implementation of the 6- week undulating and IIT (Table 2) during maturation period caused the most severe inflammation in lung tissue (grade 3) and the occurrence of this inflammation was similarly and significantly more frequent in comparison to the control and basic groups.


The results showed that the implementation of the three kinds of taper exercise training programs (frequency, repetition, intensity) following the IIT, could significantly decrease the amount and severity of lung tissue inflammation ($p=0.001$) in the studied times in comparison to interval training (Figure 4). After two weeks of frequency, repetition and intensity tapering, the degree of lung inflammation decreased 32, 49 and 52 percent respectively compared to the interval training and this decrease continued until the third week of the tapering (51, 52 and

Figure 1. Microscopic view of lung tissue in different groups (H&E, ×200).




A) Normal lung tissue in the basic group; B) Lung tissue in the interval control group; C) IIT group with aggregation of lymphoid tissue and inflammatory cells around respiratory air way (solid arrow) and vessels (hollow arrow); D) two weeks control group without *NS*; E) two weeks frequency taper group without *NS*; F) two weeks repetition taper group without *NS*; G) three weeks frequency taper group with *NS*; H) three weeks repetition taper group with *NS*; I) three weeks intensity taper group with *NS*.

Figure 2. Comparison of lung tissue inflammation in control groups.




* and ** show a significant difference comparing with basic and interval control groups (respectively $p < 0.05$ and $p < 0.01$).

Figure 3. Comparison of lung tissue inflammation in the basic, control and increasing interval training groups.



*** shows a significant difference ($p < 0.001$) comparing with basic and interval control groups.

Figure 4. Comparison of lung tissue inflammation in interval and frequency, repetition and intensity taper groups without *N. sativa* over two and three weeks of tapering.




*** shows a significant difference ($p < 0.001$) in three type of tapering comparing with increasing interval training group. ### shows a significant difference ($p < 0.001$) in third week comparing with second week of frequency taper group.

59 percent respectively in frequency, repetition and intensity taper). The results as summarized in Figure 4 also revealed that, compared to frequency and repetition taper groups, intensity decreasing taper group acted more effectively in the reduction of lung tissue inflammation.

The evaluation of the time effect on lung tissue inflammation showed that lung tissue inflammation was reduced 27, 6 and 15 percent respectively in the frequency, repetition and intensity taper groups of the third week compared to the second week (Figure 4). This reduction was significant only in frequency taper ($p = 0.001$, $p = 0.60$ and $p = 0.195$ respectively in frequency, repetition, and intensity groups).


The evaluation of interactive effect of *NS* hydro-alcoholic extract use (as a supplement) and the implementation of different patterns of taper revealed the same amount of decrease in lung tissue inflammation as the groups without *NS* (Figure 5). The frequency, repetition and intensity groups, with a decrease of respectively 61%, 75% and 72% in the second

Figure 5. Comparison of lung tissue inflammation in interval groups and frequency, repetition and intensity taper groups treated with *N. sativa* during the taper period.



*** shows a significant difference ($p < 0.001$) in three type of tapering comparing with increasing interval training group.

Figure 6. Comparison of lung tissue inflammation in all groups of the study.



* shows a significant difference comparing with basic group. & shows a significant difference comparing with increasing interval training group. # shows a significant difference between 2nd and 3rd week of frequency taper group.

week and a decrease of 77%, 82% and 80% in the third week, showed a similar and significant decrease in lung tissue inflammation comparing with interval exercise training group ($p = 0.001$). In the survey of the time effect, a decrease of 43%, 28% and 30% in lung tissue inflammation was observed in the third week frequency, repetition and intensity taper groups compared to the second week taper groups. This change was significant only in frequency taper group ($p = 0.002$, $p = 0.195$, $p = 0.120$ respectively in frequency, repetition, intensity groups).

The results of the present study showed that the degree of lung tissue inflammation in all taper groups was significantly higher comparing to their counterpart ($p = 0.04$) and basic ($p = 0.001$) groups. As it is shown in Figure 6, the use of hydro-alcoholic extract of *NS* in the taper groups could decrease the amount of lung tissue inflammation caused by interval exercise. Lung tissue inflammation in the frequency, repetition and intensity groups treated with *NS* in the second week decreased

respectively 42%, 54%, and 50% comparing with the groups untreated with *NS* and this decrease was respectively 62%, 41% and 52% in the third week. In short, it can be concluded that the increasing interval exercise along with the implementation of intensity and repetition tapers and using the hydro-alcoholic extract of *NS* in a time period of three weeks had the most decrease in lung tissue inflammation.

Discussion

This study analyzed the microscopical changes and structural remodeling of lung tissue caused by inflammation in it through implementation of a period of increasing interval exercise followed by different patterns of tapering during maturation period. Since changes in the control groups can indicate changes in the lung tissue during maturation period, this study was designed to analyze the maturation process as well. The age range for the research protocol was 5-14 weeks. According to Sengupta¹⁴ this age range in rats is equivalent to the age range of 6-16 years old in humans. Therefore the male rats in the control groups, treated and untreated with *NS*, were in their maturation period in the second week of tapering. It is assumed that the significant increase of lung tissue inflammation in these groups is due to being in the critical period of maturation which is related to developmental changes of immune system brought about by maturation and increasing efficiency of antigenic system⁹.

Our results showed that the implementation of increasing interval exercise during maturation period causes severe lung tissue inflammation and leads the inflammatory and lymphoid cells into it. Although the amount of lung tissue inflammation in different periods and different patterns has decreased in animals both treated and untreated with *NS* hydro-alcoholic extract, but it is still significantly higher in comparison to the basic and control groups. It is concluded that this is because of the downfall of immune system which occurs following a high-intensity exercise over a long period of time⁵.

Although several different studies have been done about the influence of high intensity exercise on the immune system and the inflammatory factors in the body, but this study is unique for the microscopical analysis it makes on the influence of high intensity exercise on the immune system of lung tissue and occurrence of inflammation in it, using histological and stereological methods. A few studies have been done about the effects of high intensity exercise on immune system function. The presence of pulmonary macrophages is very essential in adjusting the acute and chronic inflammatory responses and the call of into the spreadable site of inflammation in the lung²⁹. Michna *et al.*³⁰ observed that, after a period of intense training, the immigrant peritoneal macrophages of humans and mice had a better performance in responding to the chemotactic factors. The previous studies showed that acute and chronic exercise training has an effective catalytic role in many macrophage functions. On the other hand, the increasing rate of macrophage function varies according to the intensity and duration of exercise³¹.

Previous study by Sobhani *et al.*³² showed that HIIT in maturing rats cause in airway narrowing of the lung parenchyma. Yadegari *et al.*³³ also indicated that HIIT leads to parenchymal remodeling in lung tissue by induction of inflammation. Our recent research revealed that

six weeks HIIT significantly increase number of alveolar macrophage in lung tissue³⁴. Review of Ramel *et al.*³⁵, Murphy *et al.*³⁶ and Yamamoto *et al.*³⁷ studies suggested that high intensity physical activities increases the number of neutrophils, while this increase has not been observed in low intensity physical activities.

Some studies showed that, the function of lymphocytes, in long periods, is sensitive to the increase of exercise intensity in endurance activities^{38,39}. It can be concluded that, high intensity exercises decrease the function of lymphocytes and macrophages due to an increase in the circulation of stress hormones, especially cortisol, and a change in the balance of pro-inflammatory or anti-inflammatory cytokines when responding to the exercise training³⁸. In a study conducted on 18 swimmers on the national level and 11 healthy untrained volunteers, it was found out that the number of monocytes, neutrophils and dendritic cell subsets as well as the amount of IL-1 β , IL-6, and IL-12 decreases in these athletes during the training season. The results of this study supported the idea that long-term high intensity exercise may affect the innate immune cells function, reduce their capacity in responding to acute challenges, and increase the risk of URTI⁴⁰.

Previous studies have shown that daily repeated physical activity during long periods, in athletes and especially endurance athletes, induces damage to the epithelium cells and increases the inflammation in their respiratory mucosa⁴¹. Thus, it can be concluded that despite of multiple mechanisms of innate and adaptive immunity being there, the implementation of high intensity exercise training, weakens the immune system of the lung tissue. The reported severity of inflammation in the lung tissue of IIT animals in this study also confirms this possibility.

For a reduction of disorders in immune system, physiological capacity and mood state profiles of athletes following a long-term and intensive exercise, performing a taper with a gradual reduction in the load of exercise can be recommended by the sport trainers to the athletes as an appropriate approach¹³. The results of this study showed that the implementation of taper patterns (frequency, repetition and intensity) after a period of IIT, could significantly decrease the lung tissue inflammation caused by intensive exercise training but the inflammation was still significantly higher in comparison to the control group after 3 weeks of taper. Regarding the time effect of the taper, the implementation of a three-week repetition and intensity taper was more effective in enhancing the immune system and reducing the lung tissue inflammation subsequent to intensive exercise than a two-week taper. The results of this study were in line with previous studies^{3,42}.

Mujica *et al.*⁴², having observed the trained athletes during 1-3 weeks of taper, reported enhanced performance often accompanied by increased anabolic activity, reduced physiological stress and restoration of mucosal immunity and immune function. It has also been shown that a 6-day taper in the middle-distance runners improved the performance in 800 meter runners¹⁶. Two weeks taper in triathletes⁴³, one week taper in rugby league players^{44,45} and two weeks taper in judo athletes¹¹ resulted in increased T/C ratio and improved performance. It can be concluded that the recovery or the enhancement of immune system function during taper is dependent on the amount of immune system diminution during intensive exercises³.

On the other hand, by the increase of volume and intensity of training during the pre-competitive season, sport trainers will also concern

about serious matters other than increased risk of sport injuries and URTI development. One of these concerns is the increasing tendency of athletes to take sport medicines and chemical supplements, some of which are completely ineffective in long term use³. Herbal drugs and supplements as natural treatment (complementary treatment) with fewer side effects and multiple properties can be the best alternative for athletes⁴⁶.

In the last three decades, extensive research has been done on the biological effects of *NS* seeds. In numerous scientific articles, the antioxidant, anti-inflammatory, immune booster and antihistamines properties of numerous compounds in *NS* hydro-alcoholic extract have been pointed out⁴⁷. One of the special features of *NS* is its role in regulating immune function in treadmill exercised rat²⁵. The effect of *NS* hydro-alcoholic extracts use on the reduction of lung tissue inflammation^{23,24} specially induced by intensive exercise training has also been observed in this study. While the implementation of different patterns of taper reduced the lung tissue inflammation, but *NS* hydro-alcoholic extracts use enhanced this reduction.

Although the amount of lung tissue inflammation in the animals significantly decreased in all three types of taper treated with *NS*, in comparison to their counterparts, but the implementation of three weeks of intensity and repetition taper accompanied by *NS* hydro-alcoholic extracts use, was more effective in the reduction of lung tissue inflammation induced by intensive exercise. Previous studies have indicated that some of *NS* compounds have the effect of reinforcing the cellular immunity⁴⁸. Thymoquinone's anti-inflammatory properties, the major compound of *NS* extract, works through the suppression of inflammatory mediators such as prostaglandins and leukotrienes^{49,50}.

This study is among the few studies that examine the effect of taper on lung tissue safety mechanism during maturation period. The results of this study showed that although increase of interval training intensity has been undulating and gradual but the immune system of the lung tissue is not able to cope with that and it causes severe inflammation in the lung tissue. This problem during pre-maturation period may have a negative effect on the performance of the athletes and the results of the competition or even negative effects on the future of their sport. Our results indicated that the implementation of three types of taper decreases the lung tissue inflammation induced by IIT. It can be concluded that a reduction in the load of exercise alone can compensate for the induced weakness, or enhance the immune function so that the lung tissue inflammation decreases. But since the interactive effect of *NS* hydro-alcoholic extract use and taper was more effective in the reduction of lung tissue inflammation, we can conclude that reducing the exercise load accompanied by *NS* hydro-alcoholic extract use associated with particular anti-inflammatory properties, has a more prominent role in safety mechanism of lung tissue. Thus the implementation of taper along with *NS* hydro-alcoholic extract use will enhance the immune system of the lung tissue during maturation period and subsequently reduces the induced damages. It may also make the sport life span longer. Decrease of the intensity or frequency of the exercise load is a good pattern for taper program. However the results of the study revealed that the implementation of 3 weeks taper is more effective in the reduction of lung tissue inflammation than 2 weeks taper. It is suggested in the future studies, Changes in various factors of

innate and adaptive immunity of lung tissue in pre-maturation period induced by increasing interval exercise or taper as well as the optimal taper duration in human models should be studied.

Acknowledgment

This study was financially supported by Research Council of University of Mazandaran and University of Tabriz.

Conflict of interest

The authors do not declare a conflict of interest.

Bibliography

- Martin SA, Pence BD, Woods JA. Exercise and respiratory tract viral infections. *Exerc Sport Sci Rev*. 2009;37(4):157-64.
- Nieman DC. Does Exercise Alter Immune Function and Respiratory Infections? *Pres Counc Phys Fit Sports Res Dig*. 2001;3(13):1-8.
- Farhangimaleki N, Zehsaz F, Tiidus PM. The effect of tapering period on plasma pro-inflammatory cytokine levels and performance in elite male cyclists. *J Sports Sci Med*. 2009;8(4):600.
- Moldoveanu B, Otmishi P, Jani P, Walker J, Sarmiento X, Guardiola J, et al. Inflammatory mechanisms in the lung. *J Inflamm Res*. 2009;2:1-11.
- Walsh NP, Gleeson M, Pyne DB, Nieman DC, Dhabhar FS, Shephard RJ, et al. Position statement part two: maintaining immune health. *Exerc Immunol Rev*. 2011;17:64-103.
- Neville V, Gleeson M, Folland JP. Salivary IgA as a risk factor for upper respiratory infections in elite professional athletes. *Med Sci Sports Exerc*. 2008;40:1228-36.
- Matthews CE, Ockene IS, Freedson PS, Rosal MC, Merriam PA, Hebert JR. Moderate to vigorous physical activity and risk of upper-respiratory tract infection. *Med Sci Sports Exerc*. 2002;34(8):1242-8.
- Thomas L, Busso TH. A theoretical study of taper characteristics to optimize performance. *Med Sci Sports Exerc*. 2005;37(9):1615-21.
- Ploeger HE, Takken T, De Greef MH, Timmons BW. The effects of acute and chronic exercise on inflammatory markers in children and adults with a chronic inflammatory disease: a systematic review. *Exerc Immunol Rev*. 2009;15(1):6-41.
- Khubchandani KR, Snyder JM. Surfactant protein A (SP-A): the alveolus and beyond. *FASEB J*. 2001;15(1):59-69.
- Papacosta E, Gleeson M. Effects of intensified training and taper on immune function. *Rev Bras Educ Fis Esporte*. 2013;27(1):159-76.
- Scharhag J, Meyer T, Gabriel HH, Schlick B, Faude O, Kindermann W. Does prolonged cycling of moderate intensity affect immune cell function? *Br J Sports Med*. 2005;39(3):171-7.
- Mujika I. Tapering for triathlon competition. *J Human Sport Exerc*. 2011; 6(2): 264-70.
- Sengupta P. A scientific review of age determination for a laboratory rat: how old is it in comparison with human age. *Biomed Int*. 2011;2(2):81-9.
- Mujika I. Intense training: the key to optimal performance before and during the taper. *Scand J Med Sci Sports*. 2010;20(52):24-31.
- Mujika I, Goya A, Ruiz E, Grijalba A, Santisteban J, Padilla S. Physiological and performance responses to a 6-day taper in middle-distance runners: influence of training frequency. *Int J Sports Med*. 2002;23(5):367-73.
- Shepley B, MacDougall JD, Cipriano N, Sutton JR, Tarnopolsky MA, Coates G. Physiological effects of tapering in highly trained athletes. *J Appl Physiol*. 1992;72(2):706-11.
- Gleeson M. Mucosal immunity and respiratory illness in elite athletes. *Int J Sports Med*. 2000;21(Sup. 1):33-43.
- Bosquet L, Montpetit J, Arvaisis D, Mujika I. Effects of tapering on performance: a meta-analysis. *Med Sci Sports Exerc*. 2007;39(8):1358-65.
- Randhawa MA, Alghamdi MS. Anticancer activity of *Nigella sativa* (black seed) - a review. *Am J Chin Med*. 2011;39(06):1075-91.
- Salem ML. Immunomodulatory and therapeutic properties of the *Nigella sativa* L. seed. *Int Immunopharmacol*. 2005;5(13):1749-70.
- Ali BH, Blunden G. Pharmacological and toxicological properties of *Nigella sativa*. *Phytother Res*. 2003;17(4):299-305.

23. Kanter M. Effects of *Nigella sativa* seed extract on ameliorating lung tissue damage in rats after experimental pulmonary aspirations. *Acta Histochemica*. 2009;111(5):393-403.
24. Abidi A, Robbe A, Kourda N, Ben Khamsa S, Legrand A. *Nigella sativa*, a traditional Tunisian herbal medicine, attenuates bleomycin-induced pulmonary fibrosis in a rat model. *Biomed Pharmacother*. 2017;90:626-37.
25. Gholamnezhad Z, Boskabady MH, Hosseini M. Effect of *Nigella saiva* on immune response in treadmill exercised rat. *BMC Complement Altern Med*. 2014;14(1):437.
26. Mirdar Sh, Arabzadeh A, Arzani A, Ahmadi S, Neyestani F, Baqbani M. The comparison time periods and different patterns of taper with *Nigella Sativa* supplementation on body weight changes and endurance performance in male wistar rats during of maturity. *J Appl Exerc Physiol*. 2015;10(20):115-28. [full text in persian]
27. Mirdar Sh, Arabzadeh E, Hamidian Gh. Effects of two and three weeks of tapering on lower respiratory tract in the maturing rat. *Koomesh*. 2015;16(3):366-75. [full text in persian]
28. Braber S, Henricks PA, Nijkamp FP, Kraneveld AD, Folkerts G. Inflammatory changes in the airways of mice caused by cigarette smoke exposure are only partially reversed after smoking cessation. *Respir Res*. 2010;11(1):99.
29. I-Ghamdi MS. The anti-inflammatory, analgesic and antipyretic activity of *Nigella sativa*. *J Ethnopharmacol*. 2001;76(1):45-8.
30. Michna H. The human macrophage system: activity and functional morphology. *Bibl anat*. 1988;31:1-84.
31. Davis JM, Murphy EA, Brown AS, Carmichael MD, Ghaffar A, Mayer EP. Effects of moderate exercise and oat β -glucan on innate immune function and susceptibility to respiratory infection. *Am J Physiol Regul Integr Comp Physiol*. 2004;286(2):R366-72.
32. Sobhani V, Mirdar S, Arabzadeh E, Hamidian Gh, Mohammadi F. High-intensity interval training-induced inflammation and airway narrowing of the lung parenchyma in male maturing rats. *Comp Clin Path*. 20017;72(3):577-82.
33. Yadegari M, Mirdar S, Hamidian Gh. The effect of high-intensity interval training on lung parenchymal and non-parenchymal structural changes. *Daneshvar Med*. 2016;23(124):51-60. [full text in Persian]
34. Mirdar Sh, Naiestany F, Hamidian Gh, Hedayati M. Increment of alveolar macrophages and pulmonary surfactant of young male rats after six weeks interval training. *Sport Physiol*. 2018;9(36):59-72. [full text in Persian]
35. Ramel A, Wagner KH, Elmadfa I. Correlations between plasma noradrenaline concentrations, antioxidants, and neutrophil counts after submaximal resistance exercise in men. *Br J Sports Med*. 2004; 38(5): e22.
36. Murphy EA, Davis JM, Brown AS, Carmichael MD, Ghaffar A, Mayer EP. Oat beta-glucan effects on neutrophil respiratory burst activity following exercise. *Med Sci Sports Exerc*. 2007;39(4):639-44.
37. Yamamoto Y, Nakaji S, Umeda T, Matsuzaka M, Takahashi I, Tanabe M, et al. Effects of long-term training on neutrophil function in male university judoists. *Br J Sports Med*. 2008;42(4):255-9.
38. Lancaster GI, Halson SL, Khan Q, Drysdale P, Wallace F, Jeukendrup AE, et al. Effects of acute exhaustive exercise and chronic exercise training on type 1 and type 2 T lymphocytes. *Exerc Immunol Rev*. 2004;10(91):91-106.
39. Simpson RJ, Cosgrove C, Ingram LA, Florida-James GD, Whyte GP, Pircher H, et al. Senescent T-lymphocytes are mobilised into the peripheral blood compartment in young and older humans after exhaustive exercise. *Brain Behav Immun*. 2008;22(4):544-51.
40. Silva RA, Vieira RP, Duarte AC, Lopes FD, Perini A, Mauad T, et al. Aerobic training reverses airway inflammation and remodeling in an asthma murine model. *Eur Respir J*. 2010;35(5):994-1002.
41. Carlsen KH. The breathless adolescent asthmatic athlete. *Eur Respir J*. 2011; 38: 713-20.
42. Mujika I, Padilla S, Pyne D, Busso T. Physiological changes associated with the pre-event taper in athletes. *Sports Med*. 2004;34(13):891-927.
43. Coutts AJ, Wallace LK, Slattery KM. Monitoring changes in performance, physiology, biochemistry, and psychology during overreaching and recovery in triathletes. *Int J Sports Med*. 2007;28(02):125-34.
44. Coutts A, Reaburn P, Piva TJ, Murphy A. Changes in selected biochemical, muscular strength, power, and endurance measures during deliberate overreaching and tapering in rugby league players. *Int J Sports Med*. 2007;28(02):116-24.
45. Coutts AJ, Reaburn P, Piva TJ, Rowsell GJ. Monitoring for overreaching in rugby league players. *Eur J Appl Physiol*. 2007;99(3):313-24.
46. Fong HH. Integration of herbal medicine into modern medical practices: issues and prospects. *Integr Cancer Ther*. 2002;1(3):287-93.
47. Edris AE. Anti-cancer properties of *Nigella* spp. essential oils and their major constituents, thymoquinone and β -elemene. *Curr Clin Pharmacol*. 2009;4(1):43-6.
48. Haq A, Abdullatif M, Lobo PI, Khabar KS, Sheth KV, Al-Sedairy ST. *Nigella sativa*: effect on human lymphocytes and polymorphonuclear leukocyte phagocytic activity. *Immunopharmacol*. 1995;30(2):147-55.
49. Ghayur MN, Gilani AH, Janssen LJ. Intestinal, airway, and cardiovascular relaxant activities of thymoquinone. *Evid Based Complement Alternat Med*. 2012; Article ID: 305319, 13 pages.
50. Modaresi M, Poor-naji N. The effect of black seed (*Nigella sativa*) hydro-alcoholic extract on breeding factors in female mice. *J Shahrekord Univ Med Sci*. 2012;13(6):63-70. [full text in persian].

Espíritu **UCAM** Espíritu Universitario

Miguel Ángel López

Campeón del Mundo en 20 km. marcha (Pekín, 2015)
Estudiante y deportista de la UCAM



- **Actividad Física Terapéutica** ⁽²⁾
- **Alto Rendimiento Deportivo:**
 - Fuerza y Acondicionamiento Físico** ⁽²⁾
- **Performance Sport:**
 - Strength and Conditioning** ⁽¹⁾
- **Audiología** ⁽²⁾
- **Balneoterapia e Hidroterapia** ⁽¹⁾
- **Desarrollos Avanzados**
 - de Oncología Personalizada Multidisciplinar** ⁽¹⁾
- **Enfermería de Salud Laboral** ⁽²⁾
- **Enfermería de Urgencias,**
 - Emergencias y Cuidados Especiales** ⁽¹⁾
- **Fisioterapia en el Deporte** ⁽¹⁾
- **Geriatría y Gerontología:**
 - Atención a la dependencia** ⁽²⁾
- **Gestión y Planificación de Servicios Sanitarios** ⁽²⁾
- **Gestión Integral del Riesgo Cardiovascular** ⁽²⁾
- **Ingeniería Biomédica** ⁽¹⁾
- **Investigación en Ciencias Sociosanitarias** ⁽²⁾
- **Investigación en Educación Física y Salud** ⁽²⁾
- **Neuro-Rehabilitación** ⁽¹⁾
- **Nutrición Clínica** ⁽¹⁾
- **Nutrición y Seguridad Alimentaria** ⁽²⁾
- **Nutrición en la Actividad Física y Deporte** ⁽¹⁾
- **Osteopatía y Terapia Manual** ⁽²⁾
- **Patología Molecular Humana** ⁽²⁾
- **Psicología General Sanitaria** ⁽¹⁾

⁽¹⁾ Presencial ⁽²⁾ Semipresencial

Androgens from physiology, through pharmacy and pharmacology to the status of lifestyle drugs - are we going in the right direction?

Nataša Duborija-Kovačević¹, Mitar Popović¹, Rade Kovač²

¹Department of Pharmacology and Clinical Pharmacology, University of Montenegro, Podgorica, Montenegro. ²Special Hospital for Lung Diseases "Brezovik", Nikšić, Montenegro.

Received: 10.09.2018

Accepted: 31.01.2019

Summary

The aim of this review was to evaluate the current cognition about androgens (A) physiology, their pharmaceutical development and place in modern medicine. Special aspect was to explore the reasons and consequences of A use as so-called "lifestyle drugs" (LD).

To write this review, we used the scientific papers in English of a recent date on PubMed, reference textbooks, books and monographs of different disciplines, as well as official documents and reports of some internationally recognized organizations (European Medicines Agency, World Anti-Doping Agency, Medicines and Healthcare products Regulatory Agency).

Endocrinological role of A is generally known, but their non-hormonal effects are still the subject of intensive investigation. For decades, testosterone (T) and its esters have been the substitution therapy of the first choice in clearly defined clinical conditions. When it comes to pharmaceutical development, there are large number of effective and safe T formulations on the market which provide a very good patients' compliance. Regarding clinical application of synthetic A with dominant anabolic activity, the only acceptable indication nowadays is severe burn injuries, while others have to be proven by high-quality clinical studies. Particularly worrying is the wide-spread use of anabolic steroids (AS) for non-medical purposes, as so-called LD. Although numerous and serious side-effects are well-documented, general impression is that both users of AS and clinicians should know more about the risks of their use.

This review points to the need of better information and more comprehensive education at different levels, as well as implementation of additional preventive strategies, especially in the youth population, in order to avoid potentially serious consequences of the AS use.

Key words:

Androgens. Physiology.
Pharmacy. Pharmacology.
Lifestyle drugs.

Andrógenos, de la fisiología, a través de la farmacia y la farmacología al estado de las drogas de vida, ¿vamos en la dirección correcta?

Resumen

El objetivo de esta revisión fue evaluar el conocimiento actual sobre la fisiología de los andrógenos (A), su desarrollo farmacéutico y su lugar en la medicina moderna. Un aspecto especial fue estudiar las razones y consecuencias del uso de llamadas "medicamentos de estilo de vida" (LD).

Para llevar a cabo esta revisión, se realizó una búsqueda de artículos científicos en inglés recientes en PubMed, libros y monografías de diferentes disciplinas, así como documentos oficiales e informes de algunas organizaciones reconocidas internacionalmente (Agencia Europea de Medicamentos, Agencia mundial Antidopaje, Agencia Reguladora de Medicamentos y Productos Sanitarios).

El papel endocrinológico de los A es generalmente conocido, pero sus efectos no hormonales siguen siendo objeto de una investigación intensiva. Durante décadas, la testosterona (T) y sus ésteres han sido la primera elección para terapia de sustitución en condiciones clínicas claramente definidas. Cuando se trata del desarrollo farmacéutico, hay una gran cantidad de formulaciones de T eficaces y seguras en el mercado que proporcionan un muy buen cumplimiento por parte de los pacientes. Con respecto a la aplicación clínica de A sintético con actividad anabólica dominante, la única indicación aceptable en la actualidad son las lesiones por quemaduras graves, mientras que otras deben ser probadas por estudios clínicos de alta calidad. Particularmente preocupante es el uso generalizado de los esteroides anabólicos (AS) para fines no médicos, como los llamados LD. Aunque los efectos secundarios son numerosos y graves, la impresión general es que tanto los usuarios de AS como los clínicos deberían saber más sobre los riesgos de su uso.

Esta revisión apunta a la necesidad de una mejor información y una educación más integral en diferentes niveles, así como la implementación de estrategias preventivas adicionales, especialmente en la población joven, para evitar consecuencias potencialmente graves del uso de AS.

Palabras clave:

Andrógenos. Fisiología. Farmacia.
Farmacología. Medicamentos
de estilo de vida.

Correspondence: Natasa Duborija-Kovacevic

E-mail: duborijan@ac.me

Introduction

Male sex hormones or androgens (A) and their role in the human body are being learnt from elementary school. Later, at schools and faculties of health care, students learn about diseases and conditions that result from their insufficient or excessive secretion. Medical doctors and pharmacists know most about A as medicines, while other professions, as well as the general public, know very superficially or to the extent of their own needs or interests. As with many other medicines, some indications for A use are unquestionable and generally accepted from both clinicians and regulatory bodies, while others are still waiting for valid evidence.

However, unlike most other drugs it is well-known that different substances with androgenic actions were used by people without any medical diagnosis, in order to increase muscle mass, strength and endurance¹⁻⁴. Available data show that those were first German soldiers¹, later Russian and American athletes^{2,3}, and since the eighties of the last century these substances have "entered" into the widest population of young and middle-aged individuals whose physical appearance and muscle mass have become the basic life preoccupation⁴. In official literature these substances got the status of lifestyle drugs (LD)⁵. Possible risks and consequences of their use, including cases of death, are well-documented⁶⁻⁸. The extent and importance of this problem is indicated by the data that in some countries, such as Netherlands, an outpatient clinic for past and current users of those substances was established in 2011 in order to acquire more knowledge about the health risks associated with anabolic androgenic steroids abuse⁹.

The objective of this review was to evaluate critically and in detail the current cognition about A physiology, their pharmaceutical development and place in modern medicine. Special aspect was to explore the reasons and potential consequences of their use for non-medical purposes as so-called LD and to point out the importance of better education and preventive strategies in this area, especially in youth population.

Methods

Various literary sources were used to write this review. Initial data were obtained from textbooks, monographs and other books in pharmacology and clinical pharmacology, physiology, pharmacy, endocrinology and various sports sciences. The inclusion criteria for these publications were the following: a publication of a newer date which is internationally recognized (or used in university teaching), written in English and issued by a renowned publisher. Older date publications that are not of an international character and not published in English were not taken into account. For a more detailed insight into individual parts of this review, we searched the "PubMed" database, using a number of keywords and determinations. Here the criteria for inclusion were the following: publication had to be original scientific paper or review (including studies on animals and other types of experimental studies) published in English in the period 2000-2018. Priority was given to the results of randomized, controlled, double-blind clinical studies with a large number of participants. In exceptional cases, older papers were

quoted, where this was of substantial importance for the research subject. Case studies, papers without the author's name and papers which weren't written in English were not taken into consideration. The last source of information was official websites of internationally recognized regulatory institutions (European Medicines Agency-EMA, Medicines and Healthcare products Regulatory Agency-MHRA and World Anti-Doping Agency-WADA) from which the latest reports, recommendations and documents were taken. The authors accessed these sites for the last time in November 2018.

Androgens as physiologically important substances - can we live without them?

Testosterone (T) is a major natural A. It is mostly synthesized in testicular Leydig's cells and smaller amounts in the ovaries and adrenal cortex from cholesterol as the initial substance. In a healthy adult male, 4-9 mg of T is excreted daily and only 1-2% is free in plasma¹⁰.

Androgenic effects of T are spermatogenesis and development of primary and secondary sexual characteristics of a man, including characteristic hair distribution on the body, baldness, a specific voice depth, increased skin thickness and firmness of the subcutaneous tissue, increased secretion of the sebaceous glands with the appearance of acne, etc. Anabolic effects of T include protein synthesis and muscle development, bone growth and calcium storage, increased basal metabolism, increased number of erythrocytes, and others¹¹. Behavioral effects of T include the regulation of sexuality, aggression, cognition, emotion, and personality^{12,13}.

T precursors, dehydroepiandrosterone (DHEA) and androstenedion are less potent A which are synthesized in the gonads and adrenal cortex in both sexes. In the liver they are being converted into more potent T¹⁴. DHEA and its metabolites regulate glandular and neurotransmitter secretions, influence glucose homeostasis and cyclic release of GnRH, control the activity of skeletal and smooth muscle and increase the tolerance to ischemia¹⁵. Additional neuroprotective effects, positive modification of human mood, emotions and behavior were also found¹⁶. Although in recent decades the popularity of these substances has grown because of their use in doping¹⁷, existing data doesn't confirm their effectiveness regarding the lean body mass, muscle strength or performance improvement compared with placebo^{18,19} (Figures 1 and 2).

In most peripheral tissues T is being converted in more active dihydrotestosterone (DHT) by 5 α -reductase. Inhibitors of this enzyme (finasteride and dutasteride) have being traditionally used in the

Figure 1. Structural formula of dehydroepiandrosterone.





Figure 2. Structural formula of androstenedione.

treatment of benign prostatic hyperplasia. Their use reduces the incidence of low-grade prostatic carcinoma in high-risk patients^{20,21}. Recent studies in experimental animals have found some non-hormonal effects of these medicines, such as reduction of dyskinesia in Parkinson's disease, anti-nociceptive and anti-inflammatory actions and the improvement of motor, EEG and cellular changes in hepatic encephalopathy²²⁻²⁴. This could expand the existing knowledge about A physiology and open new possibilities in the treatment of some CNS diseases. Paba S, *et al.*²⁵ identified the 5 α -reductase as possible novel therapeutic target in the treatment of schizophrenia.

Androgens as medicines - is the science fully dedicated exclusively to the well-being of the patient?


There have always been pathological states that have caused a lack or complete absence of A in the body of a man, with accompanying clinical manifestations. In that sense, it was imposed the need for designing and synthesizing substances which could, if applied exogenously, replace their deficiency and improve clinical symptoms. According to recent data, approximately 2.4 million males in USA aged 40-69 years old suffer from hypogonadism²⁶. About 2.3% of men in their 40s and 3.8% of men in their 60s were taking some form of T replacement therapy in 2011²⁷.

Testosterone and testosterone esters

Knowing the fact that T administered orally is the subject of so-called "metabolism of the first pass" through the liver to a significant extent¹⁴, Butenandt and Ruzicka synthesized T in the injection for the first time in 1935, that is considered as the beginning of the golden age of steroid chemistry. Four years later these scientists got the Nobel Prize in Chemistry¹. However, T administered even in this way had a short duration of action due to intense biotransformation (half-life of 10 minutes)²⁸, so patients were forced to receive it frequently.

In order to ensure the prolonged action of T, the science went on. In the 1940s and 1950s, the first T esters were synthesized (propionate, cypionate, enanthate). These esters are still in use and they are administered as intramuscular depot injection at intervals of 1-6 weeks, according to patients individual requirement^{29,30}.

In the mid-seventies there was a new progress. It was synthesized T undecanoate, also T ester in the form of depot intramuscular injection¹⁰. It has provided stable plasma levels of T over three months, which has been a significant advantage for patients³¹ (Figure 3).

Figure 3. Structural formula of testosterone undecanoate.

Certainly, the revolutionary discovery of the 1980s was transdermal application of T as an alternative to painful injection²⁸. The first effective transdermal patches were applied in the scrotum area because of very good pharmacokinetics and potency^{10,29}. Smaller skin surface area and application challenges (hair clipping), as well as the appearance of transdermal patches which could be applied at back, abdomen, upper arms or thighs limited the use of formulations for scrotum application²⁹.

Further discoveries went in the direction of hydroalcoholic gels/liquid solutions for transdermal absorption¹⁰. In order to minimize transfer, recommended sites for their application are areas which will be covered by clothing²⁹. There are also formulations for application to buccal mucosa, from which T is gradually absorbed, bypassing hepatic metabolism¹⁰. Modern T implants for subcutaneous administration provide the replacement of T for a period of even 6 months²⁹. The latest discovery was nasal gel, as another non-invasive treatment²⁹.


Synthetic androgens

There were three main reasons for design and synthesis of synthetic androgens (SA): the possibility of oral administration, the prolongation of biological activity in vivo and the increase of anabolic compared to androgenic effects³². Those modifications were expected to improve the possibility of clinical application of SA, that can be considered justified from the professional, scientific and ethical point of view.

Oral activity was achieved by substitution of the 17 α -H on the steroid nucleus with a methyl or ethyl group, which prevented the deactivation by first-pass metabolism in the liver¹². Methyltestosterone was the first synthesized medicine in this group (1935), and others were synthesized later¹⁷. However, the attempt to introduce these medicines into the long-acting replacement therapy (LART) was unsuccessful because of their hepatotoxicity¹⁷.

Another form of structural modification of A was 5 α -reduction. Mesterolone was synthesized in this way, but its oral administration in LART was limited due to relatively weak pharmacological activity¹² and unconfirmed efficacy in standard doses³³.

Nandrolone (N) is 19-demethylated analogue of T which also cannot be taken orally because of extensive first-pass metabolism in the liver³⁴. However, the third goal (very high ratio of anabolic to androgenic action¹² was achieved with N, that could be an explanation for its current status of the most popular A in sports doping and bodybuilding³⁵ (Figure 4).

Figure 4. Structural formula of nandrolone.

In order to enable its medical application, there were developed long-acting pro-drug esters of N (but not for oral use, for intramuscular injection again!), such as N decanoate (ND). Both experimental and clinical studies confirmed positive effects of ND on bone tissue. Li X *et al.* found ND blocked bone loss by inhibition of bone resorption in ovariectomized rats with osteopenia³⁶. Newer investigation on young adult rats found a positive effect of ND on bone callus formation after a complete femoral fracture³⁷. In women with postmenopausal osteoporosis ND increases bone density and reduces the incidence of fractures^{38,39}, but according to reference recommendations, such as UK Clinical Guideline for the Prevention and Treatment of Osteoporosis⁴⁰, bisphosphonates are considered as drugs of the first choice.

Androgens as medicines - what does the evidence-based medicine say?

Substitution therapy

T is used as substitution therapy of the first choice in conditions which cause primary testicular insufficiency, such as bilateral anorchia, Klinefelter's karyotype (XXY), surgical removal of the testes, chemotherapy and radiotherapy, but also in secondary testicular insufficiency which occurs as the result of disturbance on the hypothalamic-pituitary axis. Another reason for T clinical use is delayed puberty in boys ages 16 and older¹⁰.

Substitution of T in aging men in andropause, according to some authors⁴¹⁻⁴³ has a positive effect on bone mineral density and lean mass, sexual function, mood, general sense of well-being and reduction of the amount of adipose tissue. Considering this issue in recent review, Singh P concluded that "disparate results of clinical trials suggest an incomplete picture of complex interaction between aging and A deficiency". Before commencing T treatment, he suggested clear discussion with the patient about potential benefits and risks of the therapy, as well as to consider the assessment of prostate and other risk factors⁴⁴.

Controversies regarding the diagnosis of female hypoandrogenism and possible T substitution are still unresolved⁴⁵. T therapy is not approved for women in North America⁴⁶. In EU, T-containing medicines are licensed only for the treatment of male hypogonadism⁴⁷. Additionally, the long-term side effects of T in women have not yet been fully investigated^{48,45}.

Due to relatively weak anabolic effect, as well as adverse A effects, T is rarely used in catabolic conditions. The exception is hypogonadism as the consequence of the AIDS¹⁷. Rabkin JG *et al.* found in their double-blind, placebo-controlled trial that T is effective and well tolerated in the short-term treatment of clinical hypogonadism in men with symptomatic HIV illness, restoring libido and energy, alleviating depressed mood and increasing muscle mass⁴⁹. Long-term studies regarding this issue are still missing.

Pharmacological therapy

Except in physiological doses that are used in replacement therapy, A may be administered in considerably higher doses as pharmacological therapy. This type of therapy was mainly used in some non-reproductive diseases in order to improve the patients' quality of life (QoL) by enhancing muscle, bone, or other A-sensitive functions, but without ability to influence the course of underlining disease³⁵. This type of A therapy was mentioned in the literature mostly from the last decades of the XX century for several medical indications, including anemia as a result of bone marrow aplasia or renal failure⁵⁰⁻⁵¹, osteoporosis³⁸⁻³⁹, advanced ER-positive breast cancer⁵²⁻⁵³, endometriosis⁵⁴⁻⁵⁵, burn injuries⁵⁶⁻⁵⁷, and others^{35,58}. Woerdeman J, *et al.*⁵⁹ analyzed relevant clinical studies with anabolic androgenic steroids (AAS) in the treatment of non-hormonal chronic disorders published in the period 1950-2010 and they concluded that although the beneficial effects of AAS were promising, clinically relevant endpoints such as QoL, improved physical functioning and survival were mainly missing or not significant, except for burn injuries. Both Woerdeman⁵⁹ and Handelsman³⁵ pointed to the need for additional, high-quality clinical studies with the exact clinical end points in order to confirm the long term safety and efficacy of AAS in other non-hormonal clinical conditions. An additional reason for current status of A as adjuvant, supportive or adjunctive second-line therapy is the introduction of newer, more selective and more effective medicines in these indications (erythropoietins for anemia in renal failure, bisphosphonates for osteoporosis, etc.).

After the previously stated, it can be concluded that modern, evidence-based medicine clearly positions T and its esters as the first-line therapy in LART in men, but when it comes to the pharmacological application of SA with dominant anabolic activity, the only reasonable indication is severe burn injuries, while others have to be proven.

Anabolic steroids - medical indications as a goal or a justification?

According to Thieme D, *et al.*¹, anabolic steroids (AS) were first synthesized by German scientists during the Second World War, shortly after the discovery of T itself. They are believed to have been tested on people, especially prisoners, but those results have never been published. Some personal war experiences of the surviving German soldiers speak of how they were given AS in order to increase the strength and aggressiveness. Adolf Hitler also used steroids in order to strengthen his aggressive personality, that was confirmed by notes of his doctor¹. Thus, the first use of AS in healthy people and for non-medical purposes was

in German soldiers during the Second World War in order to increase their strength, endurance and aggressiveness.

The use of AS in athletes was noted for the first time at the weight-lifting championships in Vienna in 1954 by Russian competitors². Two years later CIBA Laboratories in Basel synthesized metandienone (Dianabol®) for the US Olympic Team³. In following decades A abuse quickly spread into competition bodybuilding, track and field events and other sports where performance is dependent on muscle strength or speed of recovery during training⁶⁰.


However, at the beginning of 1980s, the conditions were changed globally and AS gradually shifted from top-class sports to the general population⁴. According to Kanayama G, *et al.*⁴, a typical "user" of AS nowadays is young to middle-aged man whose motivation for their use is primarily personal appearance and muscle mass. It is clear that the use of those substances in this context is not caused by any existing medical diagnosis.

Although 23 different definitions of LD can be found in the literature⁶¹, the most operational one is that "LD are drugs that could modify or change non-medical or non-health-related goal or conditions at the margins of health and wellbeing"^{62,63}. The examples of LD should be sildenafil citrate in the treatment of erectile dysfunction, minoxidil for the baldness, botulinum toxin for "ironing" wrinkles, etc.⁶¹ Satisfying the criteria of LD definition, AS "found their place" among substances of the modern times which alter not only the personal appearance, but also the physical and mental capabilities. Although all of these substances have the status of LD, a significant difference should be noticed: erectile dysfunction and male-pattern hair loss are official medical diagnoses that are treated by licensed medicines, but facial wrinkles and personal dissatisfaction with the amount and tonus of muscle mass in own body are not yet, that doesn't mean that they will not be soon. Or, maybe they actually exist, but we still don't know their name(s).

Designer steroids - did things get out of control?

Designer steroids (DS) are AAS which are synthesized from a known parent steroid and chemically modified with the intent to circumvent controlled substances laws⁶⁴⁻⁶⁵. They are often identical to physiological substances and/or their metabolites whose concentrations in human plasma otherwise vary significantly, which makes it difficult to detect them⁵. The first DS, tetrahydrogestrinone (THG) was discovered in 2003 in the Olympic Analytical Laboratory at the University of California, Los Angeles⁶⁶ (Figure 5).

Figure 5. Structural formula of tetrahydrogestrinone.



The disappointing thing is that, unlike the previously described AS, the discovery and development of DS have never been linked (in theory or practice) with any medical indication, which could justify their design and synthesis to a certain extent, at least from the ethical point of view. Cheating athletes have a double motivation to use DS: to achieve performance enhancement and to escape from testing positive in anti-doping tests. Both can be considered as deception in sport and human sense. The fact that the first and probably many other DS were synthesized in laboratories belonging to universities could be considered as the deception of science. In their discovery and development are working persons (scientists) for which the ethics and the truth should be the basic principles of their professional dealing. An increasing number of DS on the market which are mainly sold as dietary supplements⁶⁷ should be the next fraud in sequence. "Bypassing" a strong regulatory rules that refer to licensed medicines, DS become more easily accessible and potentially more dangerous for end-users.

According to Pope HG, *et al.* (2004), AAS users rated physicians as no more reliable source of information about those substances than their friends, Internet sites, or the persons who sold them the steroids⁶⁸. On the other side, most general practitioners had some contact with AAS users in their practice, but only 40% of them reported that they would be willing to provide harm minimization advice⁶⁹.

Previously stated indicates the seriousness and complexity of this phenomenon globally, as well as the need for additional effective actions at different fields.

How anabolic steroids act on healthy human organism - do we know possible consequences?

For years, AS have been the most commonly detected doping substances in athletes⁷⁰. In 2015, according to the WADA report, AS participated exactly 50% in all positive findings on doping⁷¹. In combination with training and high protein intake, AS undoubtedly increase muscle mass and body weight, but there is little evidence that they increase muscle strength more than it would have been possible by the training itself, or that their use would improve sports results in general⁵.

Given the disturbing fact that in the last decades AS have been increasingly used to improve the physical appearance of young people who are not primarily athletes⁴ for aesthetic, psychological, sociological and other non-medical and non-sporting reasons, then we can state regrettably that, in this sense, these substances are likely to satisfy the needs of their users, at least for a limited period. On the other hand, since they are applied for a long time in supra-physiological doses, which are usually up to 50-100 times higher compared to endogenously produced T in an adult healthy male⁷², their application is associated with long-term and serious adverse effects, of which cardiovascular, neuroendocrine and psychiatric are dominant⁶⁷. The entity of side effects depends on the sex, the dose, the duration of treatment, whether AS are taken during exercise training or under sedentary conditions, and individual susceptibility to A exposure⁷³.

Adverse *cardiovascular disorders* include hypertension, an increase of LDL and reduced HDL, potentiation of thrombosis and cardiac arr-

hythmias, cardiomyopathy, left ventricular hypertrophy and myocardial ischemia^{74,75}. Melchert and Welder⁷⁶ categorized cardiovascular effects of AS in four types of actions: vasospastic, atherogenic, thrombotic and direct myocardial injuries. Frati P, et al. (2015) identified nineteen sudden cardiac deaths (SCD) in AS users in the period 1990-2012⁸, which probably could be attributed to an imbalance of autonomous nervous system activity⁷⁷. Some authors suggest that in the cases of SCD in athletes the use of AS should be excluded⁷⁴.

The *endocrinological effects* of AS depend on the age and sex of the person using them¹⁴. In adult males, anabolic effects may be accompanied by salt and water retention. The skin becomes thick and often darker, the sebaceous glands become more active, resulting in the appearance of acne¹⁴. AS use causes gynecomastia, priapism, hypogonadism, inhibits spermatogenesis and reduces the man's fertility⁷⁸. These influences adversely affect the mood, that forces a man to take these substances repeatedly. BPH and prostate cancer in AS users were also described⁷⁸. In women AS cause hairiness, male pattern baldness, menstrual cycle disorders, breast atrophy, changes in voice depth, acne, etc.^{14,78}

At the *psychological plane*, AS cause a sense of well-being, increase competitive spirit and aggressiveness, in some cases even to the level of psychosis¹⁴. Violence, aggression and impulsivity are explained by some authors as the consequence of decrease in the functional correlation of key centers in the brain responsible for emotional and cognitive behavioral regulation⁷⁹. Depression which usually requires long-term psychiatric treatment develops very often at the end of applied "cycle" of AS⁵. A numerous cases of suicide in AS users with some specificities in comparison to suicides caused by psychoactive substances were described^{80,81}. AS users more easily take other substances in order to increase muscle mass, reduce fat tissue and improve body appearance⁸². These substances, so-called "body image drugs" include other hormones (growth hormone, insulin), beta-agonists (clenbuterol), stimulants (amphetamine, ephedrine), laxatives, diuretics, etc.⁸² Even greater problem is that AS users are more likely to reach to so-called "street" drugs, predominantly opiates, compared to people who do not use AS⁸³. Particularly worrying are the results of Arvary and Pope⁸⁴ according to which 9% of male sex heroic users began using it during the use of AS, that as many as 81% of respondents bought opiate for the first time from the same dealer who sold them AS, and 67% of the respondents took opiates in order to fight with AS abstinence syndrome. Ten years ago, Graham et al. accused the medical profession who didn't accept the fact that AAS use dependency is a psychiatric condition⁸⁵.

Adverse effects of AS on the liver include adenomas, hepatocellular carcinoma, cholestasis and peliosis hepatis^{86,87}. Bond P, et al.⁸⁸ assume oxidative stress as the causative factor of AS-induced hepatotoxicity.

At the end of this chapter, the question arises is there the need to use other medicines in order to cure the "lifestyle improvement" caused by AS?

Conclusions

The hormonal effects of A are generally known in the professional, but also in general population, through various ways and levels of formal and informal education and experience. Scientists' attention is

now occupied by non-hormonal effects of A, metabolites and enzymes involved in their biotransformation. These substances and enzymes have been shown to influence the various aspects of human functioning, primarily the CNS, that opens up the new possibilities in the treatment of certain diseases.

In clinical terms, T and its esters are clearly positioned as the first line replacement therapy in male hypoandrogenism, while T-containing medicines, in the absence of effective evidence, are still not licensed for women. When it comes to the type and selection of T pharmaceutical formulations, it can be concluded that there is a fairly wide selection of effective and safe preparations, both traditional and modern, which ensures a good patients compliance and minimal negative impact on their QoL.

When SA are concerned, it can be concluded that in the pharmacokinetic sense their discovery and development fulfilled the expectations, but in clinical terms it is still not the case. Although there are undoubtedly promising results, scientific and professional community are still expecting the outcomes of high-quality clinical studies in order to confirm the long-term efficacy and safety of SA in most non-hormonal clinical conditions they are proposing.

However, thanks to their anabolic properties, SA have long been present in the population of healthy people without medical diagnosis. First of all they were soldiers, and then athletes. In the past decades, tight muscle and physical appearance were imposed as superior living values, which SA "have dropped" into the population of "ordinary" people around the world and give them the status of LD in reference literature. It went so far that nowadays DS are synthesized and used as dietary supplements without any medical reason, in order to deceive in achieving sports results and to circumvent the institutional control, that is disappointing from many aspects.

Existing research shows that both healthcare professionals and AS users don't know enough about them. Users don't have too much trust in doctors, and doctors admit that they don't know enough about these substances. Although they have the status of LD, numerous side effects, including deaths, warn and point to the need for development of additional comprehensive measures and clear strategies at different levels in order to prevent or minimize potential risks and consequences of AS abuse.

Conflict of interest

The authors do not declare a conflict of interest.

Bibliography

1. Muller RK. History of Doping and Doping Control. In: Thieme D, Hemmersbach P, editors. *Doping in Sports*. Berlin Heidelberg: Springer-Verlag 2010:1-18.
2. Wade N. Anabolic Steroids: Doctors Denounce Them, but Athletes Aren't Listening. *Science* 1972;176(4042):1399-403.
3. McDevitt ER. Ergogenic drugs in sports. In: DeLee J, Drez D, eds. *Orthopaedic Sports Medicine: Principles and Practice*. 2nd ed. Philadelphia, PA: WB Saunders 2003:471-83.
4. Kanayama G, Pope HG Jr. History and epidemiology of anabolic androgens in athletes and non-athletes. *Mol Cell Endocrinol*. 2018;464:4-13.
5. Rang HP, Ritter JM, Flower JR, Henderson G. Lifestyle drugs and drugs in sport. In: Rang HP, Ritter JM, Flower JR, Henderson G. *Rang and Dale's Pharmacology*, 8th ed. Churchill Livingstone 2015:703-7.

6. Kanayama G, Hudson JI, Pope HG. Illicit anabolic-androgenic steroid use. *Horm Behav.* 2010;58(1):111–21.
7. Kanayama G, Pope HG. Illicit use of androgens and other hormones: recent advances. *Current Opinion in Endocrinology, Diabetes and Obesity.* 2012;19:211–9.
8. Frati P, Busardo FP, Cipolloni L, De Dominicis E, Fineschi V. Anabolic Androgenic Steroid (AAS) Related Deaths: Autoptic, Histopathological and Toxicological Findings. *Curr Neuroparmacol.* 2015;13(1):146–59.
9. Smit DL, de Ronde W. Outpatient clinic for users of anabolic androgenic steroids: an overview. *Neth J Med.* 2018;76(4):167.
10. Bennett PN, Brown MJ. Hypothalamic, pituitary and sex hormones. In: Bennett PN, Brown MJ. *Clinical Pharmacology*, 10th ed. Churchill Livingstone Elsevier. 2008;637–59.
11. Guyton AC, Hall JE. Reproductive and hormonal functions of a man. In: Guyton AC, Hall JE, editors. *Medical physiology*. Belgrade, Savremena Administracija. 2003;922–6.
12. Kicman AT. Pharmacology of anabolic steroids. *Br J Pharmacol.* 2008;154(3):502–21.
13. Rubinow DR, Schmidt PJ. Androgens, brain, and behavior. *Am J Psychiatry.* 1996;153(8):974–84.
14. Rang HP, Ritter JM, Flower JR, Henderson G. The Reproductive System. In: Rang HP, Ritter JM, Flower JR, Henderson G. *Rang and Dale's Pharmacology*, 8th ed. Churchill Livingstone. 2015:425–39.
15. Hill M, Dušková M, Stárka L. Dehydroepiandrosterone, its metabolites and ion channels. *J Steroid Biochem Mol Biol.* 2015;145:293–314.
16. Stárka L, Dušková M, Hill M. Dehydroepiandrosterone: a neuroactive steroid. *J Steroid Biochem Mol Biol.* 2015;145:254–60.
17. Snyder PJ. Androgens. In: Brunton L, Chabner B, Knollman B, editors. *Goodman and Gilman's The Pharmacological Basis of Therapeutics*. New York McGraw Hill. 2011: 1195–206.
18. Juhn M. Popular sports supplements and ergogenic aids. *Sports Med.* 2003; 33(12):921–39.
19. Smurawa TM, Congeni JA. Testosterone precursors: use and abuse in pediatric athletes. *Pediatr Clin North Am.* 2007;54(4):787–96.
20. Andriole GL, Bostwick DG, Brawley OW, Gomella LG, Marberger M, Montorsi F, et al. Effect of dutasteride on the risk of prostate cancer. *New Eng J Med.* 2010;362(13):1192–202.
21. Thompson IM, Goodman PJ, Tangen CM, Lucia MS, Miller GJ, Ford LG, et al. The influence of finasteride on the development of prostate cancer. *New Eng J Med.* 2003;349(3):215–24.
22. Frau R, Savoia P, Fanni S, Fiorentini C, Fidalgo C, Tronci E, et al. The 5-alpha reductase inhibitor finasteride reduces dyskinesia in a rat model of Parkinson's disease. *Exp Neurol.* 2017;291:1–7.
23. Duborjija-Kovacevic N, Jakovljevic V, Sabo A, Tomic Z. Anti-nociceptive and anti-inflammatory properties of 5alpha-reductase inhibitor finasteride in experimental animals. *Eur J Drug Metab Pharmacokinet.* 2008;33(3):181–6.
24. Mladenovic D, Hrcic D, Petronijevic N, Jevtic G, Radosavljevic T, Rasic-Markovic A, et al. Finasteride improves motor, EEG, and cellular changes in rat brain in thioacetamide-induced hepatic encephalopathy. *Am J Physiol Gastrointest Liver Physiol.* 2014;307(9):G931–40.
25. Paba S, Frau R, Godar SC, Devoto P, Marrosio F, Bortolato M. Steroid 5alpha-reductase as a novel therapeutic target for schizophrenia and other neuropsychiatric disorders. *Curr Pharm Des.* 2011;17(2):151–67.
26. Araujo AB, O'Donnell AB, Brambilla DJ, Simpson WB, Longcope C, Matsumoto AM, et al. Prevalence and incidence of androgen deficiency in middle-aged and older men: estimates from the Massachusetts Male Aging Study. *J Clin Endocrinol Metab.* 2004;89:5920–6.
27. Baillargeon J, Urban RJ, Ottenbacher KJ, Pierson KS, Goodwin JS. Trends in androgen prescribing in the United States, 2001 to 2011. *JAMA Intern Med.* 2013;173:1465–6.
28. Behre HM, Wang CC, Handelsman DJ, Nieschlag E. Pharmacology of testosterone preparations. In: Nieschlag E, Behre HM (eds). *Testosterone-Action, deficiency, substitution*, 3rd edn. Cambridge University Press. 2004;405–44.
29. Shoskes JJ, Wilson MK, Spinner ML. Pharmacology of testosterone replacement therapy preparations. *Transl Androl Urol.* 2016;5(6):834–43.
30. Testosterone Enantate 250mg/ml, solution for injection. Summary of Product Characteristics. Available in: <https://www.medicines.org.uk/emc/product/3733/smpc>, November 2018.
31. Nebido 1000mg/4ml, solution for injection. Summary of Product Characteristics. Available in: <https://www.medicines.org.uk/emc/product/3873/smpc>, November 2018.
32. Shahidi NT. A review of the chemistry, biological action, and clinical applications of anabolic-androgenic steroids. *Clin Ther.* 2001;23(9):1355–90.
33. Luisi M, Franchi E. Double-blind group comparative study of testosterone undecanoate and mesterolone in hypogonadal male patients. *J Endocrinol Invest.* 1980;3(3):305–8.
34. Sneader W. Hormone Analogues. In: Sneader W. *Drug Discovery: A History*. John Wiley & Sons 2005:188–226.
35. Handelsman DJ. *Androgen Physiology, Pharmacology and Abuse*. ENDOTEXT (online edition) 2016 Jan. Available in: <https://www.ncbi.nlm.nih.gov/books/NBK279000/>.
36. Li X, Takahashi M, Kushida K, Shimizu S, Hoshino H, Suzuki M, et al. The effects of nandrolone decanoate on bone mass and metabolism in ovariectomized rats with osteopenia. *J Bone Miner Metab.* 2000;18(5):258–63.
37. Guimarães APFGM, Butezloff MM, Zamarioli A, Issa JPM, Volpon JB. Nandrolone decanoate appears to increase bone callus formation in young adult rats after a complete femoral fracture. *Acta Cir Bras.* 2017;32(11):924–34.
38. Frisoli A Jr, Chaves PH, Pinheiro MM, Szejnfeld VL. The effect of nandrolone decanoate on bone mineral density, muscle mass, and hemoglobin levels in elderly women with osteoporosis: a double-blind, randomized, placebo-controlled clinical trial. *J Gerontol A Biol Sci Med Sci.* 2005;60(5):648–53.
39. Gennari C, Agnusdei D, Gonnelli S, Nardi P. Effects of nandrolone decanoate therapy on bone mass and calcium metabolism in women with established post-menopausal osteoporosis: a double-blind placebo-controlled study. *Maturitas.* 1989;11(3):187–97.
40. Compston J, Cooper A, Cooper C, Gittoes N, Gregson C, Harvey N, et al. UK clinical guideline for the prevention and treatment of osteoporosis. *Arch Osteoporosis.* 2017;12(1):43.
41. Amory JK, Watts NB, Easley KA, Sutton PR, Anawalt BD, Matsumoto AM, et al. Exogenous testosterone or testosterone with finasteride increases bone mineral density in older men with low serum testosterone. *J Clin Endocrinol Metab.* 2004;89(2):503–10.
42. Neto WK, Gama EF, Rocha LY, Ramos CC, Taets W, Scapini KB, et al. Effects of testosterone on lean mass gain in elderly men: systematic review with meta-analysis of controlled and randomized studies. *Age (Dordt).* 2015;37(1):9742.
43. Kenny AM, Prestwood KM, Gruman CA, Marcello KM, Raisz LG. Effects of transdermal testosterone on bone and muscle in older men with low bioavailable testosterone levels. *J Gerontol A Biol Sci Med Sci.* 2001;56(5):M266–72.
44. Singh P. Andropause: Current concepts. *Indian J Endocrinol Metab.* 2013;17(Suppl 3):S621–9.
45. Reis SL, Abdo CH. Benefits and risks of testosterone treatment for hypoactive sexual desire disorder in women: a critical review of studies published in the decades preceding and succeeding the advent of phosphodiesterase type 5 inhibitors. *Clinics (Sao Paulo).* 2014;69(4):294–303.
46. Basson R. Testosterone therapy for reduced libido in women. *Ther Adv Endocrinol Metab.* 2010;1(4):155–64.
47. European Medicines Agency (EMA). Testosterone-containing medicines. Available in: <https://www.ema.europa.eu/en/medicines/human/referrals/testosterone-containing-medicines>, November 2018
48. Gregersen N, Hilmand CB, Jensen PT, Giraldo AG. Sexual dysfunction in the menopause. Incidence, pharmacological treatment and side effects. *Ugeskr Laeger.* 2006;168(6):559–63.
49. Rabkin JG, Wagner GJ, Rabkin R. A double-blind, placebo-controlled trial of testosterone therapy for HIV-positive men with hypogonadal symptoms. *Arch Gen Psychiatry.* 2000;57(2):141–7.
50. Kraft D. Long-term treatment of renal anaemia with mesterolone (author's transl). *Dtsch Med Wochenschr.* 1980;105(23):830–2.
51. Gascón A, Belvis JJ, Berisa F, Iglesias E, Estopiñán V, Teruel JL. Nandrolone decanoate is a good alternative for the treatment of anemia in elderly male patients on hemodialysis. *Geriatr Nephrol Urol.* 1999;9(2):67–72.
52. Schifeling DJ, Jackson DV, Zekan PJ, Muss HB. Fluoxymesterone as third line endocrine therapy for advanced breast cancer. A phase II trial of the Piedmont Oncology Association. *Am J Clin Oncol.* 1992;15(3):233–5.
53. Ingle JN, Twito DI, Schaid DJ, Cullinan SA, Krook JE, Mailliard JA, et al. Combination hormonal therapy with tamoxifen plus fluoxymesterone versus tamoxifen alone in postmenopausal women with metastatic breast cancer. An updated analysis. *Cancer.* 1991;67(4):886–91
54. Biberoglu KO, Behrman SJ. Dosage aspects of danazol therapy in endometriosis: short-term and long-term effectiveness. *Am J Obstet Gynecol.* 1981;139(6):645–54.
55. Buttram VC Jr, Reiter RC, Ward S. Treatment of endometriosis with danazol: report of a 6-year prospective study. *Fertil Steril.* 1985;43(3):353–60.
56. Diaz EC, Herndon DN, Porter C, Sidossis LS, Suman OE, Børsheim E. Effects of pharmacological interventions on muscle protein synthesis and breakdown in recovery from burns. *Burns.* 2015;41(4):649–57.
57. Murphy KD, Thomas S, Mlcak RP, Chinkes DL, Klein GL, Herndon DN. Effects of long-term oxandrolone administration in severely burned children. *Surgery.* 2004;136(2):219–24.

58. Handelsman DJ. Androgen therapy in non-gonadal disease. In: Nieschlag E and Behre HM, Editors. *Testosterone: Action, Deficiency and Substitution*. Cambridge University Press 2011;372-407.
59. Woerdeman J, de Ronde W. Therapeutic effects of anabolic androgenic steroids on chronic diseases associated with muscle wasting. Review. *Expert Opin Investig Drugs*. 2011;20(1):87-97.
60. Fitzpatrick F. *Where steroids were all the rage: A doctor's curiosity and a businessman's love of weightlifting set off a revolution in York*. Philadelphia Inquirer; Philadelphia, PA. 2002.
61. Rahman SZ, Gupta V, Sukhlecha A, Khunte Y. Lifestyle drugs: concept and impact on society. *Indian J Pharm Sci*. 2010;72(4):409-13.
62. Gilbert D, Walley T, New B. Lifestyle medicine. *Br Med J*. 2000;321:1341-4.
63. Flower RJ. Lifestyle and non-medical uses of drugs. In: Rang HP, Dale MM, Ritter JM, editors. *Rang and Dale's Pharmacology*. 6th ed. Churchill Livingstone 2007:765-9.
64. Van Thuyne W, Van Eenoo P, Delbeke FT. Nutritional supplements: prevalence of use and contamination with doping agents. *Nutr Res Rev*. 2006;19:147-58.
65. Kazlauskas R. Designer steroids. *Handb Exp Pharmacol*. 2010;195:155-85.
66. Malvey TC, Armsey TD 2nd. Tetrahydrogestrinone: the discovery of a designer steroid. *Curr Sports Med Rep*. 2005;4(4):227-30.
67. Joseph JF, Parr MK. Synthetic androgens as designer supplements. *Curr Neuropharmacol*. 2015;13(1):89-100.
68. Pope HG, Kanayama G, Ionescu-Ploggia M, Hudson JI. Anabolic steroid users' attitudes towards physicians. *Addiction*. 2004;99(9):1189-94.
69. Gupta L, Towler B. General practitioners' views and knowledge about anabolic steroid use - survey of GPs in a high prevalence area. *Drug Alcohol Rev*. 1997;16(4):373-9.
70. Geyer H, Schanzer W, Thevis M. Anabolic agents: recent strategies for their detection and protection from inadvertent doping. *Br J Sports Med*. 2014;48(10):820-6.
71. WADA. 2015 Anti-Doping Testing Figures - Laboratory Report. World Anti-Doping Agency, 2016. (<https://www.wada-ama.org>).
72. Reyes-Fuentes A, Veldhuis JD. Neuroendocrine physiology of the normal male gonadal axis. *Endocrinol Metab Clin North Am*. 1993;22:93-124.
73. Turillazzi E, Perilli G, Di Paolo M, Neri M, Riezzo I, Fineschi V. Side effects of AAS abuse: an overview. *Mini Rev Med Chem*. 2011;11(5):374-89.
74. Kindermann W. Cardiovascular side effects of anabolic-androgenic steroids. *Herz*. 2006;31(6):566-73.
75. Higgins JP, Heshmat A, Higgins CL. Androgen abuse and increased cardiac risk. *South Med J*. 2012;105(12):670-4.
76. Melchert RB, Welder AA. Cardiovascular effects of androgenic-anabolic steroids. *Med Sci Sports Exerc*. 1995;27(9):1252-62.
77. Schwartz PJ, La Rovere MT, Vanoli E. Autonomic nervous system and sudden cardiac death. Experimental basis and clinical observations for post-myocardial infarction risk stratification. *Circulation*. 1992;85(1)Suppl.:177-191.
78. Christou MA, Christou PA, Markozannes G, Tsatsoulis A, Mastorakos G, Tigas S. Effects of Anabolic Androgenic Steroids on the Reproductive System of Athletes and Recreational Users: A Systematic Review and Meta-Analysis. *Sports Med*. 2017;47(9):1869-83.
79. Westlye LT, Kaufmann T, Alnaes D, Hullstein IR, Bjornebekk A. Brain connectivity aberrations in anabolic-androgenic steroid users. *Neuroimage Clin*. 2016;13:62-9.
80. Petersson A, Garle M, Holmgren P, Druid H, Krantz P, Thiblin I. Toxicological findings and manner of death in autopsied users of anabolic androgenic steroids. *Drug Alcohol Depend*. 2006;81(3):241-9.
81. Thiblin I, Lindquist O, Rais J. Cause and manner of death among users of anabolic androgenic steroids. *J Forensic Sci*. 2000;45(1):16-23.
82. Kanayama G, Pope HG Jr, Hudson JI. "Body-image" drugs: a growing psychosomatic problem. *Psychother Psychosom*. 2001b;70:61-5.
83. Huang EY, Chen YH, Huang TY, Chen YJ, Chow LH. Chronic administration of nandrolone increases susceptibility to morphine dependence without correlation with LWV-hemorphin 7 in rats. *Neuropeptides*. 2016;59:63-9.
84. Arvary D, Pope HG Jr. Anabolic-androgenic steroids as a gateway to opioid dependence. *New Engl J Med*. 2000;342(20):1532.
85. Graham MR, Davies B, Grace FM, Kicman A, Baker JS. Anabolic steroid use: patterns of use and detection of doping. *Sports Med*. 2008;38(6):505-25.
86. Solimini R, Rotolo MC, Mastrobattista L, Mortali C, Minutillo A, Pichini S, et al. Hepatotoxicity associated with illicit use of anabolic androgenic steroids in doping. *Eur Rev Med Pharmacol Sci*. 2017;21(1 Suppl):7-16.
87. Sánchez-Osorio M, Duarte-Rojo A, Martínez-Benítez B, Torre A, Uribe M. Anabolic-androgenic steroids and liver injury. *Liver Int*. 2008;28(2):278-82.
88. Bond P, Liewellyn W, Van Moi P. Anabolic androgenic steroid-induced hepatotoxicity. *Med Hypotheses*. 2016;93:150-3.

Sport classification regulations for athletes with differences in sexual development (DSD)

Pedro Manonelles Marqueta¹, José Luis Terreros Blanco², Juan José Rodríguez Sendín³

¹President of the Spanish Society of Sports Medicine (SEMED-FEMEDE). Extraordinary Professor of Sports Medicine. Catholic University of Murcia (UCAM). ²Director of the Spanish Agency for the Protection of Health in Sport (AEPHAD). ³President of the Central Committee of Deontology, Medical Law and Oversight of the Spanish Collegiate Medical Organization (OMC).

Received: 11.02.2019

Accepted: 20.03.2019

Summary

The classification systems for sports competition are based primarily on the sex of the athlete and generate the male and female categories in almost all existing sports.

There have been some cases of fraud, in which men have competed in the female category, and others, in which some female competitors have caused suspicions about their sex. The last known case is the South African athlete Caster Semenya, who won the final of the 800 m in the World Championships in Berlin in 2009 with 2.45 seconds less than the second classified, with a distance of 16 m in the final straight.

After a multitude of studies, it was verified that the athlete presented a medical condition called difference of the sexual development (DSD), with a production of high levels of testosterone.

High testosterone levels, with sensitivity to this hormone in women, may represent a huge advantage in sports performance, which has been quantified by a range over 9%. The International Athletics Federation (IAAF) has promulgated a rule requiring female athletes with high levels of testosterone and sensitive to it, who want to participate in 400m to a mile tests, to decrease testosterone levels by using estrogens. This work analyses sports classification systems, the physiological effects of testosterone, the basis of sexual differentiation, and presents the medical and deontological arguments to refute the obligation of hormonal treatment of women to be able to compete in sports.

Key words:

Sports classification.
Differentiation of sexual state.
DSD. Testosterone. Performance.
Athletics. Sports regulations.

Reglamento de clasificación deportiva para atletas con diferencias en el desarrollo sexual (DSD)

Resumen

Los sistemas de clasificación para competición deportiva se basan fundamentalmente en el sexo del deportista y generan las categorías masculina y femenina, en la práctica totalidad de deportes.

Ha habido algunos casos de fraude, en los que hombres han competido en la categoría femenina, y otros, en los que algunas competidoras femeninas han suscitado sospechas sobre su sexo. El último caso conocido es el de la atleta sudafricana Caster Semenya, ganadora de la final de los 800 m en el Campeonato del Mundo de atletismo de Berlín de 2009 con 2:45 segundos menos que la segunda clasificada, a la que superó en 16 m en la recta final.

Tras multitud de estudios, se comprobó que la atleta presentaba una condición médica denominada diferencia del desarrollo sexual (DSD), con producción de elevados niveles de testosterona.

Las cifras elevadas de testosterona, con sensibilidad a esta hormona en mujeres, pueden suponer una enorme ventaja en el rendimiento deportivo, que se ha cuantificado en un rango sobre el 9%.

La Federación Internacional de Atletismo (IAAF) ha promulgado una normativa que obliga a las atletas femeninas con altos niveles de testosterona y sensibilidad a la misma, que quieren participar en pruebas de 400 m a la milla, a disminuir las cifras de testosterona mediante la utilización de estrógenos.

Este trabajo analiza los sistemas de clasificación deportiva, los efectos fisiológicos de la testosterona, las bases de la diferenciación sexual, y presenta los argumentos médicos y deontológicos para rebatir la obligación de tratamiento hormonal de mujeres para poder competir en especialidades deportivas.

Palabras clave:

Clasificación deportiva. Diferenciación del estado sexual. DSD. Testosterona. Rendimiento. Atletismo. Normas deportivas.

Correspondence: Pedro Manonelles
E-mail: manonelles@telefonica.net

Introduction

On 23 April, 2018, the International Association of Athletics Federations (IAAF) published the following: ELIGIBILITY REGULATIONS FOR FEMALE CLASSIFICATION (ATHLETES WITH DIFFERENCES OF SEX DEVELOPMENT)¹ that was scheduled to come into force on 1 November, 2018, but this was suspended on the basis of the appeal made by the female athlete, Caster Semenya, before the Court of Arbitration for Sport (CAS) and conditional upon its final ruling.

These regulations were drawn up as a consequence of the debate which arose due to the sports results of women athletes who had achieved very significant advantages over their opponents. All this was particularly significant in the case of the South African athlete mentioned above. This female athlete won the women's 800 m trial in the 2009 World Championship in Berlin with a time of 1:55.45 min, while the athlete coming in second took 1: 57.90: a difference of 2.45 seconds i.e. a distance of more than 16 meters (Figure 1). In this 800 m trial Semenya won the final of the Olympic Games in Rio de Janeiro 2016 and was World Champion in Berlin in 2009 and in London in 2017. Her best time was 1:54.25 and she also won international competitions between 400 m and 1500 m. However, it should be borne in mind that she was also defeated on the track in the 800 m women's race in the London Olympics in 2012 and the 2011 Daegu World Championship, both times by an athlete who was subsequently stripped of both gold medals for having used doping and who was suspended for that reason for 4 years. Both medals were then also awarded to Caster Semenya.

The case of Semenya has been giving rise to speculations in the media, like she had used doping substances, or that she was in fact a man.

As a result of all this, Semenya has undergone a number of different medical and genetic studies, some of whose results seem unfortunately to have been leaked to the media. In the case of other studies, numerous speculations have been reported. What seems clear is that the athlete suffers from androgenism and that she has a blood testosterone level higher than the reference values for women of her characteristics.

Before moving forward, it must be considered that other elite athletes have also been subjected to intense media exposure for the same reasons: their physical appearance and exceptional sports performance.

Figure 1. Arrival of the women's 800m final of the World Athletics Championships, Berlin 2009.



The perception that several of them are competing or have competed at the highest level in the 800 m trial, like the holder herself of the world record for that distance (1: 53.28 since 1983) seems very strange.

Back in 2010, Semenya underwent a thorough examination by a group of specialists who determined that she should compete as a woman and without any limitation. However, her exceptional performance and its media impact led the IAAF to look for a method to avoid such major differences between competitors, to the point of producing the regulations of 2018, already cited¹.

As a result of all the above, this report aims to analyse the regulations that the IAAF intends to impose in order to reduce the differences in performance in women's athletics, from the points of view of Sports Ethics and especially of Medicine.

Physiological aspects

It is undeniably clear that men have significant advantages in muscle size, strength and power, compared with women. Table 1, produced by the authors, shows the percentage differences between men and women, in some of the world records in athletics. To this end, only trials where women and men competed under equal conditions (ruling out obstacle races, which have different height, and throwing with different weights) have been considered. In order to compare these differences in jumps and races, the records of female athletes have been expressed by transforming the record time into the race speed (metres per second⁻¹).

Generally, this difference is quantified as a 10-12% advantage for men¹. This is attributed to man's higher testosterone levels from puberty, among other possible causes. The cellular and molecular mechanisms of the advantage that a higher rate of testosterone provides are clearly described².

Table 1. Analysis of the percentage differences between men and women for some world records in athletics.

Trial	World record male	Speed (msec ⁻¹) or distance (m)	World record female	Speed (mseg ⁻¹) or distance (m)	Difference (%)
100 m	9,58	10,44	10,49	9,53	8,7
200 m	19,19	10,42	21,34	9,37	10,1
400 m	43,03	9,30	47,60	8,40	9,6
800 m	1:40,91	7,93	1:53,28	7,08	10,7
1500 m	3:26:00	7,28	3:50,07	6,52	10,5
5000 m	12:37,35	6,60	14:11,15	5,87	11,0
10000 mm	26:17,53	6,34	29:17,46	5,69	10,2
Marathon	2:01,39	5,75	2:15,25	5,17	10,2
20km walk	1:16,36	4,35	1:24,38	3,94	9,5
Relay 4x100	36,84	10,86	40,82	9,80	9,8
Relay 4x400	2:54,29	9,17	3:15,17	8,19	10,7
Long jump	8,95	8,95	7,52	7,52	16,0
Triple jump	18,29	18,29	15,50	15,50	15,3
High jump	2,45	2,45	2,09	2,09	14,7
Pole vault	6,16	6,16	5,06	5,06	17,9

In Table 1 it can be seen how, in race trials, the advantage of male athletes would be between 8.7% and 11.0%, however specifically in the distances between 400 m and one mile (including the 4x400m relay) they do not seem to have a wider difference (between 9.6 and 10.7%) than the rest of the distances. In addition, it can be seen how, in the case of jumps, where rapid use of muscle strength is of major importance, the differences are greater (advantages for male athletes are between 14.7% and 17.9%). Although this cannot be compared, one could venture that it is very possible that in throwing that difference would be even greater.

Gender and sport. Classification and verification systems

Sport-based classification, separated by gender, in practically all sports and sports specialties, is something totally accepted today. The almost insurmountable differences in performance between women and men is the clear reason for this separation.

To maintain this separation and avoid the pitfalls in gender classification, in 1950 the IAAF set down a regulation including physical medical examinations, regulations that, in the long run, were extended to other sports. Back in 1950, an athlete was prevented from continuing her sports career due to refusing to undergo one of these medical examinations. The standards developed further and in 1966 Barr's chromatin began to be studied in female athletes who wanted to participate in female trials, using the karyotype in a saliva sample as an initial test. If in this test some Y chromosome was detected, medical examinations were carried out, including the morphological verification of external genitalia.

Although these physical explorations ended up being considered degrading and were happily repealed in all sports. In the early 1990s the use of the Polymerase Chain Reaction (PCR) technique was advocated for the extension and examination of DNA strands, as a method which was more objective and scientific.

Finally, in 2011 the IAAF set down two different regulations, one for people with sex reassignment³, and another for female athletes with hyperandrogenism⁴. However, in 2015, a female Indian athlete brought an appeal to the CAS regarding her suspension due to a diagnosis of hyperandrogenism, after a process described by her, as "horrible and humiliating". The athlete obtained a favourable ruling, since the CAS finally accepted that there was not enough evidence of the advantage that a higher level of testosterone could produce in female athletes⁵. Thus, the IAAF regulation on hyperandrogenism was repealed.

As can be seen, this is a recurrent issue, which appears to be harming a number of women athletes, until institutions are persuaded that the matter should be left off the agenda. Now, in 2018 and 2019, we are once again in the worst possible position as regards the rights of certain athletes.

Sexual differentiation

Biological gender is a general term that includes different aspects of chromosomal, gonadal, hormonal and phenotypic sex, each of which is fixed in an individual. Generally, all these classifications of gender are aligned in a conventional binary: male and female. However, it must

be considered that there are congenital conditions that cause misalignment and atypical development of the chromosomal, gonadal and / or anatomical genders. As a medical description of these conditions, terms such as "disorders or anomalies of sexual development"^{6,7} and later "intersexual states" or "intersex" began to be used. The term "anomalies of sexual differentiation" was introduced in 2006⁶, however all of these have been widely contested in the scientific literature⁶⁻¹⁰.

Currently they are classified as "differences in sexual development" or DSD, although later the term "variations in sexual development" (VSD) was proposed¹⁰.

As will be seen later, DSD may involve ambiguity of external genitalia and various combinations of chromosomal genotypes and sexual phenotypes other than XX-woman and XY-man^{7,11}. In the administrative field, there are a number of national systems that recognize legal genders which are different from "man" and "woman", such as "intersex", "X" or "other".

Different institutions with powers in the field of the protection of human rights attempt to avoid harmful practices and discrimination against people with these conditions. In 2015, the United Nations High Commissioner launched a campaign for the free equality of rights of people classified as "intersex"¹².

Equally, in 2015 the Council of Europe made a statement asking member states for a non-binary gender classification and to seriously consider the implications of a new and better classification of "intersex" individuals¹³. In 2016, numerous Committees and U.N. subcommittees (Against Torture, of the Rights of the Child, of the Rights of Persons with Disabilities), together with the Council of Europe and the Human Rights Commissions of Africa and the Americas, launched a document calling for an end to the violent and harmful medical practices on "intersex" children and adults¹⁴.

We should not forget that disclosing information about the medical history of any patient can lead to very serious consequences¹⁵. Furthermore, in these sensitive cases public knowledge of this protected data will have irreparable consequences for the normal physical development of young people and the psychosocial sphere of all those affected^{16,17}, as established by the World Health Organization¹⁸.

Regarding the embryological aspect of foetal sexual differentiation, it should be observed that this encompasses a series of processes whose determination and regulation involve numerous genes, proteins and hormones. Starting from a first stage of gonadal and genital development (6 weeks post-fertilization), which is common to both sexes, it is in the period of differentiation when these conditions may occur and have been given the controversial classifications previously identified. These comprise a broad spectrum of discrepancies between chromosomal, gonadal and phenotypic (genital) criteria that define sexual differentiation and are now considered "Differences in Sexual Development (DSD)". Their classification can be seen in Table 2¹⁹.

As regards the involvement of testosterone, it should be remembered that the majority of women (including elite athletes) have circulating blood testosterone levels (0.12-1.79 nmol.L⁻¹) which are lower than those of post-pubertal males ((7.7-29.4 nmol.L⁻¹). It is accepted that, in the absence of DSD or tumour, no woman should have serum testosterone levels equal to or greater than 5 nmol.L⁻¹(²⁰).

Table 2. Classification of Differences in Sexual Development (DSD)¹⁹.


Chromosome changes	47,XXY: Klinefelter syndrome and variants 45,X0 and mosaics 45,X0 / 46, XX: Turner syndrome and variants 45,X0 / 46,XY: Mixed gonadal dysgenesis 46,XX / 46,XY: ovotesticular ADS/DSD, 47,XYY chimerism
Karyotype 46,XY Anomalies in gonadal development	Gonadal dysgenesis 46,XY (complete or partial) (SRY, SOX9, NR5A1, WT1, DHH, etc.) ovotesticular 46, XY Testicular regression syndrome (includes anorchy and testicular leakage syndrome)
Anomalies in genital development due to changes in hormonal synthesis or action	Changes in androgen synthesis Mutations in the LH receptor (plasia or aplasia of Leydig cells; LHCGR) Smith-Lemli-Opitz syndrome (deficit in 7-dehydrocholesterol reductase: DHCR7) Defects in the synthesis of testosterone Congenital lipoid suprarenal hyperplasia (StAR) Deficiency in cholesterol desmolase (CYP11A1) Deficiency in 3_-hydroxysteroid dehydrogenase (HSD3B2) Deficiency in 17_-hydroxylase / 17-20 lyase (CYP17A1) Deficiency P450 oxidoreductase (POR) Deficiency in cytochrome b5 (CYB5) Deficiency in 17_-hydroxysteroid dehydrogenase (HSD17B3) Deficiency in 5_-reductase type 2 (SRD5A2) Changes in the action of androgens Insensitivity to androgens (AR; total or partial = CAIS or PAIS) Drugs and environmental modulators Changes in the synthesis or action of the antimullerian hormone Persistent Mullerian duct syndrome (AMH / AMHR2)
Others	Malformation syndromes with changes in male genital development (e.g.: cloacal anomalies, Aarskog syndrome, Robinow syndrome, etc.) Severe early-onset intrauterine growth restriction Isolated hypospadias (CXorf6 or MAMLD1) Congenital hypogonadotropic hypogonadism Cryptorchidism (INSL3, RXFP2 [o INSL3R or GREAT])
Karyotype 46,XX Anomalies in gonadal development	Gonadal dysgenesis 46,XX Ovotesticular 46,XX Testicular 46,XX ADS/DSD (SRY, dup SOX9, RSPO1) or male 46,XX Foetal production Deficiency in 21-hydroxylase (CPY21A2) Deficiency in 11_-hydroxylase (CYP11B1) P450 oxidoreductase deficiency (POR) Deficiency in cytochrome b5 (CYB5) Deficiency in 3_-hydroxysteroid dehydrogenase (HSD3B2) Mutations of the glucocorticoid receptor (NR3C1) Fetoplacental production Deficiency in placental and foetal aromatase (CYP19A1) P450 oxidoreductase deficiency (POR) Foetal or placental tumours that produce androgens Maternal production Androgenic drugs Maternal virilising tumours (e.g. luteomas, Krukenberg tumour)
Others	Malformation syndromes Hypoplasia/agenesis of Mullerian structures (Rokitansky-Hauser syndrome type I and type II - MURCS) Uterine anomalies (e.g. MODY 5) Vaginal atresia Adhesions of vaginal labia

People with DSD may have very high levels of natural testosterone that may be similar to, or higher than, normal male values²⁰. There is a broad medical and scientific consensus that, if these people are sensitive to androgens, i.e. if they have normo-functioning androgen receptors, and record such high levels of natural testosterone, their muscle mass, strength and haemoglobin level may also be much higher and, therefore, significantly improve their sports potential²¹⁻³⁵. It can be considered

that an increase of blood testosterone rate in women between 0.9 and 7.3 nmol.L⁻¹ produces an increase in muscle mass of 4% and in muscle strength of 12-26%. The increase in testosterone to levels of: 5, 7, 10 and 19 nmol.L⁻¹, increases the amount of haemoglobin by 6.5%, 7.8%, 8.9% and 11%, respectively.

It is therefore estimated that the ergogenic advantage of testosterone levels in the male range instead of the female range is over 9%²¹.

Figure 2. Best annual times in the 800 m for an elite female athlete with DSD between 2008 and 2016, with control of her testosterone rate between 2010 and 2014²¹.



All this indicates that, for the most part, the competitive advantages of men are attributable to the action of male sex hormones, in such a way that the increase in testosterone in women from the range of female values to the male range would cause this increase in muscle mass, strength and the rate of haemoglobin.

The ergogenic power of testosterone and that of all its derivatives and other substances with similar chemical structure or biological effects, is manifested by its inclusion in the World Anti-Doping Agency's (WADA) List of Prohibited Substances and Methods³⁶. These substances are prohibited at all times (in and out of competition) and in all sports, as Non-Specific Substances in Category S1: Anabolic Agents. These are included in that list since it has existed, and their use is considered to be a case of utmost seriousness.

However, from the legal point of view, the CAS ruling of 2015 should be recalled⁵, stating that there is not enough evidence of the advantage that a higher rate of testosterone may give to female athletes.

For its part, the IAAF has compiled observational data on the effects of artificial suppression of high testosterone levels on athletes with DSD, depending on whether or not their testosterone levels are suppressed. The suppression of circulating testosterone levels in the case of three female athletes with DSD from between 21 and 25 nmol.L⁻¹ down to a rate below 2 nmol.L⁻¹ led to a decrease in their performance on average of 5.7%, as can be observed in Figure 2^{21,37}.

This study shows that athletes with DSD with circulating testosterone levels in the normal male range have a significant competitive advantage over athletes with testosterone levels in the normal female range, and for the IAAF this would justify forcing these athletes with DSD to reduce testosterone levels to the normal female range to continue competing in the female category.

Application of the IAAF regulations¹

Sphere

The IAAF data would indicate that the advantages of some athletes with DSD have had a significant effect on mid-distance track events, so

it is inferred from this that the rule only applies to events from 400 m to a mile (which the regulations term "Restricted Events"). In international competition and outdoor track events, these events are: 400 m, 400 m hurdles, 800 m, 1500 m and one mile, alone or as part of a relay race or a combined trial.

The special eligibility requirements described apply only to "Relevant Athletes" in the female classification in a Restricted Event in International Competition.

The regulations describe as "Relevant Athlete" those who meet the following three criteria:

- Women with one of the following DSDs:
 - Deficiency of 5 α -reductase type 2.
 - Partial androgen insensitivity syndrome (PAIS), or Deficiency of 17 β -hydroxysteroid dehydrogenase type 3) or Congenital adrenal hyperplasia or Deficiency of 3 β -hydroxysteroid dehydrogenase.
- Ovotesticular DSD, or Another genetic disorder with impaired gonadal steroidogenesis.
- Has circulating blood testosterone levels of ≥ 5 nmol.L⁻¹.
- Has sufficient sensitivity to androgens so that these testosterone levels have an androgenizing effect.

Conditions for eligibility

- Being recognized by law as a woman or as intersexual (or equivalent).
- Reducing the rate of circulating blood testosterone to below 5nmol.L⁻¹ for a continuous period of at least 6 months, using, for example, contraceptives.
- Maintaining the circulating blood testosterone level below 5 nmol.L⁻¹ continuously (in or outside competition) for as long as wished to maintain eligibility to compete in the female classification in Restricted Events in international competition (or setting a world record in a Restricted Event in a competition that is not an international contest).

Taking part without suppressing high levels of testosterone

The regulations allow women athletes who do not want to follow the above guidelines to participate in the following circumstances:

- In the female classification:
 - In any competition that is not an international competition.
 - In international competitions: in any discipline other than track events of between 400 m and one mile.
- In the men's classification: In any competition at any level, in any discipline, without restriction.
- In any 'intersex' classification (or similar) that the organizer of the event can offer in any competition at any level, in any discipline, without restriction.

Reasons given by the IAAF for implementing this rule

The arguments expressed by the IAAF in the text¹, are as follows:

- To guarantee fair and valid competition in athletics within categories that create a level playing field and to guarantee that success

is determined by talent, dedication, hard work, other values and characteristics that sport embodies and celebrates.

- To benefit a broad class of female athletes.
- To encourage athletes to achieve the great commitment and sacrifice required to excel in sport, and thus inspire new generations to take up sport and to aspire to the same excellence. It does not wish to risk discouraging those aspirations by having conditions of unfair competition, which would deny female athletes a fair chance of succeeding.
- Because of the significant advantages in size, strength and power that men generally possess over women from puberty, largely due to much higher levels of circulating testosterone in males, and the impact these advantages may have on sport performance, it is generally accepted that competition between male and female athletes would not be fair and would risk discouraging women from participating in sport. Therefore, in addition to the separate competition categories based on age, the IAAF has also created separate competition categories for male and female athletes.

Sports ethics arguments against the IAAF regulations

We must begin with the very concept of competitive sport as a human activity based on inequality between people. Competition attempts to measure precisely that inequality that is what gives rise to the victory of some over others. Setting down rules that try to eliminate or reduce this inequality would transform athletic competition into something else.

Sports performance is an extremely complex phenotypic trait that in turn is influenced, although not determined, by many other traits, such as the distribution of muscle fibre type, aerobic power and capacity, strength and anaerobic capacity, and the ability to train physical skills³⁸. Although the extrinsic determinants of human athletic performance, such as training, nutrition, living conditions, etc., have an undoubted and significant impact on sports performance, the importance of extrinsic determinants must not be trivialized, among these the genetic ones. It is clear that it is impossible to establish a single formula for anyone to become a sports champion and that quantifying the contribution of each of these determinants continues to be a challenge to research in sports science³⁹⁻⁴².

However, certain people who have not undergone scheduled training is already able to demonstrate extraordinarily high levels of physical performance. Some people also show a better response to training and improve their performance much more than others. when following the same working programme^{40,41}.

The fact that genetics is a very significant intrinsic factor in athletic performance is shown by the analysis of athlete rankings (year 2018 on 17 December) where it can be seen that, in the 1500 m, 14 of the 15 best all-time records were set by athletes born on the African continent, and that the first 7 athletes in this trial in 2018 were born in Ethiopia, while only one thrower with one of the top 60 world times (15 in each of the weight-lifting, discus, hammer and javelin trials) was born in Africa, and only 4 in total (7%) had a phenotype for that continent.

The IAAF intends to delimit a group of Restricted Events (400 m to one mile) to apply this regulation, without stating any objective reason why this is done in these trials and not in other events. It should be borne in mind that, as has already been explained, there is no evidence that high testosterone levels in women may be more advantageous in these trials than in others, and it has already been seen that in the jump disciplines this advantage may be greater. In this way the IAAF would be legislating in violation of the principle of generality by derogating a la carte and committing injustice towards a very specific population, almost singling out specific female athletes.

In addition, it seems essential to consider that the artificial reduction of the advantage that nature has granted to certain women in the form of higher blood testosterone levels, could be somewhat random, since there is no evidence of a direct relationship between the reduction in the rate of testosterone in female athletes, hyperandrogenism and the impact on sports performance. Consequently, with the same decrease in blood testosterone, even taking this at the same level, this could affect the running speed to a different extent in one female athlete from another.

And what happens in men who have circulating testosterone levels higher than the reference values? In the 80s, the genetic basis of Polycystic Ovarian Syndrome began to be studied⁴³, a basis that has become much better known over the last decades⁴⁴⁻⁴⁵. Today, numerous studies are advancing in the description of a genetic load similar to the cases of female hyperandrogenism with Polycystic Ovarian Syndrome and various types of male hyperandrogenism⁴⁶⁻⁵⁰. Why should not men who could obtain good results in those Restricted events or any other trial have applied to them regulations similar to those that the IAAF wants to apply to women?

We cannot conclude without taking into account the efforts that the governments of the world and the sports movement have been making to eradicate the scourge of doping. As the World Anti-Doping Code says in justifying this effort in its Fundamentals: *“Anti-doping programs seek to preserve what is intrinsically valuable about sport. This intrinsic value is often referred to as “the spirit of sport.” it is the essence of Olympism, the pursuit of human excellence through the dedicated perfection of each person’s natural talents. It is how we play true. The spirit of sport is the celebration of the human spirit, body and mind, and is reflected in values we find in and through sport, ...”*

As can be seen, natural talent is the excellence that is sought to be preserved, so how can this fight be justified, if when these natural talents appear, the attempt is to eliminate them in an artificial way? That way we cannot defend fair play⁵¹.

Medical ethics arguments against the IAAF regulations

If sports ethics arguments against these regulations seem overwhelming evidence, medical arguments that are very clear and concise seem much more important - and any medical professional will undoubtedly understand these.

The rules of Medicine prohibit the use of medication that is not destined to treat a disease or pathology. Moreover, medications must be used following the established indications and not for any other purposes. The use of medication outside these circumstances is con-

trary to medical practice and is therefore an offence on the part of the prescribing physician⁵².

These regulations, which aim to reduce performance in people who have innate qualities of genetic origin and have not been obtained by illegal means, would undoubtedly apply in these cases, and doctors who prescribe treatments for this purpose could clearly be committing an offence.

But above all, it should be noted that in the mind of every doctor, in their relationship with their patients, the side effects, contraindications and risks that prescribed medication could have should take pride of place. It should not be forgotten that the drugs that affect the hormonal sphere of individuals involve significant health risks, risks that are accentuated exponentially if these drugs are used beyond their medical indications⁵³.

Finally, it is necessary to add the problem that arises in cases like these with respect to the protection of data and even individuals' rights. We are seeing women who, as a result of wanting to exercise their right to practice sports, find themselves involved in the maximum possible circulation internationally of their tests results, examinations and medical diagnoses, to then be exhibited in the media at the same level as cheating athletes who use doping to excel in sports.

Conclusion

Faced with these regulations that clearly fly in the face of unquestionable evidence from the point of view of sportsmanship, against the oath of medical practice, and even against the rights of individuals, we can only demand their immediate repeal.

Conflict of interest

The authors do not declare a conflict of interest.

Bibliography


- IAAF. Eligibility regulations for the female classification (athletes with Differences of Sex Development). 2018. (Consultado 7/1/2019). Disponible en: <https://www.documentcloud.org/documents/4449932-IAAF-Eligibility-Regulations-for-the-Female.html>
- Kadi A. Cellular and molecular mechanisms responsible for the action of testosterone on human skeletal muscle. A basis for illegal performance enhancement. *Br J Pharmacol* 2008;154:522-8.
- IAAF Regulations governing eligibility of athletes who have undergone sex reassignment to compete in women's competition. 2011. (Consultado 7/1/2019). Disponible en: <https://www.iaaf.org/responsive/download/downloadregistration?token=vzim4unobddtpci2exbluh0mg9u87fpubur6dl>
- IAAF Regulations governing eligibility of females with hyperandrogenism to compete in women's competition. 2011. (Consultado 7/6/2018). Disponible en: <https://www.iaaf.org/about-iaaf/documents/health-science>
- Court of Arbitration for Sport. CAS2014/A/3759. Dutee Chand v. Athletics Federation of India (AFI) & The International Association of Athletics Federations (IAAF). 2015. (Consultado 7/1/2019). Disponible en: <https://playwired.files.wordpress.com/2015/09/dutee-chand-v-athleticsfederation-of-india-afi-the-international-association-of-athletics-federationsiaaf.pdf>.
- Holmes M. The intersex encliridion: Naming and knowledge. *Somatechnics* 2014;1:388-411.
- Money J, Ehrhardt AA. Man & woman boy & girl. Differentiation and dimorphism of gender identity from conception to maturity. The John Hopkins University Press. USA 1972.
- Davis G. Contesting Intersex: The Dubious Diagnosis. New York University Press. USA 2015.
- Houk CP, Hughes IA, Ahmed SF, Lee PA. Summary of Consensus Statement on Intersex disorders and their management. Writing Committee for the International Intersex Consensus. *Pediatrics* 2006;118: 753-7.
- Diamond M, Beh HG. Variations of sex development instead of disorders of sex development. *Arch Dis Child*. 2006 Electronic Letter, 27 July 2006. (Consultado 17/1/2019). Disponible en: <http://www.hawaii.edu/PCSS/biblio/articles/2005to2009/2006-variations.html>.
- Domurat Dreger A. Hermaphrodites and the medical invention of sex. Harvard University Press. USA 2001.
- United Nations Office of the High Commissioner for Human Rights. "Free & Equal Campaign Fact Sheet: Intersex". Fact sheet 2015. (Consultado 7/1/2019). Disponible en: https://unfhe.org/system/unfhe-65-Intersex_Factsheet_ENGLISH.pdf.
- Council of Europe. Commissioner for Human Rights. Human rights and intersex people. Issue Paper April 2015. (Consultado 7/1/2019). Disponible en: (Consultado 7/1/2019). Disponible en: <https://rmcoe.int/16806da5d4>.
- UN Committee against Torture, UN Committee on the Rights of the Child, UN Committee on the Rights of People with Disabilities, UN Subcommittee on Prevention of Torture and other Cruel, Inhuman or Degrading Treatment or Punishment, Méndez J, Puras D; Šimonovič D; Santos M, African Commission on Human and Peoples' Rights, Council of Europe Commissioner for Human Rights, Inter-American Commission on Human Rights. Intersex Awareness Day – Wednesday 26 October. End violence and harmful medical practices on intersex children and adults, UN and regional experts urge. Office of the High Commissioner for Human Rights, 2016. Consultado 14/1/2019). Disponible en: <https://www.ohchr.org/EN/NewsEvents/Pages/DisplayNews.aspx?NewsID=20739&LangID=E>
- European Parliament, Council of the European Union. Regulation (EU) 2016/679 of the European Parliament and the Council of 27 April 2016 on the protection of natural persons with regard to the processing of personal data and the free movement of such data, thus repealing Directive 95/46/EC (General Data Protection Regulation). *Journal of the European Union*, 4-5-2016: 119/5-119/88.
- Swiss National Advisory Commission on Biomedical Ethics NEK-CNE On the management of differences of sex development. Ethical issues relating to "intersexuality". 2012. Consultado 7/2/2019). Disponible en: https://web.archive.org/web/20150423213245/http://www.nekcne.ch/fileadmin/nek-cnedateien/Themen/Stellungnahmen/en/NEK_Intersexualitaet_En.pdf
- Carpenter Morgan. Intersex and ageing. International Human Rights Australia. 2015. Consultado 7/1/2019). Disponible en: <https://ihra.org.au/28385/intersex-and-ageing/>.
- World Health Organization (2015). Sexual health, human rights and the law. Geneva: World Health Organization Berna, Switzerland. Consultado 7/1/2019). Disponible en: https://apps.who.int/iris/bitstream/handle/10665/175556/9789241564984_eng.pdf;jsessionid=5F48D6D9CE4CDA961433E6E89A5F6EB9?sequence=1
- Guerrero-Fernández J, Azcona San Juliána C, Barreiro Condea J, Bermúdez de la Vega JA, Carcavilla Urquía A, Castaño González LA, et al. Guía de actuación en las anomalías de la diferenciación sexual (ADS) / desarrollo sexual diferente (DSD). *An Pediatr (Barc)*. 2018;89(5):315.e1-315.e19.
- Bermon S, Garnier PY, Hirschberg AL, Robinson N, Giraud S, Nicolí R, et al. Serum androgen levels in elite female athletes. *J Clin Endocrinol Metab*. 2014;99:4328-35.
- Handelsman DJ, Hirschberg AL, Bermon S. Circulating testosterone as the hormonal basis of sex differences in athletic performance. *Endocr Rev* 2018;39:803-29.
- Auchus RJ. Endocrinology and women's sports: the diagnosis matters. *Law & Contemp Probs*. 2017;4:127.
- Allen DB. Hormonal eligibility criteria for 'includes females' competition: A practical but problematic solution. *Horm Res Paediatr*. 2016;85:278-82.
- Bermon S, Vilain E, Fénichel P, Ritzén M Women with hyperandrogenism in elite sports: scientific and ethical rationales for regulating. *J Clin Endocrinol Metab*. 2015;100:828-30.
- Ritzén M, Ljungqvist A, Budgett R, Garnier PY, Bermon S, Lindén-Hirschberg A, Vilain E, Martínez-Patiño MJ. The regulations about eligibility for women with hyperandrogenism to compete in women's category are well founded. A rebuttal to the conclusions by Healy et al. *Clin Endocrinol (Oxf)*. 2015;82:307-8.
- Sánchez FJ, Martínez-Patiño MJ, Vilain E. The new policy on hyperandrogenism in elite female athletes is not about "sex testing". *J Sex Res*. 2013;50:112-5.
- Wood RI, Stanton SJ. Testosterone and sport: current perspectives. *Horm Behav* 2012;61:147-55.
- Ballantyne KN, Kayser M, Grootegoed JA. Sex and gender issues in competitive sports: investigation of a historical case leads to a new viewpoint. *Br J Sports Med*. 2012;46:614-7.
- Gooren L. The significance of testosterone for fair participation of the female sex in competitive sports. *Asian J Androl*. 2011;13:653-4.
- Hercher L. Gender verification: a term whose time has come and gone. *J Genet Couns* 2010;19:551-3.

31. Handelsman DJ, Gooren LJ. Hormones and sport: physiology, pharmacology and forensic science. *Asian J Androl*. 2008;10:348-50.
32. Hipkin LJ. The XY female in sport: the controversy continues. *Br J Sports Med*. 1993;27:150-6.
33. Healy ML, Gibney J, Pentecost C, Wheeler MJ, Sonksen PH. Endocrine profiles in 693 elite athletes in the postcompetition setting. *Clin Endocrinol (Oxf)*. 2014;81:294-305.
34. Sonksen P, Ferguson-Smith MA, Bavington LD, Holt RI, Cowan DA, Catlin DH, Kidd B, Davis G, Davis P, Edwards L, Tamar-Mattis A. Medical and ethical concerns regarding women with hyperandrogenism and elite sport. *J Clin Endocrinol Metab*. 2015;100:825-7.
35. Huang G, Basaria S. Do anabolic-androgenic steroids have performance-enhancing effects in female athletes? *Mol Cell Endocrinol*. 2018;464:56-64.
36. World Antidoping Agency. La Lista de Prohibiciones. El Código Mundial Antidopaje, Estándar Internacional. Enero de 2018. Consultado 7/1/2019). Disponible en: https://www.wadaama.org/sites/default/files/prohibited_list_2018_sp.pdf.
37. Berman S. Androgens and athletic performance of elite female athletes. *Curr Opin Endocrinol Diabetes Obes*. 2017;24:246-51.
38. Boucharde C, Rankinen T, Timmons JA. Genomics and genetics in the biology of adaptation to exercise. *Compr Physiol*. 2011;1:1603-48.
39. Kiss MAPDM, Böhme MTS, Mansoldo AC, Degaki E, Regazzini M. Performance and sports talent. *Rev Paul Educ Fis*. 2004;19:89-100.
40. Gibson WT. Key concepts in human genetics: understanding the complex phenotype. *Med Sport Sci*. 2009; 54:1-10.
41. Tucker R, Collins M What makes champions? A review of the relative contribution of genes and training to sporting success. *Br J Sports Med* 2012; 46:555-61.
42. Eynon N, Ruiz JR, Oliveira J, Duarte JA, Birk R, Lucia A. Genes and elite athletes: a roadmap for future research. *J Physiol*. 2011; 589:3063-70.
43. Lunde O, Magnus P, Sandvik L, Høglø S. Familial clustering in the Polycystic Ovarian syndrome. *Gynecol Obstet Invest*. 1989; 28:23-30.
44. Fratantonio E, Vicari E, Pafumi C, Calogero AE. Genetics of polycystic ovarian syndrome. *Reprod Biomed Online*. 2009; 10:713-20.
45. Barber TM, Franks S. Genetics of polycystic ovary syndrome. *Front Hormone Res*. 2013;40:28-39.
46. Carey AH, Chan KL, Short F, White D, Williamson R, Franks S. Evidence for a single gene effect causing polycystic ovaries and male pattern baldness. *Clin Endocrinol*. 1993; 38:653-8.
47. Legro RS. Is there a male phenotype in polycystic ovary syndrome families? *J Pediatr Endocrinol Metab*. 2000; 13:1307-9.
48. Dusková M, Cermáková I, Hill M, Vanková M, Sámáliková P, Stárka L. What may be the markers of the male equivalent of polycystic ovary syndrome? *Physiol Res*. 2004; 53:287-95.
49. Stárka L, Hill M, Poláček V. Hormonal profile of men with premature androgenetic alopecia. *Sbornik Lékařsky*. 2000; 101:17-22.
50. Cannarella R, Condorelli RA, Mongioì LM, La Vignera S, Calogero AE Does a male polycystic ovarian syndrome equivalent exist? *J Endoc Invest*. 2018; 41:49-57.
51. Agencia Mundial Antidopaje. Código Mundial Antidopaje 2015. Consultado 7/1/2019). Disponible en: <https://aepsad.culturaydeporte.gob.es/dam/jcr:eb761b8c-17a1-4f7f-b20664828c4cd86e/codigomundialantidopaje2015.PDF>.
52. Consejo General de Colegios Oficiales de Médicos. Código de Deontología Médica. Guía de Ética Médica. Madrid.2011.
53. The 2017 hormone therapy position statement of The North American Menopause Society. The NAMS 2017 Hormone Therapy Position Statement Advisory Panel. Collaborators: Pinkerton JV, Sánchez Aguirre F, Blake J, Cosman F, Hodis HN, Hoffstetter S, Kaunitz AM, Kingsberg SA, Maki PM, Manson JE, Marchbanks P, McClung MR, Nachtigall LE, Nelson LM, PACE DT, Reid RL, Sarrel PM, Shifren JL, Stuenkel CA, Utian WH. *Menopause*. 2017;24:728-53.

VIII JORNADAS NACIONALES DE MEDICINA DEL DEPORTE

MEDICINA DEL BALONCESTO

22-23 DE NOVIEMBRE DE 2019



SOCIEDAD ESPAÑOLA DE MEDICINA DEL DEPORTE (SEMED)
REGIDORÍA D'ESPORTS / AJUNTAMENT DE REUS

COMITÉ ORGANIZADOR

Presidente:	Pedro Manonelles Marqueta
Vicepresidente y Pte Comité Organizador local:	Luis Franco Bonafonte
Secretario General:	Francisco Javier Rubio Pérez
Tesorero:	Javier Pérez Ansón
Vocales:	Daniel Brotons Cuixart Antoni Castro Salomó Juan N. García-Nieto Portabella Teresa Gaztañaga Aurrekoetxea Mateu Huguet Recasens José Fernando Jiménez Díaz Carlos Miñarro García María Pilar Oyón Belaza Salvador Sarrá Moretó Paola Ugarte Peyrón

COMITÉ CIENTÍFICO

Presidente:	Miguel Del Valle Soto
Secretario:	Oriol Abellan Aynés
Vocales:	Gonzalo Correa González Carlos De Teresa Galván Emilio Luengo Fernández Lluís Masana Marín Zigor Montalvo Zenarruzabeitia Manuel Montero Jaime José Naranjo Orellana Eduardo Ortega Rincón Jordi Salas Salvado Ángel Sánchez Ramos José Luis Terreros Blanco

PROGRAMA CIENTÍFICO (PRELIMINAR)

DÍA 22 DE NOVIEMBRE, VIERNES

- 09.00-10.30 PONENCIA: La Medicina del Deporte en el Baloncesto.**
Moderador: **Francisco Javier Rubio Pérez**
Baloncesto femenino. **Silvia Treviño Monjas**
Organización y control médico en Selecciones Españolas. **Pilar Doñoro Cuevas**
Baloncesto en la discapacidad – baloncesto en silla de ruedas. **Josep Oriol Martínez Ferrer**
- 11.00 -12.30 PONENCIA: Lesiones y Baloncesto**
Moderador: **Alfredo Rodríguez Gangoso**
La rodilla. **Jaume Perramon Llavina**
El tobillo. **Cristóbal Rodríguez Hernández**
Músculo y tendón. **Javier Valle López**
- 12.30 -13.30 CONFERENCIA INAUGURAL**
Presentación: **Luis Franco Bonafonte**
La historia del dopaje en el deporte olímpico
Eduardo Henrique De Rose
- 15.30 -17.00 PONENCIA: Muerte Súbita y Deporte**
Moderador: **J. María Alegret Colomé**
Recomendaciones sobre participación deportiva en la cardiopatía isquémica. **Mats Borjesson**
El electrocardiograma en la prevención de la muerte súbita del deportista. **Gonzalo Grazioli**
Arritmias y muerte súbita del deportista.
Xavier Viñolas Prat
- 17.30 -19.00 TALLER**
Electrocardiograma en deportistas.
Emilio Luengo Fernández

DÍA 23 DE NOVIEMBRE, SÁBADO

- 10.00 -11.30 PONENCIA: Controversias: Nutrición - Ayudas Ergogénicas. Los mitos de la alimentación en el deporte.**
Moderador: **Mónica Bulló**
¿Influye el tipo de dieta en la microbiota y el rendimiento deportivo?
Teresa Gaztañaga Aurrekoetxea
Dietas detox y antioxidantes alimentarios en la práctica deportiva. **Nuria Rosique**
Ayudas ergogénicas, realidad o mito.
Begoña Manuz González
- 12.00 – 13.00 PONENCIA: Manejo del dolor en Medicina del Deporte.**
Moderador: **Isabel Tello Galindo**
Bloqueos nerviosos en lesiones del aparato locomotor en Medicina del Deporte.
Eduardo Marco Sánchez
Distrofia Simpático Refleja y Lumbalgia – Síndrome facetario en deportistas. ¿Qué ofrece la Unidad de Dolor? **Guillem Bujosa Portells**
- 13.00 -13.45 CONFERENCIA DE CLAUSURA**
Presentación: **Pedro Manonelles Marqueta**
Actualización en dopaje. **José Luis Terreros Blanco**

COMUNICACIONES CIENTIFICAS

El Comité Científico invita a todos los participantes a remitir comunicaciones científicas (comunicaciones orales y póster-presentación interactiva) a las VIII Jornadas Nacionales de la Sociedad Española de Medicina del Deporte.

Temas para presentación de Comunicaciones Científicas:

- Medicina del deporte.
- Entrenamiento y mejora del rendimiento.
- Biomecánica.
- Cardiología del deporte.
- Fisiología del esfuerzo.
- Nutrición y ayudas ergogénicas.
- Cineantropometría.
- Lesiones deportivas: diagnóstico, prevención y tratamiento.
- Actividad física y salud.

INFORMACIÓN GENERAL

22-23 de noviembre de 2019

Lugar


Auditorio y aulas
Hospital Universitari Sant Joan de Reus
Av. del Dr. Josep Laporte, 1
43204 – Reus (Tarragona)
Tfno: 977310300
Unidad de Medicina del Deporte.
Tfno: 977308305
Fax: 977337753
Correo electrónico: lfranco@grupsagessa.com
Localización del hospital: <http://www.hospitalsantjoan.cat/contacteu/>

Secretaría Científica

Sociedad Española de Medicina del Deporte
Dirección: Apartado de correos 1207. 31080 Pamplona
Teléfono: +34 948 26 77 06
Fax: +34 948 17 14 31
Correo electrónico: congresos@femede.es
<http://www.femede.es/page.php?/interno/OtrasActividades>

Secretaría Técnica


Viajes El Corte Inglés S.A.
División Eventos Deportivos
C/ Tarifa, nº 8. 41002 Sevilla
Teléfono: + 34 954 50 66 23
Correo electrónico: areaeventos@viajeseci.es
Personas de contacto: Silvia Herreros



Derechos de inscripción	Antes del 18-7-2019	Del 18-7-2019 al 19-9-2019	Desde 27-9-19 y en sede Jornadas
Cuota general	125 euros	150 euros	200 euros
Miembros ARAMEDE/ FEMEDE	100 euros	125 euros	175 euros
Médicos MIR*	60 euros	75 euros	125 euros
Estudiantes**	30 euros	30 euros	30 euros

*Es necesaria acreditación.

**Grados, Licenciaturas y Diplomaturas: Medicina, CC Actividad Física y Deporte, CC de la Salud...). Es necesaria acreditación. No se considera estudiantes los profesionales que cursen estudios, ni a graduados, licenciados y/o diplomados.



PERFECCIONA TU ENTRENAMIENTO

Por: Nick Tumminello
 Edita: Ediciones Tutor. Editorial El Drac.
 Impresores 20. P.E. Prado del Espino. 28660 Boadilla del Monte. Madrid.
 Telf. 915 599 832 - Fax: 915 410 235
 E-mail: info@edicionestutor.com Web: www.edicionestutor.com
 Madrid 2019. 320 páginas. P.V.P: 29,95 euros

Cada sesión de entrenamiento pretende solo una cosa: maximizar los resultados. Tanto si los resultados son la forma física, la función y el rendimiento, la pérdida de grasa o mejorar el físico, tu mayor deseo es que el esfuerzo merezca la pena.

Esta obra contiene 243 ejercicios y 71 programas, e incluso sesiones

alternativas para desarrollarlas en casa o en el gimnasio de un hotel y en las que se utiliza el peso corporal. Además, el lector podrá aprender: los errores que pudiera estar cometiendo; pequeños cambios en la técnica que pueden producir grandes resultados; por qué ciertos ejercicios son preferibles a otros; si el hombre y la mujer deben entrenar

de modo diferente; ejercicios que han de evitarse y principios probados que han de seguirse; y combinaciones y secuencias que maximizan los resultados. Altamente didáctico y de aplicación inmediata, este libro ayudará al deportista a ajustar el progreso en la forma y acondicionamientos físicos.



CÓMO RESPIRAR PARA MEJORAR EL RENDIMIENTO

Por: Eric Franklin
 Edita: Ediciones Tutor. Editorial El Drac.
 Impresores 20. P.E. Prado del Espino. 28660 Boadilla del Monte. Madrid.
 Telf. 915 599 832 - Fax: 915 410 235
 E-mail: info@edicionestutor.com Web: www.edicionestutor.com
 Madrid 2019. 72 páginas. P.V.P: 15 euros

El maestro internacional Eric Franklin enseña, a través de las páginas de este libro, a desarrollar más energía y atención y a mejorar el rendimiento físico y deportivo optimizando algo que solemos dar por hecho: la respiración. Respirar es necesario para producir energía; una respiración sumamente funcional abre oportunidades para alcanzar un rendimiento máximo. La presente obra es un texto conciso y ampliamente ilustrado que ayudará al lector a aprender a: mejorar su función respiratoria para beneficiar su salud e incrementar el rendimiento

deportivo; comprender la anatomía de la respiración, todos los músculos implicados y cómo interactúan entre sí; practicar 35 ejercicios respiratorios para mejorar la técnica respiratoria funcional; estudiar y entrenar el músculo vital de la respiración, el diafragma; comprender la función y el movimiento de la caja torácica y su relación con la respiración; e integrar todos los elementos involucrados en la respiración para conseguir una óptima función respiratoria.

Durante más de 30 años de enseñanza, el autor ha puesto a prueba

las ideas y ejercicios presentados en este libro. Bailarines, practicantes de yoga, monitores de pilates, actores, logopedas, profesores de canto, fisioterapeutas, nadadores, corredores y otros muchos profesionales han utilizado sus principios. Su descripción detallada de la anatomía implicada, su presentación de los ejercicios y su capacidad de integrar esta información de manera práctica convierten este texto en una referencia importante para quienes persiguen alcanzar su máximo rendimiento en el deporte y en la vida.



MUERTE SÚBITA EN EL DEPORTE. MÉTODOS DE CRIBADO

Por: E. Luengo, P. Manonelles
 Edita: Esmon Publicidad S.A. Sociedad Española de Medicina del Deporte. Apartado 1207. 31080 Pamplona.
 Telf. 948 267 706 - Fax: 948 171 431
 Email: femedede@femedede.es Web: www.femedede.es
 Barcelona, 2016. 72 páginas. P.V.P: 25 euros

La muerte súbita es un hecho fatal en la práctica deportiva, especialmente por la trascendencia que tiene, tanto para el que la sufre como por la repercusión que tiene. Su familia, sus compañeros de actividad física, los

practicantes de su mismo deporte, la sociedad en general, los dirigentes deportivos y las autoridades, todos quedan afectados.

La muerte súbita es un episodio frecuentemente cardiovascular. Este

manual está orientado a apoyar la decisión de los médicos que harán el reconocimiento a los deportistas, para detectar las anomalías cardiovasculares más frecuentes que pueden desembocar en ese fatal acontecimiento.

2019		
The International Conference on Sport, Education & Psychology	2-3 Mayo Bucarest (Rumanía)	web: www.futureacademy.org.uk
International Conference on Medicine and Science in Athletics	3-5 Mayo Doha (Qatar)	web: www.aspetar.com
1er Congreso Internacional de Podología Deportiva	10-11 Mayo Plasencia (Cáceres)	web: www.sepod.es
3rd International Conference Sport, Recreation, Health	10-11 Mayo Belgrado (Serbia)	E-mail: conference@vss.edu.rs
12th Biennial ISAKOS	12-16 Mayo Cancún (México)	web: www.isakos.com
57º Congreso SERMEF	15-18 Mayo Sevilla	web: http://congresoserfef.com/index.php
Segunda Conferencia Internacional de Medicina Deportiva de Serbia	23-25 Mayo Belgrado (Serbia)	web: www.sportsmedicineserbia-conference.rs/srb/home
VIII Congreso de la Sociedad Española de Tratamientos Médicos con Ondas de Choque (SETOC)	14-15 Junio Lorca (Murcia)	web: http://grafismoautoedicion.com/Setoc.html
22nd International Symposium on Adapted Physical Activity (ISAPA)	14-18 Junio Charlottesville (EE.UU.)	web: http://isapa2019.org
2019 AIESEP International Conference	19-22 Junio Nueva York (EE.UU.)	web: https://aiesep2019.adelphi.edu
Curso Nacional de Rehabilitación de deformidades del Raquis	20-21 Junio Barcelona	web: http://www.aulavhebron.net/aula/index.php?go=info_cursos&curso=223&idioma=es
30 Jornadas AEMB	20-22 Junio Madrid	web: http://aemeb.es/madrid-2019/
XL Juegos Mundiales de la Medicina-International Sports Medicine Symposium	22-29 Junio Budva (Montenegro)	web: http://www.medigames.com
VIII Congreso Iberoamericano de Nutrición	3-5 Julio Pamplona	web: http://www.academianutricionydietetica.org/congreso.php?id=7#
24th Annual Congress of the European College of Sport Science	3-6 Julio Praga (Rep. Checa)	E-mail: office@sport-science.org
II Congreso Mexicano de Medicina del Deporte	3-6 Julio Mérida-Yucatán (México)	web: https://comede.mx/
13th Congreso Mundial de la International Society of Physical and Rehabilitation Medicine	9-13 Julio Kobe (Japón)	web: http://www.isprm.org
2nd International Conference on Physical Education, Sports Medicine and Doping Studies	15-16 Julio Sídney (Australia)	web: https://sportsmedicine.conferenceseries.com/
15th European Congress of Sport and Exercise Psychology	15-20 Julio Münster (Alemania)	web: https://www.fepsac2019.eu

Agenda

Congreso colombiano de nutrición y dietética y II Internacional en alimentación y nutrición	15-17 Agosto Manizales (Colombia)	web: https://acodin.org/congreso-2019/
9th VISTA Conference	4-7 Septiembre Amsterdam (Países Bajos)	web: www.paralympic.org/news/amsterdam-host-vista-2019
Congress on Healthy and Active Children	11-14 Septiembre Verona (Italia)	web: http://i-mdrc.com/fourth-assembly/
14th International Congress of shoulder and elbow surgery (ICSES)	17-20 Septiembre Buenos Aires (Argentina)	web: www.icses2019.org
56º Congreso SECOT	25-27 Septiembre Zaragoza	web: www.secot.es
IX Congreso de la Sociedad Cubana de Medicina Física y Rehabilitación	1-4 Octubre La Habana (Cuba)	web: http://www.rehabilitacioncuba.com
11th European Congress on Sports Medicine	3-5 Octubre Portorose (Eslovenia)	web: http://www.efsma.eu
13th European Nutrition Conference On Malnutrition In An Obese World	13-18 Octubre Dublín (Irlanda)	web: www.fens2019.org
50 Congreso Nacional de Podología y VI Encuentro Iberoamericano	18-19 Octubre Santander	web: https://50congresopodologia.com/
Congreso Internacional de Fisioterapia	25-26 Octubre Toledo	web: congreso@coficam.org
5th World Conference on Doping in Sport	5-7 Noviembre Katowice (Polonia)	web: http://www.wada-ama.org
26th Word Congress TAFISA	13-17 Noviembre Tokyo (Japón)	web: www.tafisa.org
2019 FIP World Congress of Podiatry	14-16 Noviembre Miami (EEUU)	web: www.podiatry2019.org
VIII Jornadas Nacionales de Medicina del Deporte: "Medicina del Baloncesto"	22-23 Noviembre Reus (Tarragona)	E-mail: femede@femede.es web: www.femede.es
10th Annual International Conference: Physical Education Sport & Health	23-24 Noviembre Pitesti (Rumanía)	web: http://sportconference.ro/
56 Congreso Argentino de COT	28 Noviembre-1 Diciembre Buenos Aires (Argentina)	web: www.congresoaaot.org.ar
2020		
14th ISPRM World Congress – ISPRM 2020	4-9 Marzo Orlando (EE.UU.)	web: http://www.isprm.org/congress/14th-isprm-world-congress
Congreso FESNAD	11-13 Marzo Zaragoza	web: http://www.fesnad.org/
IOC World Conference Prevention of Injury & Illness in Sport	12-14 Marzo Mónaco (Principado de Mónaco)	web: http://ioc-preventionconference.org/
37º Congress International Society for Snowsports Medicine-SITEMSH	1-3 Abril Andorra la Vella (Principat d'Andorra)	E-mail: andorra2020@sitemnsh.org

9º Congrés Societat Catalana de Medicina de l'Esport-SCME	3-4 Abril Andorra la Vella (Principat d'Andorra)	E-mail: andorra2020@sitemnsh.org
25th Annual Congress of the European College of Sport Science	1-4 Julio Sevilla	E-mail: office@sport-science.org
International Congress of Dietetics	15-18 Septiembre Cape Town (Sudáfrica)	web: http://www.icda2020.com/
XXXVI Congreso Mundial de Medicina del Deporte	24-27 Septiembre Atenas (Grecia)	web: www.globalevents.gr
26th TAFISA World Congress	13-17 Noviembre Tokyo (Japón)	web: www.icsspe.org/sites/default/files/e9_TAFISA%20World%20Congress%202019_Flyer.pdf
VIII Congreso HISPAMEF	15-17 Octubre Cartagena de Indias (Colombia)	web: http://hispacef.com/viii-congreso-hispacef-15-17-de-2020/
XVIII Congreso Internacional SEMED-FEMEDE	Murcia	web: www.femede.es
2021		
Congreso Mundial de Psicología del Deporte	1-5 Julio Taipei (Taiwan)	web: https://www.issponline.org/index.php/events/next-world-congress
26th Annual Congress of the European College of Sport Science	7-10 Julio Glasgow (Reino Unido)	E-mail: office@sport-science.org
22nd International Congress of Nutrition (ICN)	14-19 Septiembre Tokyo (Japón)	web: http://icn2021.org/
European Federation of Sports Medicine Associations (EFSMA) Conference 2021	28-30 Octubre Budapest (Hungria)	web: http://efsma.eu/
Congreso Mundial de Psicología del Deporte	Taipei (Taiwan)	
Congreso Mundial de Podología	Barcelona	web: https://cgcop.es/newweb/eventos/
2022		
8th IWG World Conference on Women and Sport	5-8 Mayo Auckland (N. Zelanda)	web: http://iwgwomenandsport.org/world-conference/
XXXVII Congreso Mundial de Medicina del Deporte FIMS	Septiembre Guadalajara (México)	web: www.femmede.com.mx

Curso "ENTRENAMIENTO, RENDIMIENTO, PREVENCIÓN Y PATOLOGÍA DEL CICLISMO"

Curso dirigido a los titulados de las diferentes profesiones sanitarias y a los titulados en ciencias de la actividad física y el deporte, destinado al conocimiento de las prestaciones y rendimiento del deportista, para que cumpla con sus expectativas competitivas y de prolongación de su práctica deportiva, y para que la práctica deportiva minimice las consecuencias que puede tener para su salud, tanto desde el punto de vista médico como lesional.

Curso "ELECTROCARDIOGRAFÍA PARA MEDICINA DEL DEPORTE"

ACREDITADO POR LA COMISIÓN DE FORMACIÓN CONTINUADA (ON-LINE 1/5/2018 A 1/5/2019) CON 2,93 CRÉDITOS

Curso dirigido a médicos destinado a proporcionar los conocimientos específicos para el estudio del sistema cardiocirculatorio desde el punto de vista del electrocardiograma (ECG).

Curso "FISIOLOGÍA Y VALORACIÓN FUNCIONAL EN EL CICLISMO"

Curso dirigido a los titulados de las diferentes profesiones sanitarias y a los titulados en ciencias de la actividad física y el deporte, destinado al conocimiento profundo de los aspectos fisiológicos y de valoración funcional del ciclismo.

Curso "AYUDAS ERGOGÉNICAS"

Curso abierto a todos los interesados en el tema que quieren conocer las ayudas ergogénicas y su utilización en el deporte.

Curso "CARDIOLOGÍA DEL DEPORTE"

ACREDITADO POR LA COMISIÓN DE FORMACIÓN CONTINUADA (ON-LINE 1/5/2018 A 1/5/2019) CON 6,60 CRÉDITOS

Curso dirigido a médicos destinado a proporcionar los conocimientos específicos para el estudio del sistema cardiocirculatorio desde el punto de vista de la actividad física y deportiva, para diagnosticar los problemas cardiovasculares que pueden afectar al deportista, conocer la aptitud cardiológica para la práctica deportiva, realizar la prescripción de ejercicio y conocer y diagnosticar las enfermedades cardiovasculares susceptibles de provocar la muerte súbita del deportista y prevenir su aparición.

Curso "ALIMENTACIÓN, NUTRICIÓN E HIDRATACIÓN EN EL DEPORTE"

Curso dirigido a médicos destinado a facilitar al médico relacionado con la actividad física y el deporte la formación precisa para conocer los elementos necesarios para la obtención de los elementos energéticos necesarios para el esfuerzo físico y para prescribir una adecuada alimentación del deportista.

Curso "ALIMENTACIÓN Y NUTRICIÓN EN EL DEPORTE"

Curso dirigido a los titulados de las diferentes profesiones sanitarias (existe un curso específico para médicos) y para los titulados en ciencias de la actividad física y el deporte, dirigido a facilitar a los profesionales relacionados con la actividad física y el deporte la formación precisa para conocer los elementos necesarios para la obtención de los elementos energéticos necesarios para el esfuerzo físico y para conocer la adecuada alimentación del deportista.

Curso "ALIMENTACIÓN Y NUTRICIÓN EN EL DEPORTE" Para Diplomados y Graduados en Enfermería

ACREDITADO POR LA COMISIÓN DE FORMACIÓN CONTINUADA (NO PRESENCIAL 15/12/2015 A 15/12/2016) CON 10,18 CRÉDITOS

Curso dirigido a facilitar a los Diplomados y Graduados en Enfermería la formación precisa para conocer los elementos necesarios para la obtención de los elementos energéticos necesarios para el esfuerzo físico y para conocer la adecuada alimentación del deportista.

Curso "CINEANTROPOMETRÍA PARA SANITARIOS"

Curso dirigido a sanitarios destinado a adquirir los conocimientos necesarios para conocer los fundamentos de la cineantropometría (puntos anatómicos de referencia, material antropométrico, protocolo de medición, error de medición, composición corporal, somatotipo, proporcionalidad) y la relación entre la antropometría y el rendimiento deportivo.

Curso "CINEANTROPOMETRÍA"

Curso dirigido a todas aquellas personas interesadas en este campo en las Ciencias del Deporte y alumnos de último año de grado, destinado a adquirir los conocimientos necesarios para conocer los fundamentos de la cineantropometría (puntos anatómicos de referencia, material antropométrico, protocolo de medición, error de medición, composición corporal, somatotipo, proporcionalidad) y la relación entre la antropometría y el rendimiento deportivo.

Más información:
www.femede.es

Guidelines of publication Archives of Sports Medicine

The ARCHIVES OF SPORTS MEDICINE Journal (Arch Med Deporte) with ISSN 0212-8799 is the official publication of the Spanish Federation of Sports Medicine. This journal publishes original works about all the features related to Medicine and Sports Sciences from 1984. This title has been working uninterruptedly with a frequency of three months until 1995 and two months after this date. Arch Med Deporte works fundamentally with the system of external review carried out by two experts (peer review). It includes regularly articles about clinical or basic research, reviews, articles or publishing commentaries, brief communications and letters to the publisher. The articles may be published in both SPANISH and ENGLISH. The submission of papers in English writing will be particularly valued.

Occasionally oral communications accepted for presentation in the Federation's Congresses will be published.

The Editorial papers will only be published after an Editor requirement.

The manuscripts accepted for publication will become FEMEDE property and their reproduction, total or partial, must be properly authorized. All the authors will have to send a written letter conceding these rights as soon as the article is accepted for publication.

Submit of manuscripts

1. The papers must be submitted at the Editor in Chief's attention, written in double space in a DIN A4 sheet and numbered in the top right corner. It is recommended to use Word format, Times New Roman and font size 12. They must be sent by e-mail to FEMEDE's e-mail address: femede@femede.es.
2. On the first page exclusively it should include: title (Spanish and English), authors' first name, initial of the second name (if applicable), surname and optionally the second one; Main official and academic qualifications, workplace, full address and corresponding author e-mail. Supports received in order to accomplish the study – such as grants, equipments, medicaments, etc- have to be included. A letter in which the first author on behalf of all signatories of the study, the assignment of the rights for total or partial reproduction of the article, once accepted for publication shall be attached. Furthermore, the main author will propose up to four reviewers to the editor. According to the reviewers, at least one must be from a different nationality than the main author. Reviewers from the same institutions as the authors, will not be accepted.

3. On the second page the abstract of the work will appear both in Spanish and English, and will have an extension of 250-300 words. It will include the intention of the work (aims of the research), methodology, the most out-standing results and the main conclusions. It must be written in such a way to allow the understanding of the essence of the article without reading it completely or partially. After the abstract, from three to ten key words will be specified in Spanish and English, derived from the Medical Subject Headings (MeSH) of the National Library of Medicine (available in: <http://www.nlm.nih.gov/mesh/MBrowser.html>).
4. The extension of the text will change according to the section applicable:
 - a. Original research: maximum 5.000 words, 6 figures and 6 tables.
 - b. Review articles: maximum 5.000 words, 5 figures and 4 tables. In case of needing a wider extension it is recommended to contact the journal Editor.
 - c. Editorials: they will be written by Editorial Board request.
 - d. Letters to the Editor: maximum 1.000 words.
5. Structure of the text: it will change according to the section applicable:
 - a. **ORIGINALS RESEARCH:** It will contain an introduction, which must be brief and will contain the aim of the work, written in such a way that the reader can understand the following text.
Material and method: the material used in the work will be exposed, as well as its characteristics, selection criteria and used techniques, facilitating the necessary data in order to allow the reader to repeat the experience shown. The statistical methods will be detailed described.
Results: Results must report and not describe the observations made with the material and method used. This information can be published in detail in the text or using tables and figures. Information given in the tables or figures must not be repeated in the text.
Discussion: The authors will expose their opinions about the results, their possible interpretation, relating the observations to the results obtained by other authors in similar publications, suggestions for future works on the topic, etc. Connect the conclusions with the aims of the study, avoiding free affirmations and conclusions not supported by the information of the work.
The acknowledgments will appear at the end of the text.

- b. **REVIEWS ARTICLES:** The text will be divided in as much paragraphs as the author considers necessary for a perfect comprehension of the topic treated.
- c. **LETTERS TO THE EDITOR:** Discussion about published papers in the last two issues, with the contribution of opinions and experiences briefed in 3 pages, will have preference in this Section.
- d. **OTHERS:** Specific sections commissioned by the Journal's Editorial Board.
6. **Bibliography:** it will be presented on pages apart and will be ordered following their appearance in the text, with a correlative numeration. In the text the quote's number will be presented between parentheses, followed or not by the authors' name; if they are mentioned, in case the work was made by two authors both of them will figure, and if there are more than two authors only the first will figure, followed by "et al".

There will not be personal communication, manuscripts or any unpublished information included in the bibliographical appointments.

The official citation for the journal Archives of Sports Medicine is Arch Med Sport.

References will be exposed in the following way:

- **Journal: order number;** surnames and name's initial of the article authors with no punctuation and separated with a comma (if the number of authors is higher than six, only the six first will figure, followed by "et al"); work's title in its original language; abbreviated journal name, according to the World Medical Periodical; year of publication; volume number; first and last page of the quoted extract. Example: Calbet JA, Radegran G, Boushel R and Saltin B. On the mechanisms that limit oxygen uptake during exercise in acute and chronic hypoxia: role of muscle mass. *J Physiol.* 2009;587:477-90.
 - **Book chapter:** Authors, chapter title, editors, book title, city, publishing house, year and number of pages. Example: Iselin E. Maladie de Kienbock et Syndrome du canal carpien. En : Simon L, Alieu Y. Poignet et Medecine de Reededucation. Londres : Collection de Pathologie Locomotrice Masson; 1981. p162-6.
 - **Book.** Authors, title, city, publishing house, year of publication, page of the quote. Example: Balius R. Ecografía muscular de la extremidad inferior. Sistemática de exploración y lesiones en el deporte. Barcelona. Editorial Masson; 2005. p 34.
 - **World Wide Web,** online journal. Example: Morse SS. Factors in the emergence of infectious diseases. *Emerg Infect Dis* (revista electrónica) 1995 JanMar (consultado 0501/2004). Available in: <http://www.cdc.gov/ncidod/EID/eid.htm>
7. **Tables and figures.** Tables and figures will be sent on separate files in JPEG format. Tables must be sent in word format.

Tables shall be numbered according to the order of appearance in the text, with the title on the top and the abbreviations described on the bottom. All nonstandard abbreviations which may be used in the tables shall be explained in footnotes.

Any kind of graphics, pictures and photographs will be denominated figures. They must be numbered correlatively by order of appearance in the text and will be sent in black and white (except in those works in which colour is justified). Color printing is an economic cost that has to be consulted with the editor.

All tables as well as figures will be numbered with Arabic numbers following the order of appearance in the text.

At the end of the text document the tables and figures captions will be included on individual pages.

8. The Journal's Editorial Staff will communicate the reception of submitted articles and will inform about its acceptance and possible date of publication.
9. After hearing the reviewers' suggestions (journal uses peer correction system), may reject the works which are not suitable, or indicate the author the modifications which are thought to be necessary for its acceptance.
10. The Editorial Board is not responsible for the concepts, opinions or affirmations supported by the authors.
11. Submissions of the papers: Archives of Sports Medicine. By e-mail to FEMEDE'S e-mail address: femedede@femedede.es. The submission will come with a cover letter on which the work's examination for its publication in the Journal will be requested, article type will be specified, and it will be certified by all authors that the work is original and has not been partially or totally published before.

Conflicts of interests

If there should be any relation between the work's authors and any public or private entity, from which a conflict of interests could appear, it must be communicated to the Editor. Authors must fulfil a specific document.

Ethics

All authors that sign the articles accept the responsibility defined by the World Association of Medical Editors.

The papers sent to the journal for evaluation must have been elaborated respecting the international recommendations about clinical and laboratory animals' researches, ratified in Helsinki and updated in 2008 by the American Physiology.

For the performance of controlled clinic essays the CONSORT normative shall be followed, available at <http://www.consort-statement.org/>

Campaña de aptitud física, deporte y salud



La **Sociedad Española de Medicina del Deporte**, en su incesante labor de expansión y consolidación de la Medicina del Deporte y, consciente de su vocación médica de preservar la salud de todas las personas, viene realizando diversas actuaciones en este ámbito desde los últimos años.

Se ha considerado el momento oportuno de lanzar la campaña de gran alcance, denominada **CAMPAÑA DE APTITUD FÍSICA, DEPORTE Y SALUD** relacionada con la promoción de la actividad física y deportiva para toda la población y que tendrá como lema **SALUD – DEPORTE – DISFRÚTALOS**, que aúna de la forma más clara y directa los tres pilares que se promueven desde la Medicina del Deporte que son el practicar deporte, con objetivos de salud y para la mejora de la aptitud física y de tal forma que se incorpore como un hábito permanente, y disfrutando, es la mejor manera de conseguirlo.



UCAM Universidad Católica San Antonio de Murcia

Campus de los Jerónimos,
Nº 135 Guadalupe 30107

(Murcia) - España

Tlf: (+34)968 27 88 01 · info@ucam.edu



UCAM
UNIVERSIDAD
CATÓLICA DE MURCIA